

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(3-Chlorobenzoyl)-2-nitrobenzene-sulfonamide

P. A. Suchetan,^a Sabine Foro,^b B. Thimme Gowda^{a*} and M. Shet Prakash^c

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany, and ^cDepartment of Chemistry, University College of Science, Tumkur University, Tumkur 572 102, India
Correspondence e-mail: gowdabt@yahoo.com


Received 8 February 2012; accepted 13 February 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.074; wR factor = 0.147; data-to-parameter ratio = 14.3.

In the title compound, $\text{C}_{13}\text{H}_9\text{ClN}_2\text{O}_5\text{S}$, the N—C bond in the C—SO₂—NH—C segment has a *gauche* torsion with respect to the S=O bonds. The conformation between the N—H bond and the *ortho*-nitro group in the sulfonyl benzene ring is *syn*, and that between the C=O and the *meta*-Cl atom in the benzoyl ring is *anti*. The molecule is twisted at the S—N bond, with a torsion angle of 65.41 (38)°. The dihedral angle between the sulfonyl benzene ring and the —SO₂—NH—C—O segment is 75.0 (1)°, and that between the sulfonyl and the benzoyl benzene ring is 89.1 (1)°. The crystal structure features inversion-related dimers linked by pairs of N—H···O(S) hydrogen bonds.

Related literature

For our studies of the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Gowda *et al.* (1999, 2006); *N*-(aryl)-methanesulfonamides, see: Gowda *et al.* (2007); *N*-(substitutedbenzoyl)-arylsulfonamides, see: Suchetan *et al.* (2012); *N*-chloroarylamides, see: Jyothi & Gowda (2004) and *N*-bromoarylsulfonamides, see: Usha & Gowda (2006)..



Experimental

Crystal data

$\text{C}_{13}\text{H}_9\text{ClN}_2\text{O}_5\text{S}$
 $M_r = 340.73$
 Orthorhombic, *Pbca*
 $a = 12.2046$ (8) Å
 $b = 12.6121$ (9) Å
 $c = 18.433$ (1) Å
 $V = 2837.3$ (3) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.44$ mm⁻¹
 $T = 293$ K
 $0.28 \times 0.28 \times 0.08$ mm

Data collection

Oxford Xcalibur diffractometer with Sapphire CCD detector
 Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2009)
 $T_{\min} = 0.886$, $T_{\max} = 0.966$
 11298 measured reflections
 2889 independent reflections
 1911 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.074$
 $wR(F^2) = 0.147$
 $S = 1.20$
 2889 reflections
 202 parameters
 1 restraint
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.29$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N···O1 ⁱ	0.86 (2)	2.41 (3)	3.193 (4)	153 (4)

Symmetry code: (i) $-x, -y + 2, -z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the University Grants Commission, Government of India, New Delhi, for a special grant under a UGC-BSR one-time grant to faculty.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2268).

References

- Gowda, B. T., Bhat, D. K., Fuess, H. & Weiss, A. (1999). *Z. Naturforsch. Teil A*, **54**, 261–267.
 Gowda, B. T., Foro, S. & Fuess, H. (2007). *Acta Cryst.* **E63**, o2597.
 Gowda, B. T., Kozisek, J. & Fuess, H. (2006). *Z. Naturforsch. Teil A*, **61**, 588–594.
 Jyothi, K. & Gowda, B. T. (2004). *Z. Naturforsch. Teil A*, **59**, 64–68.
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Suchetan, P. A., Foro, S. & Gowda, B. T. (2012). *Acta Cryst.* **E68**, o274.
 Usha, K. M. & Gowda, B. T. (2006). *J. Chem. Sci.* **118**, 351–359.

supporting information

Acta Cryst. (2012). E68, o786 [doi:10.1107/S160053681200640X]

***N*-(3-Chlorobenzoyl)-2-nitrobenzenesulfonamide**

P. A. Suchetan, Sabine Foro, B. Thimme Gowda and M. Shet Prakash

S1. Comment

As part of our studies on the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Gowda *et al.*, 1999, 2006), *N*-(aryl)-methanesulfonamides (Gowda *et al.*, 2007), *N*-(substitutedbenzoyl)-arylsulfonamides (Suchetan *et al.*, 2012), *N*-chloroarylsulfonamides (Jyothi & Gowda, 2004) and *N*-bromoarylsulfonamides (Usha & Gowda, 2006), in the present work, the crystal structure of *N*-(3-chlorobenzoyl)-2-nitrobenzenesulfonamide has been determined (Fig.1).

The conformation between the N—H and C=O bonds in the C—SO₂—NH—C(O) segment is *anti* and the N—C bond in the segment has *gauche* torsion with respect to the S=O bonds (Fig. 1), similar to that observed in *N*-(2-chlorobenzoyl)-2-nitrobenzenesulfonamide (I) (Suchetan *et al.*, 2012). In the title compound, the conformation between the N—H bond and the *ortho*-nitro group in the sulfonyl benzene ring is *syn*, similar to that observed in (I). Further, the conformation of the C=O is *anti* to the *meta*-Cl atom in the benzoyl ring, similar to that observed between the C=O and the *ortho*-Cl atom in (I).

The molecule is twisted at the S—N bond with the torsional angle of 65.41 (38)°, compared to the value of -59.68 (17)° in (I).

The dihedral angle between the sulfonyl benzene ring and the —SO₂—NH—C—O segment is 75.0 (1)°, compared to the value of 77.5 (1)° in (I). Furthermore, the dihedral angle between the sulfonyl and the benzoyl benzene rings is 89.1 (1)°, compared to the value of 71.2 (1)° in (I).

In the crystal structure two molecules each are linked by pairs of intermolecular N—H···O (S) hydrogen bonds into dimers that are located around centers of inversion (Fig. 2 and Table 1).


S2. Experimental

The title compound was prepared by refluxing a mixture of 3-chlorobenzoic acid (0.02 mole), 2-nitrobenzenesulfonamide (0.02 mole) and excess phosphorous oxychloride for 3 h on a water bath. The resultant mixture was cooled and poured into crushed ice. The solid, *N*-(3-chlorobenzoyl)-2-nitrobenzenesulfonamide, obtained was filtered, washed thoroughly with water and then dissolved in sodium bicarbonate solution. The compound was later reprecipitated by acidifying the filtered solution with dilute HCl. It was filtered, dried and recrystallized.


Rod like colourless single crystals of the title compound used in X-ray diffraction studies were obtained by slow evaporation of the solvent from its toluene solution at room temperature.

S3. Refinement

The H atom of the NH group was located in a difference map and later restrained to N—H = 0.86 (2) %A. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the U_{eq} of the parent atom).

**Figure 1**

Molecular structure of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

View of the inversion-related dimers linked by pairs of N—H···O(S) hydrogen bonds (Hydrogen bonding is shown as dashed lines)..

***N*-(3-Chlorobenzoyl)-2-nitrobenzenesulfonamide**

Crystal data

$C_{13}H_9ClN_2O_5S$

$M_r = 340.73$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 12.2046$ (8) Å

$b = 12.6121$ (9) Å

$c = 18.433$ (1) Å

$V = 2837.3$ (3) Å³

$Z = 8$

$F(000) = 1392$

$D_x = 1.595$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1818 reflections

$\theta = 2.6$ – 27.9°

$\mu = 0.44$ mm⁻¹

$T = 293$ K

Rod, colourless

$0.28 \times 0.28 \times 0.08$ mm

Data collection

Oxford Xcalibur

diffractometer with Sapphire CCD detector

Radiation source: fine-focus sealed tube

Graphite monochromator

Rotation method data acquisition using ω and φ scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2009)

$T_{\min} = 0.886$, $T_{\max} = 0.966$

11298 measured reflections
 2889 independent reflections
 1911 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

$\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.6^\circ$
 $h = -13 \rightarrow 15$
 $k = -15 \rightarrow 11$
 $l = -22 \rightarrow 23$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.074$
 $wR(F^2) = 0.147$
 $S = 1.20$
 2889 reflections
 202 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0362P)^2 + 4.7309P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.29 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. CrysAlis RED (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.0112 (3)	0.7252 (3)	0.0676 (2)	0.0347 (9)
C2	-0.0862 (3)	0.7030 (3)	0.0131 (2)	0.0402 (10)
C3	-0.1369 (4)	0.6063 (4)	0.0079 (3)	0.0503 (12)
H3	-0.1854	0.5924	-0.0298	0.060*
C4	-0.1152 (4)	0.5307 (4)	0.0590 (3)	0.0552 (13)
H4	-0.1498	0.4652	0.0563	0.066*
C5	-0.0427 (4)	0.5506 (4)	0.1143 (3)	0.0537 (13)
H5	-0.0286	0.4986	0.1488	0.064*
C6	0.0092 (4)	0.6469 (3)	0.1188 (2)	0.0450 (11)
H6	0.0582	0.6598	0.1564	0.054*
C7	-0.0490 (3)	0.9234 (3)	0.1845 (2)	0.0389 (10)
C8	-0.1239 (3)	1.0094 (3)	0.2089 (2)	0.0354 (9)
C9	-0.1284 (3)	1.1061 (3)	0.1738 (2)	0.0390 (10)
H9	-0.0812	1.1205	0.1354	0.047*
C10	-0.2036 (4)	1.1812 (3)	0.1960 (2)	0.0400 (10)
C11	-0.2730 (4)	1.1613 (4)	0.2528 (2)	0.0456 (11)
H11	-0.3240	1.2119	0.2672	0.055*
C12	-0.2663 (4)	1.0654 (4)	0.2884 (2)	0.0499 (12)

H12	-0.3127	1.0518	0.3273	0.060*
C13	-0.1921 (4)	0.9897 (3)	0.2670 (2)	0.0452 (11)
H13	-0.1879	0.9254	0.2915	0.054*
N1	-0.0153 (3)	0.9315 (3)	0.11212 (18)	0.0400 (9)
H1N	-0.050 (3)	0.966 (3)	0.0794 (18)	0.048*
N2	-0.1148 (3)	0.7825 (3)	-0.0428 (2)	0.0528 (10)
O1	0.0765 (2)	0.8844 (2)	-0.00038 (15)	0.0478 (8)
O2	0.1576 (2)	0.8265 (2)	0.11536 (16)	0.0514 (8)
O3	-0.0201 (3)	0.8509 (2)	0.22210 (16)	0.0538 (8)
O4	-0.1559 (3)	0.8640 (3)	-0.0218 (2)	0.0730 (11)
O5	-0.0955 (3)	0.7592 (3)	-0.10577 (18)	0.0747 (11)
Cl1	-0.20968 (12)	1.29932 (10)	0.14917 (7)	0.0697 (4)
S1	0.06425 (8)	0.84464 (8)	0.07160 (6)	0.0381 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.036 (2)	0.030 (2)	0.038 (2)	0.0025 (18)	0.0033 (19)	0.0001 (18)
C2	0.037 (2)	0.044 (3)	0.040 (2)	0.007 (2)	0.0002 (19)	0.011 (2)
C3	0.042 (3)	0.054 (3)	0.055 (3)	-0.004 (2)	-0.008 (2)	0.004 (2)
C4	0.063 (3)	0.037 (3)	0.066 (3)	-0.011 (2)	-0.001 (3)	0.007 (2)
C5	0.068 (3)	0.041 (3)	0.052 (3)	0.005 (2)	-0.005 (3)	0.013 (2)
C6	0.052 (3)	0.043 (3)	0.040 (2)	0.007 (2)	-0.010 (2)	0.003 (2)
C7	0.044 (3)	0.037 (2)	0.035 (2)	-0.004 (2)	0.0004 (19)	0.0011 (19)
C8	0.039 (2)	0.036 (2)	0.031 (2)	-0.003 (2)	0.0033 (18)	-0.0040 (17)
C9	0.043 (3)	0.041 (2)	0.033 (2)	-0.004 (2)	0.0097 (19)	-0.0018 (18)
C10	0.049 (3)	0.033 (2)	0.037 (2)	-0.004 (2)	0.001 (2)	-0.0022 (18)
C11	0.047 (3)	0.048 (3)	0.042 (2)	0.002 (2)	0.009 (2)	-0.013 (2)
C12	0.060 (3)	0.050 (3)	0.040 (2)	-0.008 (3)	0.018 (2)	-0.005 (2)
C13	0.061 (3)	0.037 (2)	0.037 (2)	-0.006 (2)	0.009 (2)	-0.0048 (19)
N1	0.048 (2)	0.040 (2)	0.0329 (19)	0.0065 (18)	0.0043 (16)	0.0036 (16)
N2	0.049 (2)	0.057 (3)	0.053 (3)	-0.004 (2)	-0.012 (2)	0.016 (2)
O1	0.0518 (19)	0.0507 (18)	0.0411 (16)	-0.0011 (15)	0.0137 (14)	0.0022 (14)
O2	0.0366 (17)	0.060 (2)	0.0577 (19)	0.0028 (16)	-0.0076 (15)	-0.0045 (16)
O3	0.068 (2)	0.0485 (19)	0.0450 (17)	0.0083 (18)	0.0041 (16)	0.0094 (16)
O4	0.090 (3)	0.052 (2)	0.077 (3)	0.018 (2)	-0.007 (2)	0.0162 (19)
O5	0.087 (3)	0.098 (3)	0.0394 (19)	-0.011 (2)	-0.0108 (19)	0.0131 (19)
Cl1	0.0894 (11)	0.0479 (7)	0.0718 (9)	0.0188 (7)	0.0221 (7)	0.0146 (6)
S1	0.0379 (6)	0.0399 (6)	0.0365 (5)	0.0018 (5)	0.0035 (5)	-0.0007 (5)

Geometric parameters (Å, °)

C1—C6	1.388 (5)	C8—C9	1.382 (5)
C1—C2	1.389 (6)	C9—C10	1.381 (6)
C1—S1	1.767 (4)	C9—H9	0.9300
C2—C3	1.371 (6)	C10—C11	1.369 (6)
C2—N2	1.479 (5)	C10—Cl1	1.723 (4)
C3—C4	1.366 (6)	C11—C12	1.378 (6)

C3—H3	0.9300	C11—H11	0.9300
C4—C5	1.373 (6)	C12—C13	1.374 (6)
C4—H4	0.9300	C12—H12	0.9300
C5—C6	1.372 (6)	C13—H13	0.9300
C5—H5	0.9300	N1—S1	1.643 (4)
C6—H6	0.9300	N1—H1N	0.857 (19)
C7—O3	1.201 (5)	N2—O4	1.207 (5)
C7—N1	1.399 (5)	N2—O5	1.221 (5)
C7—C8	1.489 (6)	O1—S1	1.426 (3)
C8—C13	1.379 (5)	O2—S1	1.414 (3)
C6—C1—C2	117.8 (4)	C8—C9—H9	120.2
C6—C1—S1	119.0 (3)	C11—C10—C9	120.8 (4)
C2—C1—S1	123.1 (3)	C11—C10—C11	121.0 (3)
C3—C2—C1	121.9 (4)	C9—C10—C11	118.2 (3)
C3—C2—N2	116.7 (4)	C10—C11—C12	119.2 (4)
C1—C2—N2	121.5 (4)	C10—C11—H11	120.4
C4—C3—C2	119.0 (4)	C12—C11—H11	120.4
C4—C3—H3	120.5	C13—C12—C11	120.8 (4)
C2—C3—H3	120.5	C13—C12—H12	119.6
C3—C4—C5	120.7 (4)	C11—C12—H12	119.6
C3—C4—H4	119.7	C12—C13—C8	119.7 (4)
C5—C4—H4	119.7	C12—C13—H13	120.2
C6—C5—C4	120.2 (4)	C8—C13—H13	120.2
C6—C5—H5	119.9	C7—N1—S1	123.9 (3)
C4—C5—H5	119.9	C7—N1—H1N	124 (3)
C5—C6—C1	120.4 (4)	S1—N1—H1N	108 (3)
C5—C6—H6	119.8	O4—N2—O5	126.1 (4)
C1—C6—H6	119.8	O4—N2—C2	116.9 (4)
O3—C7—N1	121.3 (4)	O5—N2—C2	117.0 (4)
O3—C7—C8	124.1 (4)	O2—S1—O1	120.19 (19)
N1—C7—C8	114.5 (4)	O2—S1—N1	108.94 (18)
C13—C8—C9	119.9 (4)	O1—S1—N1	104.51 (18)
C13—C8—C7	118.3 (4)	O2—S1—C1	107.76 (19)
C9—C8—C7	121.7 (4)	O1—S1—C1	108.41 (19)
C10—C9—C8	119.5 (4)	N1—S1—C1	106.23 (18)
C10—C9—H9	120.2		
C6—C1—C2—C3	-1.8 (6)	C11—C10—C11—C12	179.3 (3)
S1—C1—C2—C3	174.5 (3)	C10—C11—C12—C13	-0.7 (7)
C6—C1—C2—N2	178.6 (4)	C11—C12—C13—C8	-0.4 (7)
S1—C1—C2—N2	-5.0 (6)	C9—C8—C13—C12	1.7 (6)
C1—C2—C3—C4	1.8 (7)	C7—C8—C13—C12	-176.5 (4)
N2—C2—C3—C4	-178.7 (4)	O3—C7—N1—S1	0.6 (6)
C2—C3—C4—C5	-0.7 (7)	C8—C7—N1—S1	-178.1 (3)
C3—C4—C5—C6	-0.2 (7)	C3—C2—N2—O4	119.2 (5)
C4—C5—C6—C1	0.1 (7)	C1—C2—N2—O4	-61.2 (6)
C2—C1—C6—C5	0.9 (6)	C3—C2—N2—O5	-60.2 (6)

S1—C1—C6—C5	-175.6 (3)	C1—C2—N2—O5	119.4 (5)
O3—C7—C8—C13	-22.3 (6)	C7—N1—S1—O2	-50.4 (4)
N1—C7—C8—C13	156.3 (4)	C7—N1—S1—O1	179.9 (3)
O3—C7—C8—C9	159.5 (4)	C7—N1—S1—C1	65.4 (4)
N1—C7—C8—C9	-21.9 (6)	C6—C1—S1—O2	17.3 (4)
C13—C8—C9—C10	-1.7 (6)	C2—C1—S1—O2	-159.0 (3)
C7—C8—C9—C10	176.4 (4)	C6—C1—S1—O1	148.9 (3)
C8—C9—C10—C11	0.5 (6)	C2—C1—S1—O1	-27.4 (4)
C8—C9—C10—C11	-178.1 (3)	C6—C1—S1—N1	-99.3 (3)
C9—C10—C11—C12	0.7 (7)	C2—C1—S1—N1	84.4 (4)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1N...O1 ⁱ	0.86 (2)	2.41 (3)	3.193 (4)	153 (4)

Symmetry code: (i) $-x, -y+2, -z$.