

2,2'-[(2*S**,6*R**)-Piperidine-2,6-diyl]dipropan-2-ol

Guillaume Journot,^a Reinhard Neier^{a*} and Helen Stoeckli-Evans^b

^aInstitute of Chemistry, University of Neuchâtel, Avenue de Bellevaux 51, CH-2000 Neuchâtel, Switzerland, and ^bInstitute of Physics, University of Neuchâtel, Rue Emile-Argand 11, CH-2000 Neuchâtel, Switzerland

Correspondence e-mail: reinhard.neier@unine.ch


Received 4 February 2012; accepted 9 February 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.086; wR factor = 0.146; data-to-parameter ratio = 17.2.

In the title compound, $\text{C}_{11}\text{H}_{23}\text{NO}_2$, the piperidine ring has a chair conformation. The two hydroxy H atoms are disordered over two positions with fixed occupancy ratios of 0.57:0.43 and 0.63:0.37. In the molecule, there are two short $\text{N}-\text{H}\cdots\text{O}$ interactions. In the crystal, four symmetry-related molecules are linked by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds to form a cage-like arrangement, centered about the point of intersection of three twofold axes. These cages stack along the [100] direction.

Related literature

For literature on ligands of the pincer-type family, see: van Koten (1989); Albrecht & van Koten (2001). For metal complexes of such pincer ligands, see: Hofmeier & Schubert (2004); Li *et al.* (2007). For the synthesis of the starting material 2,2'-(pyridine-2,6-diyl)dipropan-2-ol, see: Klein *et al.* (2009). For an example of the transformation of bis-benzylic alcohols of 2,6-disubstituted pyridines, see: Klein *et al.* (2009). For the crystal structure of *cis*-(piperidine-2,6-diyl)dimethanol, see: Hartung *et al.* (2007).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{23}\text{NO}_2$
 $M_r = 201.30$
 Orthorhombic, *Fddd*

$a = 12.0713$ (9) Å
 $b = 23.4762$ (10) Å
 $c = 34.496$ (2) Å

$V = 9775.8$ (10) Å³
 $Z = 32$
 Mo $K\alpha$ radiation

$\mu = 0.07$ mm⁻¹
 $T = 173$ K
 $0.45 \times 0.45 \times 0.40$ mm

Data collection

Stoe IPDS 2 diffractometer
 Absorption correction: multi-scan
 (*MULScanABS* in *PLATON*;
 Spek, 2009)
 $T_{\min} = 0.911$, $T_{\max} = 1.000$

32226 measured reflections
 2319 independent reflections
 1499 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.135$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.086$
 $wR(F^2) = 0.146$
 $S = 1.16$
 2319 reflections
 135 parameters
 4 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.16$ e Å⁻³
 $\Delta\rho_{\min} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}'$	0.84 (3)	2.38 (3)	2.792 (3)	111 (2)
$\text{N1}-\text{H1}\cdots\text{O1}''$	0.84 (3)	2.43 (3)	2.814 (3)	109 (2)
$\text{O1}'-\text{H1A}\cdots\text{O1}''^i$	0.82	1.99	2.805 (3)	169
$\text{O1}'-\text{H1B}\cdots\text{O1}''^ii$	0.83	1.99	2.807 (4)	167
$\text{O1}''-\text{H1C}\cdots\text{O1}''^i$	0.82	2.00	2.805 (3)	171
$\text{O1}''-\text{H1D}\cdots\text{O1}''^iii$	0.83	2.03	2.762 (5)	148

Symmetry codes: (i) $x, -y + \frac{1}{4}, -z + \frac{1}{4}$; (ii) $-x + \frac{5}{4}, -y + \frac{1}{4}, z$; (iii) $-x + \frac{5}{4}, y, -z + \frac{1}{4}$.

Data collection: *X-Area* (Stoe & Cie, 2009); cell refinement: *X-Area*; data reduction: *X-RED32* (Stoe & Cie, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97*, *PLATON* and *publCIF* (Westrip, 2010).

HSE thanks the staff of the XRD Application Laboratory, CSEM, Neuchâtel, for access to the X-ray diffraction equipment.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2389).

References

- Albrecht, M. & van Koten, G. (2001). *Angew. Chem. Int. Ed.* **40**, 3750–3781.
 Hartung, J., Stapf, G. & Bergsträsser, U. (2007). *Acta Cryst.* **E63**, o2586–o2587.
 Hofmeier, H. & Schubert, U. S. (2004). *Chem. Soc. Rev.* **33**, 373–399.
 Klein, A., Elmas, S. & Butsch, K. (2009). *Eur. J. Inorg. Chem.* pp. 2271–2281.
 Koten, G. van (1989). *Pure Appl. Chem.* **61**, 1681–1694.
 Li, Y., Huffman, J. C. & Flood, A. H. (2007). *Chem. Commun.* pp. 2692–2694.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Stoe & Cie. (2009). *X-Area* and *X-RED32*. Stoe & Cie GmbH, Darmstadt, Germany.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o857 [doi:10.1107/S1600536812005879]

2,2'-[(2*S,6*R**)-Piperidine-2,6-diyl]dipropan-2-ol****Guillaume Journot, Reinhard Neier and Helen Stoeckli-Evans****S1. Comment**

Terpyridine and its derivatives are prototypical ligands of the pincer type family (Van Koten, 1989; Albrecht & van Koten, 2001). They have been widely used in coordination chemistry (Hofmeier & Schubert, 2004). The metal complexes obtained from pincer ligands are conformationally restricted and often thermodynamically highly stable (Hofmeier & Schubert, 2004; Li *et al.*, 2007). The bis-benzylic alcohols of 2,6-disubstituted pyridines belonging to this class of ligands can be easily transformed (Klein *et al.*, 2009).

The modification of these ligands by the hydrogenation of the pyridine ring installs chirality into the structure and increases the basicity and the strength of the ligand. In contrast to the 2,6-pyridinedicarboxylic acids and its derivatives, which have been extensively used, studies on the corresponding tridentate ONO-piperidine ligands containing the bis-alcohols have been very rare so far. The title compound (2) was prepared by the stereoselective *cis*-reduction of 2,2'-(pyridine-2,6-diyl)dipropan-2-ol (1). Herein we report on the synthesis and the crystal structure of the title compound, (2).

The molecular structure of the title molecule is illustrated in Fig. 1. The geometric parameters are very similar to those found for *cis*-(piperidine-2,6-diyl)dimethanol (Hartung *et al.*, 2007). The piperidine ring has a chair conformation, with atoms N1 and C4 being displaced from the plane through atoms C2/C3/C5/C6 by 0.667 (2) and -0.662 (3) Å, respectively.

In the molecule the amine (N1) H atom is involved in two short interactions with the hydroxyl O atoms, O1' and O1'' (Table 1). The hydroxyl H atoms are each disordered over two positions, H1A/H1B and H1C/H1D. Their occupancies were initially refined before being fixed at 0.57/0.43 and 0.63/0.37, respectively. The ¹H NMR signal for the hydroxyl H atoms [δ 2.88 (bs, 2 H, OH); see archived CIF] is a broad singlet, which indicates some fluxionality of these protons in solution.

In the crystal, four symmetry related molecules are linked by O—H...O hydrogen bonds to form a cage-like arrangement, centered about the point of intersection of three 2-fold axes (Fig 2). These cages are arranged in stacks along direction [100], as shown in Fig. 3.


S2. Experimental

The synthesis of the title compound (2) is illustrated in Fig. 4. The starting material, 2,2'-(pyridine-2,6-diyl)dipropan-2-ol (1), was prepared in one step from the commercially available dimethyl pyridine-2,6-dicarboxylate, according to the method described by (Klein *et al.*, 2009). The title compound (2), was synthesized by heating 0.5 g (2.56 mmol) of compound (1), together with 10% Pd/C (430 mg), methanol (10 ml) and acetic acid (10 ml), in an autoclave under hydrogen (50 atm), with stirring at 323 K for 12 h. For workup the reaction was filtered through a pad of celite and washed three times with dichloromethane. The solution was concentrated under vacuum to give a colourless slurry. The slurry was dissolved in dichloromethane and washed with 5% sodium hydroxide and the mixture was stirred for 5 min.

The organic layer was separated and the aqueous layer was extracted three times with dichloromethane. The combined organic layers were washed with brine, dried with sodium sulfate and concentrated under vacuum to yield 0.494 g (96%) of compound (2). Melting point: 345.3 K. HRMS calcd. for $[C_{11}H_{23}NO_2^+H^+]$ 224.1621; found 224.1621. Colourless rod-like crystals were obtained by slow evaporation of a solution of (2) in dichloromethane. Spectroscopic data for the title compound (2), are given the archived CIF.

S3. Refinement

The NH H-atom was located in a difference Fourier map and was freely refined. The OH H atoms are disordered over two positions. They were located in a difference Fourier map and were initially freely refined, including their occupancies, before being refined with distance restraints of 0.84 (2) Å. In the final cycles of refinement they were refined with fixed occupancies of 0.57/0.43 and 0.63/0.37, and allowed to ride on the parent O atom with $U_{iso}(H) = 1.5U_{eq}(O)$. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.98, 0.99 and 1.00 Å for CH₃, CH₂ and CH H-atoms, respectively, with $U_{iso}(H) = k \times U_{eq}(\text{parent C-atom})$, where $k = 1.5$ for CH₃ H-atoms and $k = 1.2$ for all other H-atoms.

**Figure 1**

A view of the molecular structure of the title molecule (2), with displacement ellipsoids drawn at the 30% probability level. (The O···H dashed lines indicate the positions of the minor components of the hydroxyl H atoms.)




Figure 2

A view of the hydrogen bonded cage formed by four symmetry related molecules of the title compound. The C-bound H atoms have been omitted for clarity. The O—H \cdots O and N—H \cdots O hydrogen bonds are shown as dashed cyan lines (see Table 1 for details).




Figure 3

A view along the *a* axis of the crystal packing of the title compound. The C-bound H atoms have been omitted for clarity. The O—H \cdots O and N—H \cdots O hydrogen bonds are shown as dashed cyan lines (see Table 1 for details).




Figure 4

Reaction scheme for the synthesis of the title compound, (2).

2,2'-[(2*S**,6*R**)-Piperidine-2,6-diyl]dipropan-2-ol

Crystal data

C₁₁H₂₃NO₂ $M_r = 201.30$ Orthorhombic, *Fddd*

Hall symbol: -F 2uv 2vw

 $a = 12.0713 (9) \text{ \AA}$ $b = 23.4762 (10) \text{ \AA}$ $c = 34.496 (2) \text{ \AA}$ $V = 9775.8 (10) \text{ \AA}^3$ $Z = 32$ $F(000) = 3584$ $D_x = 1.094 \text{ Mg m}^{-3}$

Melting point: 345.3 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 13139 reflections

 $\theta = 2.0\text{--}24.3^\circ$ $\mu = 0.07 \text{ mm}^{-1}$ $T = 173 \text{ K}$

Rod, colourless

 $0.45 \times 0.45 \times 0.40 \text{ mm}$

Data collection

Stoe IPDS 2

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ & ω scans

Absorption correction: multi-scan

(MULscanABS in PLATON; Spek, 2009)

 $T_{\min} = 0.911$, $T_{\max} = 1.000$

32226 measured reflections

2319 independent reflections

1499 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.135$ $\theta_{\max} = 25.7^\circ$, $\theta_{\min} = 2.0^\circ$ $h = -14 \rightarrow 14$ $k = -28 \rightarrow 28$ $l = -42 \rightarrow 41$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.086$ $wR(F^2) = 0.146$ $S = 1.16$

2319 reflections

135 parameters

4 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0468P)^2 + 7.437P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.16 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.15 \text{ e \AA}^{-3}$

Special details

Experimental. Spectroscopic data for 2,2'-[(2*S**,6*R**)-piperidine-2,6-diyl]dipropan-2-ol (2):

¹H NMR (CDCl₃, 298 K, p.p.m.) δ 2.88 (bs, 2 H, OH), 2.47 (d, ³J(2, 3 b) and ³J(6, 5 b) = 11.4 Hz, 2 H, C—H(2, 6)), 1.97 (dq, ³J(4 b, 4a) = 13.3 Hz, ³J(4 b, 3 b) = 3 J(4 b, 5 b) = 3.3 Hz, ³J(4 b, 3a) = ³J(4 b, 5a) = 3.3 Hz, 1 H, C—H(4 b)), 1.74 (dd, ³J(3a, 3 b) = 12.8 Hz and ³J(5a, 5 b) = 12.8 Hz, ³J(3a, 4 b) = 3.0 Hz and ³J(5a, 4 b) = 3.0 Hz, 2 H, C—H(3a, 5a)), 1.45 (qt, ³J(4a, 4 b) = 13.0 Hz, ³J(4a, 3 b) = 13.0 Hz and ³J(4a, 5 b) = 13.0 Hz, 3 J(4a, 3a) = 3.7 Hz and ³J(4a, 5a) = 3.7 Hz, 1 H, C—H(4a)), 1.25 (s, 6 H, CH₃), 1.16 (s, 6 H, CH₃), 1.06 (qd, ³J(3 b, 2) = 12.3 Hz and ³J(5 b, 6) = 12.3 Hz, ³J(3 b, 3a) = 12.3 Hz and ³J(5 b, 5a) = 12.3 Hz, ³J(3 b, 4a) = 12.3 Hz and ³J(5 b, 4a) = 12.3 Hz, ³J(3 b, 4 b) = 3.3 Hz and ³J(5 b, 4 b) = 3.3 Hz, 2 H, CH₂(3 b, 5 b));

¹³C NMR (CDCl₃, 298 K, p.p.m.) δ 71.67 (C(2, 2')), 65.45 (C(2, 6)), 27.45 (CH₃), 26.91 (C(3, 5)), 24.69 (C(4)), 24.36 (CH₃).

IR (KBr, cm⁻¹): 3377 b s, 2982 s, 2944 s, 2855m, 2782m, 2694w, 2586w, 1456m, 1442m, 1380 s, 1130 s, 931 s, 822 s, 537w.

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. The NH H-atom was located in a difference Fourier map and was freely refined. The OH H atoms are disordered over two positions with fixed occupancies of 0.57/0.43 and 0.63/0.37. They were located in a difference Fourier map and were initially freely refined, including their occupancies, before being refined with distance restraints of 0.84 (2) Å. In the final cycles of refinement they were refined with fixed occupancies of 0.57/0.43 and 0.63/0.37, and allowed to ride on the parent O atom with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.98, 0.99 and 1.00 Å for CH₃, CH₂ and CH H-atoms, respectively, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{parent C-atom})$, where $k = 1.5$ for CH₃ H-atoms and $k = 1.2$ for all other H-atoms.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1'	0.52422 (16)	0.09518 (8)	0.04534 (5)	0.0474 (7)	
O1''	0.51927 (19)	0.22777 (8)	0.14029 (6)	0.0602 (8)	
N1	0.41660 (19)	0.19356 (10)	0.07083 (6)	0.0344 (7)	
C1'	0.4778 (3)	0.15489 (12)	−0.00855 (8)	0.0490 (10)	
C1''	0.4860 (3)	0.30648 (12)	0.09761 (9)	0.0551 (11)	
C2	0.3546 (2)	0.22205 (11)	0.10141 (8)	0.0426 (10)	
C2'	0.4316 (2)	0.11656 (11)	0.02325 (8)	0.0410 (10)	
C2''	0.4314 (3)	0.26220 (11)	0.12374 (8)	0.0496 (10)	
C3	0.2533 (3)	0.25074 (14)	0.08304 (10)	0.0602 (13)	
C3'	0.3740 (3)	0.06505 (14)	0.00542 (10)	0.0730 (14)	
C3''	0.3697 (4)	0.29130 (15)	0.15692 (10)	0.0900 (16)	
C4	0.1844 (2)	0.20695 (15)	0.06117 (11)	0.0712 (13)	
C5	0.2544 (2)	0.17559 (14)	0.03176 (10)	0.0554 (11)	
C6	0.3549 (2)	0.14843 (11)	0.05111 (8)	0.0391 (9)	
H1	0.472 (2)	0.1777 (11)	0.0809 (7)	0.035 (8)*	
H1A	0.51390	0.07430	0.06420	0.0710*	0.570
H1C	0.51740	0.20950	0.16040	0.0900*	0.630
H1L	0.51510	0.18770	0.00320	0.0740*	
H1M	0.41710	0.16820	−0.02510	0.0740*	
H1N	0.53100	0.13340	−0.02420	0.0740*	
H1O	0.53320	0.28730	0.07860	0.0830*	
H1P	0.53110	0.33230	0.11340	0.0830*	
H1Q	0.42860	0.32830	0.08410	0.0830*	
H2	0.32750	0.19230	0.11990	0.0510*	
H3A	0.20760	0.26860	0.10350	0.0720*	
H3B	0.27790	0.28100	0.06500	0.0720*	
H3C	0.42710	0.04350	−0.01030	0.1090*	
H3D	0.31270	0.07790	−0.01100	0.1090*	
H3E	0.34510	0.04060	0.02610	0.1090*	
H3F	0.32950	0.26260	0.17200	0.1350*	
H3G	0.31710	0.31900	0.14630	0.1350*	
H3H	0.42300	0.31090	0.17370	0.1350*	
H4A	0.15250	0.17930	0.07980	0.0850*	
H4B	0.12240	0.22640	0.04780	0.0850*	
H5A	0.20940	0.14570	0.01910	0.0670*	
H5B	0.27950	0.20250	0.01150	0.0670*	
H6	0.32790	0.12090	0.07110	0.0470*	

H1B	0.57870	0.11630	0.04260	0.0710*	0.430
H1D	0.57710	0.24160	0.13160	0.0900*	0.370

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1'	0.0614 (13)	0.0440 (11)	0.0367 (12)	0.0143 (10)	-0.0002 (10)	0.0061 (9)
O1''	0.0968 (18)	0.0448 (12)	0.0391 (12)	0.0016 (12)	-0.0205 (12)	0.0086 (10)
N1	0.0292 (12)	0.0401 (13)	0.0340 (13)	0.0063 (11)	-0.0009 (11)	0.0029 (11)
C1'	0.0569 (19)	0.0521 (18)	0.0381 (17)	0.0011 (15)	0.0021 (15)	0.0106 (14)
C1''	0.073 (2)	0.0422 (17)	0.050 (2)	-0.0020 (16)	-0.0034 (17)	0.0073 (15)
C2	0.0459 (17)	0.0385 (15)	0.0435 (18)	0.0099 (14)	0.0160 (14)	0.0092 (14)
C2'	0.0513 (18)	0.0373 (16)	0.0345 (16)	-0.0044 (14)	-0.0085 (14)	0.0051 (13)
C2''	0.076 (2)	0.0388 (15)	0.0340 (17)	0.0091 (16)	0.0109 (16)	0.0048 (14)
C3	0.0441 (18)	0.0564 (19)	0.080 (3)	0.0191 (15)	0.0206 (17)	0.0162 (19)
C3'	0.112 (3)	0.057 (2)	0.050 (2)	-0.026 (2)	-0.014 (2)	-0.0008 (17)
C3''	0.158 (4)	0.061 (2)	0.051 (2)	0.023 (2)	0.038 (3)	-0.0051 (19)
C4	0.0315 (17)	0.075 (2)	0.107 (3)	0.0104 (17)	-0.0006 (19)	0.037 (2)
C5	0.0342 (17)	0.066 (2)	0.066 (2)	-0.0054 (15)	-0.0126 (16)	0.0203 (18)
C6	0.0359 (15)	0.0429 (16)	0.0384 (17)	-0.0065 (13)	-0.0060 (13)	0.0128 (13)

Geometric parameters (Å, °)

O1'—C2'	1.443 (3)	C1'—H1M	0.9800
O1''—C2''	1.451 (4)	C1'—H1N	0.9800
O1'—H1B	0.8300	C1''—H1O	0.9800
O1'—H1A	0.8200	C1''—H1P	0.9800
O1''—H1C	0.8200	C1''—H1Q	0.9800
O1''—H1D	0.8300	C2—H2	1.0000
N1—C6	1.463 (3)	C3—H3A	0.9900
N1—C2	1.456 (3)	C3—H3B	0.9900
N1—H1	0.84 (2)	C3'—H3C	0.9800
C1'—C2'	1.525 (4)	C3'—H3D	0.9800
C1''—C2''	1.526 (4)	C3'—H3E	0.9800
C2—C2''	1.530 (4)	C3''—H3F	0.9800
C2—C3	1.533 (4)	C3''—H3G	0.9800
C2'—C3'	1.525 (4)	C3''—H3H	0.9800
C2'—C6	1.530 (4)	C4—H4A	0.9900
C2''—C3''	1.527 (5)	C4—H4B	0.9900
C3—C4	1.522 (5)	C5—H5A	0.9900
C4—C5	1.512 (5)	C5—H5B	0.9900
C5—C6	1.524 (4)	C6—H6	1.0000
C1'—H1L	0.9800		
C2'—O1'—H1A	120.00	H1O—C1''—H1P	109.00
C2'—O1'—H1B	110.00	H1O—C1''—H1Q	109.00
C2''—O1''—H1C	127.00	H1P—C1''—H1Q	110.00
C2''—O1''—H1D	105.00	N1—C2—H2	108.00

C2—N1—C6	114.1 (2)	C2''—C2—H2	108.00
C6—N1—H1	106.1 (18)	C3—C2—H2	108.00
C2—N1—H1	108.2 (17)	C2—C3—H3A	110.00
N1—C2—C3	108.2 (2)	C2—C3—H3B	110.00
N1—C2—C2''	109.7 (2)	C4—C3—H3A	110.00
C2''—C2—C3	114.9 (2)	C4—C3—H3B	110.00
O1'—C2'—C6	107.9 (2)	H3A—C3—H3B	108.00
O1'—C2'—C1'	107.6 (2)	C2'—C3'—H3C	110.00
O1'—C2'—C3'	106.9 (2)	C2'—C3'—H3D	109.00
C1'—C2'—C6	112.6 (2)	C2'—C3'—H3E	109.00
C3'—C2'—C6	111.4 (2)	H3C—C3'—H3D	109.00
C1'—C2'—C3'	110.2 (2)	H3C—C3'—H3E	109.00
C1''—C2''—C2	112.6 (2)	H3D—C3'—H3E	109.00
C1''—C2''—C3''	110.4 (2)	C2''—C3''—H3F	109.00
O1''—C2''—C1''	107.2 (3)	C2''—C3''—H3G	109.00
O1''—C2''—C2	107.3 (2)	C2''—C3''—H3H	109.00
O1''—C2''—C3''	108.1 (2)	H3F—C3''—H3G	110.00
C2—C2''—C3''	110.9 (3)	H3F—C3''—H3H	110.00
C2—C3—C4	110.1 (3)	H3G—C3''—H3H	110.00
C3—C4—C5	110.9 (2)	C3—C4—H4A	109.00
C4—C5—C6	110.8 (3)	C3—C4—H4B	109.00
C2'—C6—C5	114.3 (2)	C5—C4—H4A	109.00
N1—C6—C2'	109.8 (2)	C5—C4—H4B	110.00
N1—C6—C5	107.8 (2)	H4A—C4—H4B	108.00
C2'—C1'—H1L	110.00	C4—C5—H5A	110.00
C2'—C1'—H1M	110.00	C4—C5—H5B	110.00
C2'—C1'—H1N	109.00	C6—C5—H5A	109.00
H1L—C1'—H1M	110.00	C6—C5—H5B	109.00
H1L—C1'—H1N	109.00	H5A—C5—H5B	108.00
H1M—C1'—H1N	109.00	N1—C6—H6	108.00
C2''—C1''—H1O	110.00	C2'—C6—H6	108.00
C2''—C1''—H1P	109.00	C5—C6—H6	108.00
C2''—C1''—H1Q	109.00		
C6—N1—C2—C2''	172.0 (2)	C2''—C2—C3—C4	179.1 (3)
C6—N1—C2—C3	-62.0 (3)	O1'—C2'—C6—N1	56.7 (3)
C2—N1—C6—C2'	-173.0 (2)	O1'—C2'—C6—C5	177.9 (2)
C2—N1—C6—C5	62.0 (3)	C1'—C2'—C6—N1	-61.9 (3)
N1—C2—C2''—O1''	-60.2 (3)	C1'—C2'—C6—C5	59.4 (3)
N1—C2—C2''—C1''	57.6 (3)	C3'—C2'—C6—N1	173.7 (2)
N1—C2—C2''—C3''	-178.1 (2)	C3'—C2'—C6—C5	-65.0 (3)
C3—C2—C2''—O1''	177.7 (2)	C2—C3—C4—C5	-54.9 (4)
C3—C2—C2''—C1''	-64.5 (3)	C3—C4—C5—C6	55.4 (4)
C3—C2—C2''—C3''	59.8 (3)	C4—C5—C6—N1	-56.7 (3)
N1—C2—C3—C4	56.2 (3)	C4—C5—C6—C2'	-179.0 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1···O1'	0.84 (3)	2.38 (3)	2.792 (3)	111 (2)
N1—H1···O1''	0.84 (3)	2.43 (3)	2.814 (3)	109 (2)
O1'—H1A···O1'' ⁱ	0.82	1.99	2.805 (3)	169
O1'—H1B···O1'' ⁱⁱ	0.83	1.99	2.807 (4)	167
O1''—H1C···O1'' ⁱ	0.82	2.00	2.805 (3)	171
O1''—H1D···O1'' ⁱⁱⁱ	0.83	2.03	2.762 (5)	148

Symmetry codes: (i) $x, -y+1/4, -z+1/4$; (ii) $-x+5/4, -y+1/4, z$; (iii) $-x+5/4, y, -z+1/4$.