

(E)-1-[1-(3-Chlorophenyl)ethylidene]-2-(2,4-dinitrophenyl)hydrazine

Hoong-Kun Fun,^{a,*‡} Suchada Chantrapromma,^{b,§} Boonlerd Nilwanna^b and Chatchanok Karalai^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

Correspondence e-mail: hkfun@usm.my


Received 2 February 2012; accepted 7 February 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.051; wR factor = 0.183; data-to-parameter ratio = 20.7.

There are two crystallographically independent molecules in the asymmetric unit of the title compound, $\text{C}_{14}\text{H}_{11}\text{ClN}_4\text{O}_4$, with the same *E* conformation about the $\text{C}=\text{N}$ double bond. The molecules are approximately planar, with a dihedral angle between the benzene rings of $10.24(12)^\circ$ in one molecule and $4.73(12)^\circ$ in the other. In both molecules, the *ortho*-nitro groups of the 2,4-dinitrophenyl units are coplanar to their bound benzene rings, whereas the *para*-nitro groups are slightly twisted. In each molecule, intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds generate *S*(6) ring motifs. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ interactions into sheets parallel to the $(\bar{1}02)$ plane. These sheets are stacked by $\pi-\pi$ interactions, with centroid–centroid distances of $3.7008(14)$ and $3.7459(14)\text{ \AA}$. A $\text{Cl}\cdots\text{O}$ short contact [$3.111(2)\text{ \AA}$] is observed.

Related literature

For bond-length data, see: Allen *et al.* (1987). For related literature on hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: Chantrapromma *et al.* (2011); Fun *et al.* (2011, 2012); Nilwanna *et al.* (2011). For background to biological activities of hydrazones, see: Angelusiu *et al.* (2010); Cui *et al.* (2010); Gokce *et al.* (2009); Khan *et al.* (2007); Loncle *et al.* (2004); Wang *et al.* (2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{11}\text{ClN}_4\text{O}_4$	$V = 2941.8(5)\text{ \AA}^3$
$M_r = 334.72$	$Z = 8$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 13.4825(13)\text{ \AA}$	$\mu = 0.29\text{ mm}^{-1}$
$b = 15.1586(15)\text{ \AA}$	$T = 296\text{ K}$
$c = 16.1281(12)\text{ \AA}$	$0.42 \times 0.19 \times 0.18\text{ mm}$
$\beta = 116.815(6)^\circ$	

Data collection

Bruker APEX DUO CCD area-detector diffractometer	32600 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	8629 independent reflections
$T_{\min} = 0.889$, $T_{\max} = 0.951$	4617 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$	1 restraint
$wR(F^2) = 0.183$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\max} = 0.35\text{ e \AA}^{-3}$
8629 reflections	$\Delta\rho_{\min} = -0.34\text{ e \AA}^{-3}$
417 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1A—H1NA \cdots O1A	0.82	1.95	2.598 (2)	135
N1B—H1NB \cdots O1B	0.86	1.86	2.589 (2)	141
C5A—H5A \cdots O1A ⁱ	0.93	2.52	3.251 (3)	136
C5B—H5B \cdots O1B ⁱⁱ	0.93	2.33	3.196 (3)	154
C11A—H11A \cdots O3B ⁱⁱⁱ	0.93	2.55	3.402 (3)	153
C11B—H11B \cdots O3A ^{iv}	0.93	2.58	3.429 (3)	153

Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 2, y + \frac{1}{2}, -z + \frac{5}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors thank the Prince of Songkla University for financial support through the Crystal Materials Research Unit. The authors also thank the Universiti Sains Malaysia for the Research University grant No. 1001/PFIZIK/811160.

‡ Thomson Reuters ResearcherID: A-3561-2009.

§ Additional correspondence author, e-mail: suchada.c@psu.ac.th. Thomson Reuters ResearcherID: A-5085-2009.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2707).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Angelusiu, M.-V., Barbuceanu, S.-F., Draghici, C. & Almajan, G.-L. (2010). *Eur. J. Med. Chem.* **45**, 2055–2062.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chantrapromma, S., Nilwanna, B., Jansrisewangwong, P., Kobkeatthawin, T. & Fun, H.-K. (2011). *Acta Cryst. E67*, o3499–o3500.
- Cui, Z., Li, Y., Ling, Y., Huang, J., Cui, J., Wang, R. & Yang, X. (2010). *Eur. J. Med. Chem.* **45**, 5576–5584.
- Fun, H.-K., Nilwanna, B., Chantrapromma, S. & Razak, I. A. (2012). *Acta Cryst. E68*, o398–o399.
- Fun, H.-K., Nilwanna, B., Jansrisewangwong, P., Kobkeatthawin, T. & Chantrapromma, S. (2011). *Acta Cryst. E67*, o3202–o3203.
- Gokce, M., Utku, S. & Kupeli, E. (2009). *Eur. J. Med. Chem.* **44**, 3760–3764.
- Khan, S. A., Saleem, K. & Khan, Z. (2007). *Eur. J. Med. Chem.* **42**, 103–108.
- Loncle, C., Brunel, J. M., Vidal, N., Dherbomez, M. & Letourneux, Y. (2004). *Eur. J. Med. Chem.* **39**, 1067–1071.
- Nilwanna, B., Chantrapromma, S., Jansrisewangwong, P. & Fun, H.-K. (2011). *Acta Cryst. E67*, o3084–o3085.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Wang, Q., Yang, Z. Y., Qi, G.-F. & Qin, D.-D. (2009). *Eur. J. Med. Chem.* **44**, 2425–2433.

supporting information

Acta Cryst. (2012). E68, o704–o705 [doi:10.1107/S160053681200548X]

(E)-1-[1-(3-Chlorophenyl)ethylidene]-2-(2,4-dinitrophenyl)hydrazine

Hoong-Kun Fun, Suchada Chantrapromma, Boonlerd Nilwanna and Chatchanok Karalai

S1. Comment

Hydrazones are known to be bioactive compounds with antibacterial, antifungal, antitumor, anti-inflammatory as well as antioxidant (Angelusiu *et al.*, 2010; Cui *et al.*, 2010; Gokce *et al.*, 2009; Khan *et al.*, 2007; Loncle *et al.*, 2004; Wang *et al.*, 2009) activities. Within our on-going research on the bioactivity of hydrazones, the title compound (I) was synthesized in order to study and compare its biological activity with other related compounds (Chantrapromma *et al.*, 2011; Fun *et al.*, 2011; 2012; Nilwanna *et al.*, 2011). Herein we report the synthesis and crystal structure of (I).

There are two crystallographic independent molecules *A* and *B* in the asymmetric unit of (I) with differences in bond angles (Fig. 1). The molecular structure of (I) is nearly planar with the dihedral angle between the two benzene rings of 10.24 (12)° in molecule *A* and 4.73 (12)° in molecule *B*. The central ethylenedihydrazine bridge (N1/N2/C7/C14) is planar with the torsion angles N1–N2–C7–C14 = 0.8 (3) and 0.5 (3)° in molecules *A* and *B*, respectively. The mean plane through this central bridge makes dihedral angles of 6.36 (17) and 3.90 (18)° with the 2,4-dinitrophenyl and 3-chlorophenyl rings, respectively in molecule *A* whereas the corresponding values are 5.37 (15) and 0.90 (15)° in molecule *B*. In both molecules, the *ortho*-nitro group of the 2,4-dinitrophenyl is coplanar with the attached benzene ring with the *r.m.s.* deviation of 0.0164 (2) Å for the nine non H-atoms (C1–C6/N3/O1/O2), and torsion angles O1–N3–C2–C3 = 176.84 (18)° and O2–N3–C2–C3 = -1.9 (3)°, whereas the *para*-nitro group is slightly twisted with the torsion angles O3–N4–C4–C5 = 168.8 (2)° and O4–N4–C4–C5 = -11.5 (3)° in molecule *A*; the corresponding values are 0.0176 (2) Å, -177.77 (18), 3.6 (3), 171.1 (2) and -8.9 (4)° in molecule *B*. In each molecule, intramolecular N—H···O hydrogen bond (Fig. 1 and Table 1) generates S(6) ring motifs (Bernstein *et al.*, 1995) The bond distances agree with the literature values (Allen *et al.*, 1987) and are comparable with the related structures (Chantrapromma *et al.*, 2011; Fun *et al.*, 2011; 2012; Nilwanna *et al.*, 2011).


In the crystal packing (Fig. 2), the molecules are linked by weak C—H···O interactions (Table 1) into sheets parallel to the (-102) plane. These sheets are further stacked along the *a* axis by π – π interactions with distances of $Cg_1\cdots Cg_2^v$ = 3.7459 (14) Å and $Cg_1\cdots Cg_3^{vi}$ = 3.7008 (14) Å [symmetry codes (*v*) = *x*, 3/2–*y*, 1/2+*z*; (*vi*) = 2–*x*, 2–*y*, 2–*z*]; Cg_1 , Cg_2 and Cg_3 are the centroids of C1A–C6A, C8A–C13A and C8B–C13B benzene rings, respectively. A $C1B\cdots O1B^{ii}$ [3.111 (2) Å] short contact is observed.

S2. Experimental


The title compound (I) was synthesized by dissolving 2,4-dinitrophenylhydrazine (0.40 g, 2 mmol) in ethanol (10.00 ml) and H_2SO_4 (conc.) (98 %, 0.50 ml) was slowly added with stirring. 3-Chloroacetophenone (0.26 ml, 2 mmol) was then added to the solution with continuous stirring. The solution was stirred for 1 h yielding an orange solid, which was filtered off and washed with methanol. Orange block-shaped single crystals of the title compound suitable for X-ray structure determination were recrystallized from ethanol by slow evaporation of the solvent at room temperature over several days. M.p. 478–479 K.

S3. Refinement

All H atoms were positioned geometrically and allowed to ride on their parent atoms, with $d(\text{N-H}) = 0.83$ and 0.86 \AA , $d(\text{C-H}) = 0.93 \text{ \AA}$ for aromatic and 0.96 \AA for CH_3 atoms. The U_{iso} values were constrained to be $1.5U_{\text{eq}}$ of the carrier atom for methyl H atoms and $1.2U_{\text{eq}}$ for the remaining H atoms. A rotating group model was used for the methyl groups. A DFIX restraint of $2.00(1) \text{ \AA}$ was used for the $\text{H}14\text{D}\cdots\text{H}1\text{NB}$ distance. An outlier (0 2 0) was omitted.

**Figure 1**

The molecular structure of (I), showing 50% probability displacement ellipsoids. Intramolecular $\text{N—H}\cdots\text{O}$ hydrogen bonds are shown as dashed lines.

**Figure 2**

The crystal packing of (I) viewed approximately along the b axis. Hydrogen bonds are shown as dashed lines.

(E)-1-[1-(3-Chlorophenyl)ethylidene]-2-(2,4-dinitrophenyl)hydrazine

Crystal data

$C_{14}H_{11}ClN_4O_4$
 $M_r = 334.72$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 13.4825$ (13) Å
 $b = 15.1586$ (15) Å
 $c = 16.1281$ (12) Å
 $\beta = 116.815$ (6)°
 $V = 2941.8$ (5) Å³
 $Z = 8$

$F(000) = 1376$
 $D_x = 1.512 \text{ Mg m}^{-3}$
Melting point = 478–479 K
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 8629 reflections
 $\theta = 1.7\text{--}30.1^\circ$
 $\mu = 0.29 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
Block, orange
0.42 × 0.19 × 0.18 mm

Data collection

Bruker APEX DUO CCD area-detector
diffractometer
Radiation source: sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2009)
 $T_{\min} = 0.889$, $T_{\max} = 0.951$

32600 measured reflections
8629 independent reflections
4617 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 30.1^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -18 \rightarrow 18$
 $k = -21 \rightarrow 19$
 $l = -22 \rightarrow 22$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.183$
 $S = 1.01$
8629 reflections

417 parameters
1 restraint
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0916P)^2 + 0.289P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.35 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1A	0.83654 (6)	0.45629 (4)	0.39055 (4)	0.0766 (2)
O1A	0.99997 (15)	0.97920 (10)	0.69823 (12)	0.0774 (5)
O2A	1.06555 (14)	0.99861 (10)	0.84524 (12)	0.0711 (5)
O3A	1.2204 (2)	0.76410 (15)	1.05861 (12)	0.1103 (8)
O4A	1.18161 (18)	0.63048 (13)	1.01190 (12)	0.0949 (6)
N1A	0.97877 (14)	0.82236 (11)	0.62719 (11)	0.0553 (4)
H1NA	0.9589	0.8744	0.6200	0.066*
N2A	0.94483 (14)	0.76046 (11)	0.55856 (11)	0.0536 (4)
N3A	1.03978 (14)	0.95052 (11)	0.77797 (13)	0.0545 (4)
N4A	1.18006 (17)	0.70917 (15)	0.99701 (13)	0.0688 (5)
C1A	1.02714 (15)	0.79618 (12)	0.71746 (13)	0.0457 (4)
C2A	1.05891 (15)	0.85647 (12)	0.79264 (13)	0.0449 (4)
C3A	1.10886 (15)	0.82829 (13)	0.88368 (13)	0.0481 (4)
H3A	1.1298	0.8685	0.9322	0.058*
C4A	1.12712 (16)	0.73977 (13)	0.90130 (13)	0.0502 (5)
C5A	1.09807 (17)	0.67883 (13)	0.82982 (14)	0.0556 (5)
H5A	1.1124	0.6192	0.8435	0.067*
C6A	1.04898 (18)	0.70614 (12)	0.74032 (14)	0.0533 (5)
H6A	1.0293	0.6647	0.6930	0.064*
C7A	0.90349 (17)	0.78840 (14)	0.47443 (14)	0.0534 (5)
C8A	0.86585 (16)	0.71840 (15)	0.40185 (13)	0.0514 (5)
C9A	0.87083 (16)	0.63003 (14)	0.42698 (13)	0.0525 (5)
H9A	0.8993	0.6143	0.4893	0.063*
C10A	0.83347 (17)	0.56618 (15)	0.35924 (14)	0.0558 (5)
C11A	0.79092 (17)	0.58681 (18)	0.26572 (14)	0.0636 (6)
H11A	0.7651	0.5428	0.2208	0.076*
C12A	0.78777 (18)	0.67358 (19)	0.24109 (15)	0.0678 (6)
H12A	0.7610	0.6885	0.1787	0.081*
C13A	0.82399 (18)	0.73955 (17)	0.30792 (14)	0.0614 (6)
H13A	0.8203	0.7982	0.2900	0.074*

C14A	0.8916 (3)	0.88420 (17)	0.44724 (17)	0.0849 (8)
H14A	0.9563	0.9159	0.4895	0.127*
H14B	0.8270	0.9082	0.4495	0.127*
H14C	0.8839	0.8896	0.3853	0.127*
Cl1B	0.60994 (6)	1.29099 (4)	1.10126 (4)	0.0783 (2)
O1B	0.58475 (15)	0.77573 (10)	0.80697 (11)	0.0683 (4)
O2B	0.51645 (18)	0.75501 (10)	0.66003 (13)	0.0860 (6)
O3B	0.35381 (19)	0.98474 (14)	0.44260 (11)	0.1013 (7)
O4B	0.3250 (2)	1.11127 (14)	0.48825 (13)	0.1090 (8)
N1B	0.58812 (13)	0.93100 (11)	0.87512 (11)	0.0490 (4)
H1NB	0.6001	0.8750	0.8796	0.059*
N2B	0.60588 (13)	0.99163 (11)	0.94310 (10)	0.0474 (4)
N3B	0.53830 (15)	0.80306 (11)	0.72640 (13)	0.0556 (4)
N4B	0.36352 (19)	1.03694 (14)	0.50342 (13)	0.0729 (6)
C1B	0.53726 (15)	0.95592 (12)	0.78481 (12)	0.0432 (4)
C2B	0.51113 (15)	0.89601 (11)	0.71062 (13)	0.0445 (4)
C3B	0.45551 (17)	0.92247 (13)	0.61880 (13)	0.0507 (5)
H3B	0.4394	0.8822	0.5709	0.061*
C4B	0.42477 (17)	1.00861 (13)	0.59995 (13)	0.0517 (5)
C5B	0.45036 (18)	1.07026 (13)	0.67045 (14)	0.0564 (5)
H5B	0.4293	1.1289	0.6559	0.068*
C6B	0.50633 (18)	1.04464 (12)	0.76078 (14)	0.0513 (5)
H6B	0.5245	1.0865	0.8076	0.062*
C7B	0.64883 (16)	0.96382 (13)	1.02719 (13)	0.0479 (4)
C8B	0.66519 (15)	1.03272 (14)	1.09812 (13)	0.0475 (4)
C9B	0.63316 (16)	1.11936 (14)	1.07065 (13)	0.0505 (5)
H9B	0.6015	1.1343	1.0080	0.061*
C10B	0.64837 (16)	1.18318 (15)	1.13627 (14)	0.0539 (5)
C11B	0.69611 (17)	1.16370 (17)	1.22983 (14)	0.0589 (5)
H11B	0.7068	1.2075	1.2734	0.071*
C12B	0.72749 (19)	1.07829 (18)	1.25720 (14)	0.0631 (6)
H12B	0.7595	1.0640	1.3200	0.076*
C13B	0.71201 (17)	1.01338 (15)	1.19250 (14)	0.0572 (5)
H13B	0.7332	0.9557	1.2123	0.069*
C14B	0.6820 (2)	0.86989 (15)	1.05689 (15)	0.0722 (7)
H14D	0.6413	0.8311	1.0056	0.108*
H14E	0.6656	0.8557	1.1074	0.108*
H14F	0.7602	0.8629	1.0766	0.108*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1A	0.0978 (5)	0.0648 (4)	0.0690 (4)	-0.0055 (3)	0.0391 (3)	-0.0079 (3)
O1A	0.1049 (13)	0.0437 (8)	0.0721 (11)	0.0150 (8)	0.0298 (10)	0.0122 (8)
O2A	0.0885 (12)	0.0463 (8)	0.0820 (11)	0.0005 (8)	0.0417 (10)	-0.0156 (8)
O3A	0.159 (2)	0.0944 (15)	0.0468 (10)	-0.0004 (14)	0.0199 (11)	0.0000 (10)
O4A	0.1274 (16)	0.0722 (12)	0.0691 (11)	0.0088 (11)	0.0301 (11)	0.0272 (9)
N1A	0.0674 (11)	0.0427 (9)	0.0489 (9)	0.0047 (8)	0.0200 (8)	0.0013 (7)

N2A	0.0591 (10)	0.0508 (9)	0.0439 (9)	0.0023 (8)	0.0170 (8)	0.0003 (7)
N3A	0.0582 (10)	0.0393 (8)	0.0657 (11)	0.0026 (7)	0.0278 (9)	-0.0008 (8)
N4A	0.0792 (13)	0.0687 (13)	0.0522 (11)	0.0052 (10)	0.0240 (10)	0.0111 (10)
C1A	0.0472 (10)	0.0407 (9)	0.0468 (10)	0.0006 (8)	0.0192 (8)	0.0013 (8)
C2A	0.0481 (10)	0.0346 (9)	0.0529 (10)	0.0007 (7)	0.0236 (8)	-0.0007 (8)
C3A	0.0483 (10)	0.0478 (10)	0.0485 (10)	-0.0041 (8)	0.0223 (8)	-0.0054 (8)
C4A	0.0537 (11)	0.0496 (11)	0.0446 (10)	-0.0004 (9)	0.0199 (9)	0.0039 (8)
C5A	0.0666 (13)	0.0363 (9)	0.0582 (12)	0.0001 (9)	0.0233 (10)	0.0042 (8)
C6A	0.0657 (12)	0.0369 (9)	0.0504 (11)	-0.0002 (8)	0.0199 (9)	-0.0035 (8)
C7A	0.0497 (11)	0.0563 (12)	0.0512 (11)	0.0054 (9)	0.0200 (9)	0.0077 (9)
C8A	0.0441 (10)	0.0649 (13)	0.0442 (10)	0.0063 (9)	0.0192 (8)	0.0061 (9)
C9A	0.0506 (11)	0.0646 (13)	0.0408 (10)	0.0008 (9)	0.0193 (8)	0.0008 (9)
C10A	0.0503 (11)	0.0668 (13)	0.0505 (11)	0.0014 (10)	0.0231 (9)	-0.0014 (10)
C11A	0.0521 (12)	0.0900 (18)	0.0467 (11)	-0.0003 (11)	0.0205 (10)	-0.0109 (11)
C12A	0.0616 (13)	0.0988 (19)	0.0394 (11)	0.0085 (13)	0.0194 (10)	0.0060 (11)
C13A	0.0604 (12)	0.0751 (15)	0.0459 (11)	0.0089 (11)	0.0214 (9)	0.0124 (10)
C14A	0.122 (2)	0.0617 (15)	0.0613 (15)	0.0093 (15)	0.0331 (15)	0.0130 (12)
Cl1B	0.1019 (5)	0.0664 (4)	0.0699 (4)	0.0174 (3)	0.0415 (4)	-0.0006 (3)
O1B	0.0926 (12)	0.0426 (8)	0.0695 (10)	0.0050 (7)	0.0364 (9)	0.0099 (7)
O2B	0.1302 (16)	0.0440 (8)	0.0760 (11)	0.0024 (9)	0.0396 (11)	-0.0144 (8)
O3B	0.159 (2)	0.0877 (14)	0.0431 (9)	0.0147 (13)	0.0329 (11)	0.0020 (9)
O4B	0.162 (2)	0.0778 (13)	0.0645 (11)	0.0374 (14)	0.0311 (12)	0.0214 (10)
N1B	0.0603 (10)	0.0419 (8)	0.0455 (8)	-0.0016 (7)	0.0246 (7)	-0.0001 (7)
N2B	0.0531 (9)	0.0474 (9)	0.0445 (8)	-0.0053 (7)	0.0245 (7)	-0.0007 (7)
N3B	0.0685 (11)	0.0395 (9)	0.0606 (11)	-0.0029 (8)	0.0307 (9)	-0.0032 (8)
N4B	0.0972 (15)	0.0665 (13)	0.0486 (11)	0.0051 (11)	0.0273 (10)	0.0099 (10)
C1B	0.0486 (10)	0.0414 (9)	0.0452 (10)	-0.0051 (7)	0.0260 (8)	0.0002 (7)
C2B	0.0517 (10)	0.0348 (9)	0.0504 (10)	-0.0048 (7)	0.0261 (9)	-0.0017 (7)
C3B	0.0636 (12)	0.0452 (10)	0.0470 (10)	-0.0059 (9)	0.0284 (9)	-0.0049 (8)
C4B	0.0663 (13)	0.0476 (11)	0.0425 (10)	-0.0020 (9)	0.0257 (9)	0.0028 (8)
C5B	0.0775 (14)	0.0387 (10)	0.0562 (12)	0.0016 (9)	0.0331 (11)	0.0065 (9)
C6B	0.0708 (13)	0.0386 (9)	0.0481 (10)	-0.0031 (9)	0.0301 (10)	-0.0033 (8)
C7B	0.0459 (10)	0.0526 (11)	0.0448 (10)	-0.0042 (8)	0.0202 (8)	0.0046 (8)
C8B	0.0440 (10)	0.0578 (12)	0.0419 (10)	-0.0066 (8)	0.0204 (8)	0.0019 (8)
C9B	0.0500 (11)	0.0621 (12)	0.0392 (9)	-0.0009 (9)	0.0200 (8)	0.0025 (9)
C10B	0.0508 (11)	0.0626 (13)	0.0513 (11)	-0.0009 (9)	0.0258 (9)	-0.0022 (9)
C11B	0.0598 (12)	0.0763 (15)	0.0458 (11)	-0.0061 (11)	0.0284 (10)	-0.0090 (10)
C12B	0.0655 (13)	0.0870 (17)	0.0378 (10)	-0.0071 (12)	0.0243 (9)	0.0029 (11)
C13B	0.0603 (12)	0.0664 (13)	0.0451 (10)	-0.0026 (10)	0.0240 (9)	0.0083 (10)
C14B	0.0949 (18)	0.0579 (14)	0.0520 (12)	0.0059 (12)	0.0228 (12)	0.0077 (10)

Geometric parameters (\AA , $^\circ$)

Cl1A—C10A	1.736 (2)	Cl1B—C10B	1.730 (2)
O1A—N3A	1.228 (2)	O1B—N3B	1.232 (2)
O2A—N3A	1.220 (2)	O2B—N3B	1.216 (2)
O3A—N4A	1.220 (3)	O3B—N4B	1.221 (3)
O4A—N4A	1.215 (3)	O4B—N4B	1.218 (3)

N1A—C1A	1.359 (2)	N1B—C1B	1.354 (2)
N1A—N2A	1.363 (2)	N1B—N2B	1.366 (2)
N1A—H1NA	0.8247	N1B—H1NB	0.8610
N2A—C7A	1.284 (2)	N2B—C7B	1.282 (2)
N3A—C2A	1.449 (2)	N3B—C2B	1.449 (2)
N4A—C4A	1.454 (3)	N4B—C4B	1.460 (3)
C1A—C6A	1.410 (3)	C1B—C6B	1.410 (2)
C1A—C2A	1.421 (3)	C1B—C2B	1.414 (2)
C2A—C3A	1.378 (3)	C2B—C3B	1.384 (3)
C3A—C4A	1.371 (3)	C3B—C4B	1.362 (3)
C3A—H3A	0.9300	C3B—H3B	0.9300
C4A—C5A	1.389 (3)	C4B—C5B	1.390 (3)
C5A—C6A	1.353 (3)	C5B—C6B	1.361 (3)
C5A—H5A	0.9300	C5B—H5B	0.9300
C6A—H6A	0.9300	C6B—H6B	0.9300
C7A—C8A	1.489 (3)	C7B—C8B	1.490 (3)
C7A—C14A	1.505 (3)	C7B—C14B	1.504 (3)
C8A—C9A	1.392 (3)	C8B—C13B	1.391 (3)
C8A—C13A	1.395 (3)	C8B—C9B	1.391 (3)
C9A—C10A	1.374 (3)	C9B—C10B	1.379 (3)
C9A—H9A	0.9300	C9B—H9B	0.9300
C10A—C11A	1.386 (3)	C10B—C11B	1.379 (3)
C11A—C12A	1.369 (4)	C11B—C12B	1.372 (4)
C11A—H11A	0.9300	C11B—H11B	0.9300
C12A—C13A	1.388 (3)	C12B—C13B	1.380 (3)
C12A—H12A	0.9300	C12B—H12B	0.9300
C13A—H13A	0.9300	C13B—H13B	0.9300
C14A—H14A	0.9600	C14B—H14D	0.9600
C14A—H14B	0.9600	C14B—H14E	0.9600
C14A—H14C	0.9600	C14B—H14F	0.9600
C1A—N1A—N2A	119.48 (16)	C1B—N1B—N2B	119.72 (16)
C1A—N1A—H1NA	113.5	C1B—N1B—H1NB	110.5
N2A—N1A—H1NA	125.2	N2B—N1B—H1NB	129.6
C7A—N2A—N1A	117.24 (18)	C7B—N2B—N1B	117.36 (17)
O2A—N3A—O1A	122.28 (17)	O2B—N3B—O1B	122.21 (18)
O2A—N3A—C2A	118.97 (18)	O2B—N3B—C2B	119.09 (18)
O1A—N3A—C2A	118.74 (17)	O1B—N3B—C2B	118.69 (16)
O4A—N4A—O3A	123.2 (2)	O4B—N4B—O3B	123.8 (2)
O4A—N4A—C4A	118.6 (2)	O4B—N4B—C4B	118.1 (2)
O3A—N4A—C4A	118.2 (2)	O3B—N4B—C4B	118.1 (2)
N1A—C1A—C6A	120.38 (17)	N1B—C1B—C6B	120.37 (17)
N1A—C1A—C2A	122.72 (17)	N1B—C1B—C2B	122.88 (16)
C6A—C1A—C2A	116.90 (17)	C6B—C1B—C2B	116.75 (17)
C3A—C2A—C1A	121.59 (17)	C3B—C2B—C1B	121.82 (17)
C3A—C2A—N3A	116.37 (17)	C3B—C2B—N3B	116.22 (17)
C1A—C2A—N3A	122.03 (17)	C1B—C2B—N3B	121.94 (17)
C4A—C3A—C2A	118.68 (18)	C4B—C3B—C2B	118.70 (18)

C4A—C3A—H3A	120.7	C4B—C3B—H3B	120.6
C2A—C3A—H3A	120.7	C2B—C3B—H3B	120.6
C3A—C4A—C5A	121.47 (18)	C3B—C4B—C5B	121.59 (18)
C3A—C4A—N4A	119.25 (19)	C3B—C4B—N4B	119.28 (18)
C5A—C4A—N4A	119.27 (19)	C5B—C4B—N4B	119.14 (19)
C6A—C5A—C4A	120.08 (18)	C6B—C5B—C4B	119.81 (19)
C6A—C5A—H5A	120.0	C6B—C5B—H5B	120.1
C4A—C5A—H5A	120.0	C4B—C5B—H5B	120.1
C5A—C6A—C1A	121.27 (18)	C5B—C6B—C1B	121.28 (18)
C5A—C6A—H6A	119.4	C5B—C6B—H6B	119.4
C1A—C6A—H6A	119.4	C1B—C6B—H6B	119.4
N2A—C7A—C8A	115.29 (19)	N2B—C7B—C8B	114.92 (18)
N2A—C7A—C14A	124.4 (2)	N2B—C7B—C14B	125.14 (19)
C8A—C7A—C14A	120.33 (19)	C8B—C7B—C14B	119.94 (17)
C9A—C8A—C13A	118.7 (2)	C13B—C8B—C9B	118.04 (19)
C9A—C8A—C7A	120.14 (17)	C13B—C8B—C7B	121.96 (19)
C13A—C8A—C7A	121.2 (2)	C9B—C8B—C7B	120.00 (17)
C10A—C9A—C8A	119.60 (19)	C10B—C9B—C8B	120.03 (18)
C10A—C9A—H9A	120.2	C10B—C9B—H9B	120.0
C8A—C9A—H9A	120.2	C8B—C9B—H9B	120.0
C9A—C10A—C11A	122.0 (2)	C11B—C10B—C9B	121.6 (2)
C9A—C10A—Cl1A	119.52 (16)	C11B—C10B—Cl1B	118.84 (18)
C11A—C10A—Cl1A	118.49 (18)	C9B—C10B—Cl1B	119.55 (16)
C12A—C11A—C10A	118.4 (2)	C12B—C11B—C10B	118.6 (2)
C12A—C11A—H11A	120.8	C12B—C11B—H11B	120.7
C10A—C11A—H11A	120.8	C10B—C11B—H11B	120.7
C11A—C12A—C13A	120.9 (2)	C11B—C12B—C13B	120.7 (2)
C11A—C12A—H12A	119.6	C11B—C12B—H12B	119.7
C13A—C12A—H12A	119.6	C13B—C12B—H12B	119.7
C12A—C13A—C8A	120.4 (2)	C12B—C13B—C8B	121.1 (2)
C12A—C13A—H13A	119.8	C12B—C13B—H13B	119.5
C8A—C13A—H13A	119.8	C8B—C13B—H13B	119.5
C7A—C14A—H14A	109.5	C7B—C14B—H14D	109.5
C7A—C14A—H14B	109.5	C7B—C14B—H14E	109.5
H14A—C14A—H14B	109.5	H14D—C14B—H14E	109.5
C7A—C14A—H14C	109.5	C7B—C14B—H14F	109.5
H14A—C14A—H14C	109.5	H14D—C14B—H14F	109.5
H14B—C14A—H14C	109.5	H14E—C14B—H14F	109.5
C1A—N1A—N2A—C7A	-177.38 (18)	C1B—N1B—N2B—C7B	176.29 (17)
N2A—N1A—C1A—C6A	4.3 (3)	N2B—N1B—C1B—C6B	2.4 (3)
N2A—N1A—C1A—C2A	-176.33 (18)	N2B—N1B—C1B—C2B	-176.75 (17)
N1A—C1A—C2A—C3A	-179.36 (18)	N1B—C1B—C2B—C3B	177.62 (18)
C6A—C1A—C2A—C3A	0.1 (3)	C6B—C1B—C2B—C3B	-1.5 (3)
N1A—C1A—C2A—N3A	1.2 (3)	N1B—C1B—C2B—N3B	-0.6 (3)
C6A—C1A—C2A—N3A	-179.35 (18)	C6B—C1B—C2B—N3B	-179.79 (17)
O2A—N3A—C2A—C3A	-1.9 (3)	O2B—N3B—C2B—C3B	3.6 (3)
O1A—N3A—C2A—C3A	176.84 (18)	O1B—N3B—C2B—C3B	-177.77 (18)

O2A—N3A—C2A—C1A	177.57 (18)	O2B—N3B—C2B—C1B	−178.10 (19)
O1A—N3A—C2A—C1A	−3.7 (3)	O1B—N3B—C2B—C1B	0.6 (3)
C1A—C2A—C3A—C4A	−0.5 (3)	C1B—C2B—C3B—C4B	−0.4 (3)
N3A—C2A—C3A—C4A	178.92 (17)	N3B—C2B—C3B—C4B	177.93 (18)
C2A—C3A—C4A—C5A	1.0 (3)	C2B—C3B—C4B—C5B	1.6 (3)
C2A—C3A—C4A—N4A	179.80 (18)	C2B—C3B—C4B—N4B	−178.42 (19)
O4A—N4A—C4A—C3A	169.6 (2)	O4B—N4B—C4B—C3B	171.1 (2)
O3A—N4A—C4A—C3A	−10.0 (3)	O3B—N4B—C4B—C3B	−8.9 (3)
O4A—N4A—C4A—C5A	−11.5 (3)	O4B—N4B—C4B—C5B	−8.9 (4)
O3A—N4A—C4A—C5A	168.8 (2)	O3B—N4B—C4B—C5B	171.1 (2)
C3A—C4A—C5A—C6A	−1.1 (3)	C3B—C4B—C5B—C6B	−0.8 (3)
N4A—C4A—C5A—C6A	−179.8 (2)	N4B—C4B—C5B—C6B	179.3 (2)
C4A—C5A—C6A—C1A	0.6 (3)	C4B—C5B—C6B—C1B	−1.3 (3)
N1A—C1A—C6A—C5A	179.35 (19)	N1B—C1B—C6B—C5B	−176.77 (19)
C2A—C1A—C6A—C5A	−0.1 (3)	C2B—C1B—C6B—C5B	2.4 (3)
N1A—N2A—C7A—C8A	−179.01 (17)	N1B—N2B—C7B—C8B	−179.42 (15)
N1A—N2A—C7A—C14A	0.8 (3)	N1B—N2B—C7B—C14B	0.5 (3)
N2A—C7A—C8A—C9A	3.7 (3)	N2B—C7B—C8B—C13B	−179.25 (18)
C14A—C7A—C8A—C9A	−176.1 (2)	C14B—C7B—C8B—C13B	0.8 (3)
N2A—C7A—C8A—C13A	−177.09 (19)	N2B—C7B—C8B—C9B	0.8 (3)
C14A—C7A—C8A—C13A	3.1 (3)	C14B—C7B—C8B—C9B	−179.11 (19)
C13A—C8A—C9A—C10A	−0.8 (3)	C13B—C8B—C9B—C10B	0.1 (3)
C7A—C8A—C9A—C10A	178.49 (18)	C7B—C8B—C9B—C10B	−179.94 (18)
C8A—C9A—C10A—C11A	0.3 (3)	C8B—C9B—C10B—C11B	0.7 (3)
C8A—C9A—C10A—C11A	−178.40 (15)	C8B—C9B—C10B—C11B	179.10 (15)
C9A—C10A—C11A—C12A	0.8 (3)	C9B—C10B—C11B—C12B	−0.9 (3)
C11A—C10A—C11A—C12A	179.51 (17)	C11B—C10B—C11B—C12B	−179.32 (17)
C10A—C11A—C12A—C13A	−1.4 (3)	C10B—C11B—C12B—C13B	0.3 (3)
C11A—C12A—C13A—C8A	0.9 (3)	C11B—C12B—C13B—C8B	0.5 (3)
C9A—C8A—C13A—C12A	0.2 (3)	C9B—C8B—C13B—C12B	−0.7 (3)
C7A—C8A—C13A—C12A	−179.1 (2)	C7B—C8B—C13B—C12B	179.35 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1A—H1NA···O1A	0.82	1.95	2.598 (2)	135
N1B—H1NB···O1B	0.86	1.86	2.589 (2)	141
C5A—H5A···O1A ⁱ	0.93	2.52	3.251 (3)	136
C5B—H5B···O1B ⁱⁱ	0.93	2.33	3.196 (3)	154
C11A—H11A···O3B ⁱⁱⁱ	0.93	2.55	3.402 (3)	153
C11B—H11B···O3A ^{iv}	0.93	2.58	3.429 (3)	153

Symmetry codes: (i) $-x+2, y-1/2, -z+3/2$; (ii) $-x+1, y+1/2, -z+3/2$; (iii) $-x+1, y-1/2, -z+1/2$; (iv) $-x+2, y+1/2, -z+5/2$.