

4,4,5,5-Tetramethyl-2-[1,3,6,8-tetra-bromo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyren-2-yl]-1,3,2-dioxaborolane

Ying Chen,^a Wen-tao Yu,^b Zhi-qiang Liu^b and Ping Yu^{a*}

^aSchool of Chemistry and Chemical Engineering, Shandong University, Jinan 250100, People's Republic of China, and ^bState Key Laboratory of Crystal Materials, Shandong University, Jinan 250100, People's Republic of China
Correspondence e-mail: yupping@sdu.edu.cn


Received 14 January 2012; accepted 11 February 2012

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.035; wR factor = 0.109; data-to-parameter ratio = 19.0.

The complete molecule of the title compound, $\text{C}_{28}\text{H}_{28}\text{B}_2\text{Br}_4\text{O}_4$, is generated by the application of a centre of inversion. In the molecule, the BO_2 plane is perpendicular to that through the pyrene ring [dihedral angle = $86.27(13)^\circ$]. In the crystal, molecules stack into columns along the b axis, the closest contact between these being of the type $\text{C}-\text{Br}\cdots\pi$.

Related literature

For background to the reactions of pyrene, see: Miura & Yamano (1995). For the structure of the non-brominated derivative, see: Coventry *et al.* (2005).



Experimental

Crystal data

$\text{C}_{28}\text{H}_{28}\text{B}_2\text{Br}_4\text{O}_4$
 $M_r = 769.76$

Monoclinic, $P2_1/c$
 $a = 15.5047(10)\text{ \AA}$

$b = 7.5136(5)\text{ \AA}$
 $c = 13.9191(9)\text{ \AA}$
 $\beta = 113.961(1)^\circ$
 $V = 1481.78(17)\text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 5.46\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.34 \times 0.24 \times 0.16\text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)
 $T_{\min} = 0.258$, $T_{\max} = 0.475$

8745 measured reflections
3344 independent reflections
2488 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.109$
 $S = 1.01$
3344 reflections

176 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.60\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.67\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the C3–C5/C8/C9/C14 benzene ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4–Br2 \cdots $Cg1^i$	1.90 (1)	3.48 (1)	4.921 (3)	130 (1)

Symmetry code: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX-2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

We gratefully acknowledge financial support from the National Natural Science Foundation of China (grant Nos 20802026 and 50803033).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5049).

References

- Bruker (2009). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Coventry, D. N., Batsanov, A. S., Goeta, A. E. H., Judith, A. K., Marder, T. B. & Perutz, R. N. (2005). *Chem. Commun.* pp. 2172–2174.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Miura, T. & Yamano, E. (1995). *J. Org. Chem.* **60**, 1070–1073.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o771 [doi:10.1107/S1600536812006095]

4,4,5,5-Tetramethyl-2-[1,3,6,8-tetrabromo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyren-2-yl]-1,3,2-dioxaborolane

Ying Chen, Wen-tao Yu, Zhi-qiang Liu and Ping Yu

S1. Comment

The chemistry of pyrene is strongly position-dependent. For example, in the bromination reaction of 2,7-di-*t*-butylpyrene, the bromide atoms were connected at the 4,5,9,10-positions of pyrene (Miura & Yamano, 1995). However, to our surprise, when the *t*-butyl group is changed to pinacol boronate, the bromination reaction resulted in bromination at the 1,3,6,8-positions as confirmed by the crystal structure described herein.

The molecule, Fig. 1, is centrosymmetric. Before bromination, the two BO₂ groups are nearly co-planar with the pyrene ring (Coventry *et al.*, 2005). However, they become nearly perpendicular after bromination (dihedral angle 86.27 (13) $^{\circ}$).

The molecules pack into columns along the *b* axis, Fig. 2. The most prominent contacts in the structure appear to be of the type C—Br \cdots π , Table 1.

S2. Experimental

The title compound was synthesized *via* a one-step bromination reaction. The precursor compound, 2,7-di-Bpinpyrene (pin = O₂C₂Me₄) was prepared using the method of Ir-catalyzed borylation (Coventry *et al.*, 2005). To a stirred mixture of 1.36 g (3.0 mmol) of 2,7-di-Bpinpyrene and a small amount of Fe powder (*ca* 0.10 g) in 80 ml of CCl₄ was added dropwise a solution of 2.88 g (0.78 mL, 18 mmol) of bromine in 20 ml of CCl₄ at room temperature. After stirring for 5 h, the mixture was slowly poured into ice water. Then the organic layer was collected and washed with aqueous Na₂S₂O₃ and dried over MgSO₄. After evaporation, the residue was crystallized from hexane, giving 1.80 g (78% yield) of gray powdered product. Crystals were grown by slow evaporation from its hexane/dichloromethane solution.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.93–0.96 Å, U_{iso}(H) 1.2–1.5U_{eq}(C)] and were included in the refinement in the riding model approximation.





Figure 1

Molecular structure of the title compound. Displacement ellipsoids are drawn at 50% probability level.

**Figure 2**

The packing in the title molecules in crystal viewed approximately down the *b* axis. Intermolecular Br···C (3.414 (3) Å) contacts, being representative of the Br···π contacts are shown as dashed lines.

4,4,5,5-Tetramethyl-2-[1,3,6,8-tetrabromo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyren-2-yl]-1,3,2-dioxaborolane

Crystal data



$M_r = 769.76$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.5047 (10)$ Å

$b = 7.5136 (5)$ Å

$c = 13.9191 (9)$ Å

$\beta = 113.961 (1)^\circ$

$V = 1481.78 (17)$ Å³

$Z = 2$

$F(000) = 756$

$D_x = 1.725 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3344 reflections

$\theta = 2.9\text{--}27.4^\circ$

$\mu = 5.46 \text{ mm}^{-1}$

$T = 296$ K

Pod, colourless

$0.34 \times 0.24 \times 0.16$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)

$T_{\min} = 0.258$, $T_{\max} = 0.475$

8745 measured reflections

3344 independent reflections

2488 reflections with $i > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -20 \rightarrow 18$

$k = -9 \rightarrow 9$

$l = -9 \rightarrow 18$

*Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.035$ $wR(F^2) = 0.109$ $S = 1.01$

3344 reflections

176 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0653P)^2 + 0.4343P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.007$ $\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.67 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.77662 (3)	0.70726 (6)	0.97202 (4)	0.07199 (18)
Br2	0.58141 (3)	0.07842 (5)	0.79828 (3)	0.06680 (17)
O1	0.73711 (17)	0.4128 (3)	0.75325 (18)	0.0521 (6)
O2	0.81831 (17)	0.2540 (4)	0.89972 (18)	0.0603 (7)
C1	0.8075 (4)	0.3192 (8)	0.6352 (3)	0.0949 (17)
H1A	0.7619	0.2248	0.6114	0.142*
H1B	0.8652	0.2812	0.6316	0.142*
H1C	0.7835	0.4218	0.5914	0.142*
C2	0.8258 (2)	0.3653 (5)	0.7461 (3)	0.0537 (9)
C3	0.6666 (2)	0.4017 (4)	0.8923 (2)	0.0380 (6)
C4	0.5917 (2)	0.2876 (4)	0.8788 (2)	0.0395 (7)
C5	0.5247 (2)	0.3199 (4)	0.9201 (2)	0.0377 (6)
C6	0.4480 (2)	0.2033 (4)	0.9060 (3)	0.0503 (8)
H6	0.4417	0.0985	0.8681	0.060*
C7	0.3844 (2)	0.2413 (5)	0.9460 (3)	0.0522 (9)
H7	0.3348	0.1628	0.9345	0.063*
C8	0.6087 (2)	0.6003 (4)	0.9941 (2)	0.0404 (7)
C9	0.6728 (2)	0.5543 (4)	0.9499 (3)	0.0424 (7)
C10	0.8662 (3)	0.2212 (5)	0.8302 (3)	0.0588 (10)
C11	0.8343 (5)	0.0358 (6)	0.7828 (6)	0.119 (2)
H11A	0.8451	-0.0486	0.8382	0.178*
H11B	0.8697	0.0015	0.7429	0.178*
H11C	0.7683	0.0387	0.7375	0.178*
C12	0.8844 (4)	0.5382 (7)	0.7748 (5)	0.0971 (17)
H12A	0.8460	0.6359	0.7364	0.146*

H12B	0.9378	0.5261	0.7571	0.146*
H12C	0.9059	0.5600	0.8489	0.146*
C13	0.9711 (3)	0.2251 (10)	0.8940 (4)	0.113 (2)
H13A	0.9882	0.3348	0.9325	0.169*
H13B	1.0032	0.2160	0.8479	0.169*
H13C	0.9889	0.1269	0.9423	0.169*
C14	0.53370 (18)	0.4799 (4)	0.9788 (2)	0.0340 (6)
B1	0.7425 (2)	0.3542 (5)	0.8473 (3)	0.0397 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0592 (3)	0.0748 (3)	0.1085 (4)	-0.02956 (19)	0.0614 (3)	-0.0371 (2)
Br2	0.0623 (3)	0.0677 (3)	0.0883 (3)	-0.01639 (18)	0.0490 (2)	-0.0388 (2)
O1	0.0482 (13)	0.0696 (15)	0.0472 (13)	0.0173 (11)	0.0281 (11)	0.0108 (11)
O2	0.0525 (14)	0.0954 (19)	0.0458 (13)	0.0283 (13)	0.0331 (12)	0.0213 (13)
C1	0.124 (4)	0.122 (4)	0.058 (3)	0.044 (4)	0.056 (3)	0.016 (3)
C2	0.053 (2)	0.070 (2)	0.0533 (19)	0.0117 (17)	0.0365 (17)	0.0064 (17)
C3	0.0304 (14)	0.0502 (17)	0.0361 (15)	0.0019 (12)	0.0164 (12)	-0.0025 (13)
C4	0.0353 (15)	0.0459 (16)	0.0396 (15)	0.0015 (12)	0.0175 (13)	-0.0087 (13)
C5	0.0303 (14)	0.0468 (16)	0.0389 (15)	-0.0028 (12)	0.0171 (13)	-0.0058 (13)
C6	0.0470 (18)	0.0480 (19)	0.066 (2)	-0.0136 (14)	0.0337 (17)	-0.0207 (16)
C7	0.0449 (18)	0.0520 (19)	0.072 (2)	-0.0169 (15)	0.0358 (18)	-0.0202 (17)
C8	0.0342 (15)	0.0481 (17)	0.0442 (16)	-0.0061 (13)	0.0213 (13)	-0.0086 (13)
C9	0.0312 (15)	0.0519 (18)	0.0493 (18)	-0.0071 (13)	0.0218 (14)	-0.0045 (14)
C10	0.054 (2)	0.073 (2)	0.070 (2)	0.0207 (18)	0.0466 (19)	0.0189 (19)
C11	0.192 (7)	0.055 (3)	0.183 (6)	0.007 (3)	0.153 (6)	0.004 (3)
C12	0.103 (4)	0.078 (3)	0.144 (5)	-0.016 (3)	0.086 (4)	-0.003 (3)
C13	0.057 (3)	0.197 (7)	0.090 (4)	0.045 (4)	0.035 (3)	0.029 (4)
C14	0.0288 (13)	0.0408 (15)	0.0354 (15)	-0.0024 (11)	0.0162 (12)	-0.0050 (12)
B1	0.0338 (17)	0.0473 (19)	0.0427 (19)	0.0011 (14)	0.0204 (15)	-0.0029 (15)

Geometric parameters (\AA , ^\circ)

Br1—C9	1.899 (3)	C6—H6	0.9300
Br2—C4	1.899 (3)	C7—C8 ⁱ	1.432 (4)
O1—B1	1.351 (4)	C7—H7	0.9300
O1—C2	1.463 (4)	C8—C9	1.408 (4)
O2—B1	1.336 (4)	C8—C14	1.419 (4)
O2—C10	1.460 (4)	C8—C7 ⁱ	1.432 (4)
C1—C2	1.492 (5)	C10—C13	1.504 (6)
C1—H1A	0.9600	C10—C11	1.535 (7)
C1—H1B	0.9600	C11—H11A	0.9600
C1—H1C	0.9600	C11—H11B	0.9600
C2—C10	1.530 (5)	C11—H11C	0.9600
C2—C12	1.542 (6)	C12—H12A	0.9600
C3—C9	1.380 (4)	C12—H12B	0.9600
C3—C4	1.394 (4)	C12—H12C	0.9600

C3—B1	1.583 (4)	C13—H13A	0.9600
C4—C5	1.397 (4)	C13—H13B	0.9600
C5—C6	1.425 (4)	C13—H13C	0.9600
C5—C14	1.428 (4)	C14—C14 ⁱ	1.426 (5)
C6—C7	1.345 (4)		
B1—O1—C2	107.2 (3)	C3—C9—Br1	116.9 (2)
B1—O2—C10	107.7 (2)	C8—C9—Br1	118.9 (2)
C2—C1—H1A	109.5	O2—C10—C13	109.0 (3)
C2—C1—H1B	109.5	O2—C10—C2	103.1 (2)
H1A—C1—H1B	109.5	C13—C10—C2	116.5 (4)
C2—C1—H1C	109.5	O2—C10—C11	106.1 (3)
H1A—C1—H1C	109.5	C13—C10—C11	110.7 (5)
H1B—C1—H1C	109.5	C2—C10—C11	110.7 (4)
O1—C2—C1	109.7 (3)	C10—C11—H11A	109.5
O1—C2—C10	103.0 (2)	C10—C11—H11B	109.5
C1—C2—C10	118.3 (4)	H11A—C11—H11B	109.5
O1—C2—C12	104.4 (3)	C10—C11—H11C	109.5
C1—C2—C12	108.0 (4)	H11A—C11—H11C	109.5
C10—C2—C12	112.5 (4)	H11B—C11—H11C	109.5
C9—C3—C4	116.6 (3)	C2—C12—H12A	109.5
C9—C3—B1	122.0 (3)	C2—C12—H12B	109.5
C4—C3—B1	121.4 (3)	H12A—C12—H12B	109.5
C3—C4—C5	123.7 (3)	C2—C12—H12C	109.5
C3—C4—Br2	117.0 (2)	H12A—C12—H12C	109.5
C5—C4—Br2	119.2 (2)	H12B—C12—H12C	109.5
C4—C5—C6	123.7 (3)	C10—C13—H13A	109.5
C4—C5—C14	117.7 (2)	C10—C13—H13B	109.5
C6—C5—C14	118.6 (2)	H13A—C13—H13B	109.5
C7—C6—C5	121.6 (3)	C10—C13—H13C	109.5
C7—C6—H6	119.2	H13A—C13—H13C	109.5
C5—C6—H6	119.2	H13B—C13—H13C	109.5
C6—C7—C8 ⁱ	121.6 (3)	C8—C14—C14 ⁱ	119.8 (3)
C6—C7—H7	119.2	C8—C14—C5	120.4 (2)
C8 ⁱ —C7—H7	119.2	C14 ⁱ —C14—C5	119.8 (3)
C9—C8—C14	117.3 (3)	O2—B1—O1	114.1 (3)
C9—C8—C7 ⁱ	124.0 (3)	O2—B1—C3	122.8 (3)
C14—C8—C7 ⁱ	118.6 (3)	O1—B1—C3	123.1 (3)
C3—C9—C8	124.2 (3)		

Symmetry code: (i) $-x+1, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the centroid of the C3—C5/C8/C9/C14 benzene ring.

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
C4—Br2 \cdots Cg1 ⁱⁱ	1.90 (1)	3.48 (1)	4.921 (3)	130 (1)

Symmetry code: (ii) $-x+1, y-1/2, -z+3/2$.