

catena-Poly[[1,10-phenanthroline)-cobalt(II)]-di- μ -azido]

Xue-Miao Gao, Jiong-Peng Zhao and Fu-Chen Liu*

School of Chemistry and Chemical Engineering, Tianjin University of Technology, Tianjin 300384, People's Republic of China
Correspondence e-mail: fuchenliutj@yahoo.com


Received 12 January 2012; accepted 13 February 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.052; wR factor = 0.101; data-to-parameter ratio = 14.9.

In the crystal structure of the binuclear title complex, $[\text{Co}(\text{N}_3)_2(\text{C}_{12}\text{H}_8\text{N}_2)]_n$, each Co^{II} cation is coordinated by two N atoms from one chelating 1,10-phenanthroline ligand and four azide ligands in a slightly distorted octahedral coordination. The two Co^{II} cations of the binuclear complex are related by an inversion centre and are bridged by two symmetry-related azide ligands in both $\mu_{1,1}$ and $\mu_{1,3}$ modes. The $\mu_{1,3}$ bridging mode gives rise to an infinite one-dimensional chain along the a axis, whereas the $\mu_{1,1}$ bridging mode is responsible for the formation of the binuclear Co^{II} complex.

Related literature

For general background to metal–azide complexes, see: Zhao *et al.* (2009). For a closely related Ni–azide structure, see: Li *et al.* (2000).



Experimental

Crystal data

$[\text{Co}(\text{N}_3)_2(\text{C}_{12}\text{H}_8\text{N}_2)]$	$\gamma = 105.78 (3)^\circ$
$M_r = 323.19$	$V = 622.8 (2)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.0018 (14)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.049 (2)\text{ \AA}$	$\mu = 1.38\text{ mm}^{-1}$
$c = 10.491 (2)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 109.83 (3)^\circ$	$0.2 \times 0.18 \times 0.18\text{ mm}$
$\beta = 103.63 (3)^\circ$	

Data collection

Rigaku SCXmini diffractometer	2822 independent reflections
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	2087 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.720$, $T_{\max} = 1$	$R_{\text{int}} = 0.053$
6534 measured reflections	Standard reflections: 0

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	190 parameters
$wR(F^2) = 0.101$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.36\text{ e \AA}^{-3}$
2822 reflections	$\Delta\rho_{\min} = -0.42\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

$\text{Co1}-\text{N}1^{\text{i}}$	2.113 (3)	$\text{Co1}-\text{N}6^{\text{ii}}$	2.144 (3)
$\text{Co1}-\text{N}1$	2.175 (3)	$\text{Co1}-\text{N}7$	2.141 (3)
$\text{Co1}-\text{N}4$	2.202 (3)	$\text{Co1}-\text{N}8$	2.141 (3)

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $-x + 3, -y + 2, -z + 1$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *PROCESS-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors acknowledge financial support from Tianjin Municipal Education Commission (grant No. 20100502).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VN2030).

References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Li, L.-C., Liao, D.-Z., Jiang, Z.-H. & Yan, S.-P. (2000). *Polyhedron*, **19**, 1575–1578.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Americas Corporation, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Zhao, J.-P., Hu, B.-W., Sanudo, E. C., Yang, Q., Zeng, Y.-F. & Bu, X.-H. (2009). *Inorg. Chem.* **48**, 2482–2489.

supporting information

Acta Cryst. (2012). E68, m318 [doi:10.1107/S1600536812006435]

catena-Poly[[*(1,10-phenanthroline)cobalt(II)*]-di- μ -azido]

Xue-Miao Gao, Jiong-Peng Zhao and Fu-Chen Liu

S1. Comment


Some metal-azido complexes with unique structural features have been reported in the past (Zhao *et al.*, 2009). Co-ligands are often encountered in metal azido complex, e.g. the 1,10-phenanthroline ligand which are used frequently in assembling metal azido complexes. One 1D nickel-azido complex with 1,10-phenanthroline as co-ligand was reported in 2000 (Li *et al.*, 2000). However, its isomorphic Co^{II} compound was not reported until this work. That may be due to the fact that Co^{II} cations are easy oxidized to Co^{III} with 1,10-phenanthroline as co-ligand. In the title complex the Co^{II} ion is coordinated by one chelating 1,10-phenanthroline and four azido anions forming a distorted CoN₆ octahedral environment (Fig. 1). The azido anions take two different coordinated types, in which one bridges two Co^{II} ions in $\mu_{1,1}$ mode while the other one bridges two Co^{II} ions in $\mu_{1,3}$ mode. The two $\mu_{1,1}$ azido anions link two Co^{II} ions forming a binuclear dimer whereas the dimers linked by the two $\mu_{1,3}$ azido anions yield a 1D chain of bimuclear Co^{II} complexes (Fig. 2). π – π stacking of the phenanthroline aromatic rings between adjacent chains assists in the forming of a 3D supermolecular structure (Fig. 3). The smallest centroid to centroid distances between the aromatic rings of phenanthroline aromatic rings between adjacent chains are 3.589 (2) Å and 3.605 (2) Å, respectively. A weak CH···N interaction is present between the aromatic H5 proton and the terminal N3 azide nitrogen (2.54 Å), reinforcing slightly the packing of adjacent chains.

S2. Experimental


The complex was hydrothermally synthesized under auto-generated pressure. A mixture of cobalt formate (1 mmol), NaN₃ (1 mmol) and 1,10-phenanthroline (1 mmol) in methanol was sealed in a Teflon-lined stainless-steel Parr bomb that was heated at 413 K for 48 h. Red crystals of the title complex were collected after the bomb was allowed to cool to room temperature. Yield 20% based on metal salt.

S3. Refinement


H atoms were included in calculated positions and treated as riding on their parent C atoms with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

The coordinated mode and linkage of the complex. Atomic displacement ellipsoids are drawn at the 30% probability level. Symmetry codes: (i) $2 - x, 2 - y, 1 - z$; (ii) $3 - x, 2 - y, 1 - z$.

**Figure 2**


The 1D chain of the complex along the a -axis.

**Figure 3**

Packing plot of the title compound. The dotted lines indicate the weak hydrogen bonds between the azide anions and ring H atoms of adjacent chains

catena-Poly[[1,10-phenanthroline)cobalt(II)]-di- μ -azido]

Crystal data



$$M_r = 323.19$$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$$a = 7.0018 (14) \text{ \AA}$$

$$b = 10.049 (2) \text{ \AA}$$

$$c = 10.491 (2) \text{ \AA}$$

$$\alpha = 109.83 (3)^\circ$$

$$\beta = 103.63 (3)^\circ$$

$$\gamma = 105.78 (3)^\circ$$

$$V = 622.8 (2) \text{ \AA}^3$$

$$Z = 2$$

$$F(000) = 326$$

$$D_x = 1.723 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5775 reflections

$$\theta = 3.2\text{--}27.5^\circ$$

$$\mu = 1.38 \text{ mm}^{-1}$$

$T = 293\text{ K}$

Block, red

*Data collection*Rigaku SCXmini
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scansAbsorption correction: multi-scan
(*ABSCOR*; Higashi, 1995) $T_{\min} = 0.720$, $T_{\max} = 1$ $0.2 \times 0.18 \times 0.18\text{ mm}$

6534 measured reflections

2822 independent reflections

2087 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.053$ $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.2^\circ$ $h = -9 \rightarrow 9$ $k = -13 \rightarrow 13$ $l = -13 \rightarrow 13$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.052$ $wR(F^2) = 0.101$ $S = 1.07$

2822 reflections

190 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0411P)^2 + 0.0273P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.36\text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.42\text{ e \AA}^{-3}$ *Special details*

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$
Co1	1.08955 (7)	0.88775 (5)	0.39668 (4)	0.02424 (16)
N1	0.8499 (4)	0.9870 (3)	0.3808 (3)	0.0290 (6)
N2	0.7988 (4)	1.0243 (3)	0.2840 (3)	0.0321 (7)
N3	0.7510 (6)	1.0590 (4)	0.1905 (4)	0.0599 (11)
N4	1.3416 (5)	0.7993 (3)	0.4387 (3)	0.0357 (7)
N5	1.5146 (5)	0.8649 (3)	0.5268 (3)	0.0264 (6)
N6	1.6882 (5)	0.9262 (3)	0.6149 (3)	0.0362 (7)
N7	0.9662 (4)	0.7562 (3)	0.1657 (3)	0.0249 (6)
N8	0.8808 (4)	0.6720 (3)	0.3697 (3)	0.0249 (6)
C1	1.0005 (6)	0.8017 (4)	0.0652 (4)	0.0338 (8)
H1A	1.0859	0.9035	0.0945	0.041*
C2	0.9122 (6)	0.7015 (5)	-0.0844 (4)	0.0410 (9)
H2A	0.9373	0.7376	-0.1521	0.049*
C3	0.7906 (6)	0.5522 (4)	-0.1286 (4)	0.0395 (9)

H3A	0.7343	0.4850	-0.2269	0.047*
C4	0.6230 (5)	0.3430 (4)	-0.0625 (4)	0.0375 (9)
H4A	0.5630	0.2713	-0.1593	0.045*
C5	0.5901 (5)	0.2991 (4)	0.0411 (4)	0.0374 (9)
H5A	0.5115	0.1968	0.0150	0.045*
C6	0.6387 (5)	0.3701 (4)	0.3039 (4)	0.0363 (9)
H6A	0.5605	0.2695	0.2834	0.044*
C7	0.7191 (6)	0.4816 (4)	0.4423 (4)	0.0371 (9)
H7A	0.6937	0.4580	0.5167	0.045*
C8	0.8392 (5)	0.6306 (4)	0.4720 (4)	0.0324 (8)
H8A	0.8933	0.7053	0.5673	0.039*
C9	0.7490 (5)	0.4984 (4)	-0.0256 (3)	0.0289 (8)
C10	0.6746 (5)	0.4079 (4)	0.1915 (4)	0.0285 (7)
C11	0.8398 (5)	0.6068 (3)	0.1210 (3)	0.0235 (7)
C12	0.7988 (5)	0.5611 (3)	0.2306 (3)	0.0230 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0244 (3)	0.0210 (2)	0.0226 (3)	0.00471 (18)	0.00743 (18)	0.00793 (19)
N1	0.0317 (16)	0.0289 (15)	0.0245 (15)	0.0127 (13)	0.0077 (13)	0.0103 (13)
N2	0.0352 (18)	0.0192 (14)	0.0265 (16)	0.0055 (13)	0.0027 (14)	0.0020 (13)
N3	0.089 (3)	0.039 (2)	0.038 (2)	0.021 (2)	0.0026 (19)	0.0190 (17)
N4	0.0317 (18)	0.0273 (16)	0.0434 (18)	0.0082 (14)	0.0105 (15)	0.0146 (14)
N5	0.0326 (18)	0.0191 (14)	0.0332 (17)	0.0100 (13)	0.0167 (15)	0.0145 (13)
N6	0.0289 (17)	0.0355 (17)	0.0396 (18)	0.0050 (14)	0.0083 (15)	0.0195 (15)
N7	0.0248 (15)	0.0268 (15)	0.0252 (14)	0.0098 (12)	0.0112 (12)	0.0124 (12)
N8	0.0276 (15)	0.0216 (14)	0.0210 (14)	0.0072 (12)	0.0060 (12)	0.0081 (12)
C1	0.035 (2)	0.042 (2)	0.033 (2)	0.0161 (17)	0.0165 (17)	0.0215 (18)
C2	0.044 (2)	0.063 (3)	0.033 (2)	0.029 (2)	0.0211 (18)	0.027 (2)
C3	0.038 (2)	0.049 (2)	0.0268 (19)	0.0207 (19)	0.0093 (17)	0.0091 (18)
C4	0.033 (2)	0.033 (2)	0.030 (2)	0.0162 (17)	0.0035 (17)	-0.0019 (17)
C5	0.032 (2)	0.0247 (18)	0.042 (2)	0.0091 (16)	0.0064 (17)	0.0047 (17)
C6	0.032 (2)	0.0249 (18)	0.049 (2)	0.0061 (16)	0.0136 (18)	0.0165 (18)
C7	0.044 (2)	0.035 (2)	0.040 (2)	0.0110 (17)	0.0211 (18)	0.0239 (18)
C8	0.035 (2)	0.0322 (19)	0.0281 (18)	0.0114 (16)	0.0095 (16)	0.0130 (16)
C9	0.0287 (19)	0.0333 (19)	0.0232 (17)	0.0181 (16)	0.0068 (15)	0.0075 (15)
C10	0.0244 (18)	0.0258 (17)	0.0332 (19)	0.0127 (14)	0.0080 (15)	0.0096 (15)
C11	0.0223 (17)	0.0235 (17)	0.0224 (17)	0.0092 (14)	0.0071 (14)	0.0078 (14)
C12	0.0179 (16)	0.0211 (16)	0.0265 (17)	0.0074 (13)	0.0059 (14)	0.0080 (14)

Geometric parameters (\AA , $^\circ$)

Co1—N1 ⁱ	2.113 (3)	C2—C3	1.356 (5)
Co1—N1	2.175 (3)	C2—H2A	0.9300
Co1—N4	2.202 (3)	C3—C9	1.416 (5)
Co1—N6 ⁱⁱ	2.144 (3)	C3—H3A	0.9300
Co1—N7	2.141 (3)	C4—C5	1.347 (5)

Co1—N8	2.141 (3)	C4—C9	1.431 (5)
N1—N2	1.209 (4)	C4—H4A	0.9300
N1—Co1 ⁱ	2.113 (3)	C5—C10	1.440 (5)
N2—N3	1.155 (4)	C5—H5A	0.9300
N4—N5	1.175 (4)	C6—C7	1.359 (5)
N5—N6	1.178 (4)	C6—C10	1.411 (5)
N6—Co1 ⁱⁱ	2.144 (3)	C6—H6A	0.9300
N7—C1	1.329 (4)	C7—C8	1.386 (5)
N7—C11	1.364 (4)	C7—H7A	0.9300
N8—C8	1.339 (4)	C8—H8A	0.9300
N8—C12	1.362 (4)	C9—C11	1.408 (4)
C1—C2	1.411 (5)	C10—C12	1.403 (4)
C1—H1A	0.9300	C11—C12	1.434 (4)
N1 ⁱ —Co1—N8	97.49 (11)	C3—C2—H2A	120.3
N1 ⁱ —Co1—N6 ⁱⁱ	93.69 (12)	C1—C2—H2A	120.3
N8—Co1—N6 ⁱⁱ	167.62 (10)	C2—C3—C9	120.1 (3)
N1 ⁱ —Co1—N7	168.99 (10)	C2—C3—H3A	119.9
N8—Co1—N7	77.67 (10)	C9—C3—H3A	119.9
N6 ⁱⁱ —Co1—N7	92.20 (11)	C5—C4—C9	120.9 (3)
N1 ⁱ —Co1—N1	79.52 (11)	C5—C4—H4A	119.5
N8—Co1—N1	95.55 (10)	C9—C4—H4A	119.5
N6 ⁱⁱ —Co1—N1	91.67 (11)	C4—C5—C10	121.1 (3)
N7—Co1—N1	91.02 (10)	C4—C5—H5A	119.5
N1 ⁱ —Co1—N4	94.21 (11)	C10—C5—H5A	119.5
N8—Co1—N4	84.91 (10)	C7—C6—C10	119.7 (3)
N6 ⁱⁱ —Co1—N4	89.00 (11)	C7—C6—H6A	120.2
N7—Co1—N4	95.19 (11)	C10—C6—H6A	120.2
N1—Co1—N4	173.73 (10)	C6—C7—C8	119.7 (3)
N2—N1—Co1 ⁱ	127.0 (2)	C6—C7—H7A	120.2
N2—N1—Co1	121.2 (2)	C8—C7—H7A	120.2
Co1 ⁱ —N1—Co1	100.48 (11)	N8—C8—C7	123.0 (3)
N3—N2—N1	179.2 (4)	N8—C8—H8A	118.5
N5—N4—Co1	128.3 (2)	C7—C8—H8A	118.5
N4—N5—N6	177.8 (3)	C11—C9—C3	116.7 (3)
N5—N6—Co1 ⁱⁱ	118.0 (2)	C11—C9—C4	119.5 (3)
C1—N7—C11	118.1 (3)	C3—C9—C4	123.9 (3)
C1—N7—Co1	128.3 (2)	C12—C10—C6	117.2 (3)
C11—N7—Co1	113.60 (19)	C12—C10—C5	119.1 (3)
C8—N8—C12	117.6 (3)	C6—C10—C5	123.7 (3)
C8—N8—Co1	128.5 (2)	N7—C11—C9	123.1 (3)
C12—N8—Co1	113.6 (2)	N7—C11—C12	117.3 (3)
N7—C1—C2	122.5 (3)	C9—C11—C12	119.6 (3)
N7—C1—H1A	118.8	N8—C12—C10	122.9 (3)
C2—C1—H1A	118.8	N8—C12—C11	117.3 (3)
C3—C2—C1	119.4 (3)	C10—C12—C11	119.8 (3)
N1 ⁱ —Co1—N1—N2	145.8 (3)	C1—C2—C3—C9	1.4 (5)

N8—Co1—N1—N2	−117.6 (2)	C9—C4—C5—C10	2.1 (5)
N6 ⁱⁱ —Co1—N1—N2	52.4 (3)	C10—C6—C7—C8	−1.3 (5)
N7—Co1—N1—N2	−39.9 (3)	C12—N8—C8—C7	0.4 (5)
N1 ⁱ —Co1—N1—Co1 ⁱ	0.0	Co1—N8—C8—C7	−172.4 (2)
N8—Co1—N1—Co1 ⁱ	96.62 (12)	C6—C7—C8—N8	0.2 (5)
N6 ⁱⁱ —Co1—N1—Co1 ⁱ	−93.45 (12)	C2—C3—C9—C11	−0.1 (5)
N7—Co1—N1—Co1 ⁱ	174.32 (11)	C2—C3—C9—C4	−179.9 (3)
N1 ⁱ —Co1—N4—N5	−41.8 (3)	C5—C4—C9—C11	0.0 (5)
N8—Co1—N4—N5	−139.0 (3)	C5—C4—C9—C3	179.8 (3)
N6 ⁱⁱ —Co1—N4—N5	51.8 (3)	C7—C6—C10—C12	1.7 (5)
N7—Co1—N4—N5	143.9 (3)	C7—C6—C10—C5	−177.8 (3)
N1 ⁱ —Co1—N7—C1	111.0 (5)	C4—C5—C10—C12	−2.1 (5)
N8—Co1—N7—C1	175.8 (3)	C4—C5—C10—C6	177.5 (3)
N6 ⁱⁱ —Co1—N7—C1	−11.4 (3)	C1—N7—C11—C9	1.7 (5)
N1—Co1—N7—C1	80.4 (3)	Co1—N7—C11—C9	−177.4 (2)
N4—Co1—N7—C1	−100.5 (3)	C1—N7—C11—C12	−177.9 (3)
N1 ⁱ —Co1—N7—C11	−70.1 (6)	Co1—N7—C11—C12	3.0 (3)
N8—Co1—N7—C11	−5.2 (2)	C3—C9—C11—N7	−1.5 (5)
N6 ⁱⁱ —Co1—N7—C11	167.6 (2)	C4—C9—C11—N7	178.3 (3)
N1—Co1—N7—C11	−100.7 (2)	C3—C9—C11—C12	178.1 (3)
N4—Co1—N7—C11	78.4 (2)	C4—C9—C11—C12	−2.1 (5)
N1 ⁱ —Co1—N8—C8	−10.2 (3)	C8—N8—C12—C10	0.0 (4)
N6 ⁱⁱ —Co1—N8—C8	144.2 (5)	Co1—N8—C12—C10	173.9 (2)
N7—Co1—N8—C8	179.9 (3)	C8—N8—C12—C11	178.6 (3)
N1—Co1—N8—C8	−90.3 (3)	Co1—N8—C12—C11	−7.6 (3)
N4—Co1—N8—C8	83.4 (3)	C6—C10—C12—N8	−1.1 (5)
N1 ⁱ —Co1—N8—C12	176.8 (2)	C5—C10—C12—N8	178.5 (3)
N6 ⁱⁱ —Co1—N8—C12	−28.8 (6)	C6—C10—C12—C11	−179.6 (3)
N7—Co1—N8—C12	6.8 (2)	C5—C10—C12—C11	0.0 (5)
N1—Co1—N8—C12	96.6 (2)	N7—C11—C12—N8	3.1 (4)
N4—Co1—N8—C12	−89.6 (2)	C9—C11—C12—N8	−176.5 (3)
C11—N7—C1—C2	−0.3 (5)	N7—C11—C12—C10	−178.3 (3)
Co1—N7—C1—C2	178.6 (2)	C9—C11—C12—C10	2.1 (5)
N7—C1—C2—C3	−1.2 (5)		

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x+3, -y+2, -z+1$.