

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Bromobenzoic acid–6-(4-bromophenyl)-3-methyl-1,2,4-triazolo-[3,4-*b*][1,3,4]thiadiazole (1/1)

 Kamini Kapoor,^a Vivek K. Gupta,^a Satya Paul,^b Seema Sahi^b and Rajni Kant^{a*}

^aX-ray Crystallography Laboratory, Post-Graduate Department of Physics & Electronics, University of Jammu, Jammu Tawi 180 006, India, and ^bDepartment of Chemistry, University of Jammu, Jammu Tawi 180 006, India
Correspondence e-mail: rkvk.paper11@gmail.com


Received 5 March 2012; accepted 21 March 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.048; wR factor = 0.116; data-to-parameter ratio = 15.2.

In the title 1:1 co-crystal, $\text{C}_{10}\text{H}_7\text{BrN}_4\text{S}\cdot\text{C}_7\text{H}_5\text{BrO}_2$, the triazolothiadiazole system is approximately planar [with a maximum deviation of 0.030 (4) Å] and forms a dihedral angle of 8.6 (1)° with the bromophenyl ring. In the carboxylic acid molecule, the carboxyl group is rotated by 6.4 (3)° out of the benzene ring plane. The crystal structure features O—H···N and C—H···O hydrogen bonds, π – π stacking interactions [centroid–centroid distances = 3.713 (2), 3.670 (2) and 3.859 (3) Å] and short S···N [2.883 (4) Å] contacts.

Related literature

For the biological activity of triazole derivatives, thiadiazoles and triazolothiadiazole compounds, see: Chaturvedi *et al.* (1988); Holla *et al.* (2003); Bhat *et al.* (2004); Bekircan & Bektas (2006); Shawali & Sayed (2006); Mathew *et al.* (2007); Karthikeyan *et al.* (2007); Zhou *et al.* (2007). For related structures, see: Dinçer *et al.* (2005); Arshad *et al.* (2009); Jia *et al.* (2011). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{10}\text{H}_7\text{BrN}_4\text{S}\cdot\text{C}_7\text{H}_5\text{BrO}_2$
 $M_r = 496.19$
 Triclinic, $P\bar{1}$
 $a = 7.7592$ (3) Å
 $b = 8.0634$ (4) Å
 $c = 14.9076$ (7) Å
 $\alpha = 94.090$ (4)°
 $\beta = 92.961$ (3)°
 $\gamma = 99.326$ (4)°
 $V = 916.13$ (7) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 4.56$ mm⁻¹
 $T = 293$ K
 $0.3 \times 0.2 \times 0.2$ mm

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer
 Absorption correction: multi-scan (*CrysAlis PRO RED*; Oxford Diffraction, 2010)
 $T_{\min} = 0.581$, $T_{\max} = 1.000$
 8264 measured reflections
 3594 independent reflections
 2254 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.116$
 $S = 1.01$
 3594 reflections
 236 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.43$ e Å⁻³
 $\Delta\rho_{\min} = -0.51$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O24—H24···N2 ⁱ	0.82	1.87	2.674 (4)	169
C9—H9A···O23 ⁱ	0.96	2.48	3.393 (6)	159

 Symmetry code: (i) $-x - 1, -y, -z + 2$.

Data collection: *CrysAlis PRO CCD* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO CCD*; data reduction: *CrysAlis PRO RED* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009).

RK acknowledges the Department of Science & Technology for the single-crystal X-ray diffractometer sanctioned as a National Facility under project No. SR/S2/CMP-47/2003. He is also thankful to the University of Jammu, Jammu, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2421).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Arshad, M. N., Tahir, M. N., Khan, I. U., Shafiq, M. & Waheed, A. (2009). *Acta Cryst. E* **65**, o640.
 Bekircan, O. & Bektas, H. (2006). *Molecules*, **11**, 469–477.
 Bhat, K. S., Prasad, D. J., Poojary, B. & Holla, B. S. (2004). *Phosphorus Sulfur Silicon Relat. Elem.* **179**, 1595–1603.
 Chaturvedi, B., Tiwari, N. & Nizamuddin (1988). *Agric. Biol. Chem.* **52**, 1229–1232.
 Dinçer, M., Özdemir, N., Çetin, A., Cansız, A. & Büyükgüngör, O. (2005). *Acta Cryst. C* **61**, o665–o667.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.

- Holla, B. S., Veerendra, B., Shivananda, M. K. & Poojary, B. (2003). *Eur. J. Med. Chem.* **38**, 759–767.
- Jia, W., Wang, Z., Jia, X., Zhang, J. & Wang, W. (2011). *Acta Cryst.* **E67**, o1093.
- Karthikeyan, M. S., Holla, B. S., Kalluraya, B. & Kumari, N. S. (2007). *Monatsh. Chem.* **138**, 1309–1316.
- Mathew, V., Keshavayya, J., Vaidya, V. P. & Giles, D. (2007). *Eur. J. Med. Chem.* **42**, 823–840.
- Oxford Diffraction (2010). *CrysAlis PRO CCD* and *CrysAlis PRO RED*. Oxford Diffraction Ltd, Yarnton, England.
- Shawali, A. S. & Sayed, A. R. (2006). *J. Sulfur Chem.* **27**, 233–244.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Zhou, S., Zhang, L., Jin, J., Zhang, A., Lei, X., Lin, J., He, J. & Zhang, H. (2007). *Phosphorus Sulfur Silicon Relat. Elem.* **182**, 419–432.

supporting information

Acta Cryst. (2012). E68, o1185–o1186 [https://doi.org/10.1107/S1600536812012184]

4-Bromobenzoic acid–6-(4-bromophenyl)-3-methyl-1,2,4-triazolo[3,4-*b*] [1,3,4]thiadiazole (1/1)

Kamini Kapoor, Vivek K. Gupta, Satya Paul, Seema Sahi and Rajni Kant

S1. Comment

Derivatives of 1,2,4-triazole possess a wide spectrum of biological activity, such as anticancer, anticonvulsant, analgesic, antibacterial, anthelmintic, antitubercular and anti-inflammatory activities (Holla *et al.*, 2003; Bekircan & Bektas, 2006; Zhou *et al.*, 2007). Similarly 1,3,4-thiadiazoles were also found to possess antitumor, anti-inflammatory, antibacterial, antifungal, anticonvulsant and antitubercular properties (Bhat *et al.*, 2004; Mathew *et al.*, 2007). Thus triazolothiadiazole systems may be viewed as cyclic analogues of two very important components, which often display diverse pharmacological properties. Triazolothiadiazoles obtained by fusing the 1,2,4-triazole and 1,3,4-thiadiazole rings together have been reported to possess similar biological properties (Chaturvedi *et al.*, 1988; Shawali & Sayed, 2006; Karthikeyan *et al.*, 2007). Here we report the crystal structure of the 1:1 cocrystal of a triazolothiadiazole derivative and 4-bromobenzoic acid.

Bond lengths (Allen *et al.*, 1987) and angles in the title compound (Fig. 1) have normal values and also correspond to those observed in related structures (Dinçer *et al.*, 2005; Arshad *et al.*, 2009; Jia *et al.*, 2011). The triazolothiadiazole ring is planar with a maximum deviation of 0.030 (4) Å for atom C6. The plane through the benzene ring forms dihedral angle of 8.6 (1)° with the triazolothiadiazole unit. In the molecular structure, an intramolecular C15—H15⋯S7 contact leads to the formation of a five-membered ring which is fused with the phenyl ring (Fig. 1).

In the crystal structure of the title compound, intermolecular O—H⋯N and C—H⋯O hydrogen bonds (Table 2) link the triazolothiadiazole molecule with 4-bromobenzoic acid (Fig. 2). In addition to these interactions, the crystal structure contains three π – π stacking interactions. The first of these is between the thiadiazole ring and its symmetry-related partner at ($-x$, $1-y$, $-z$), with a distance of 3.713 (2) Å between the ring centroids, and a perpendicular distance between the rings of 3.468 Å. The second is between the triazole ring and the benzene ring at ($-x$, $1-y$, $-z$), with a distance of 3.670 (2) Å between the ring centroids and a perpendicular distance between the rings of 3.427 Å. The third is between the benzene rings (C10⋯C15) and (C16⋯C21) in the asymmetric unit, with a distance of 3.859 (3) Å between the ring centroids and a perpendicular distance between the rings of 3.599 Å. A short contact distance not listed in tables, yet noteworthy, is S7⋯N1 with N1 at position ($-x-1$, $-y+1$, $-z+2$), the S⋯N separation being 2.883 (4) Å, which may cause steric hindrance.

S2. Experimental

4-Amino-5-mercapto-3-methyl-1,2,4-triazole (0.130 g, 1 mmol) and 4-bromo- β -chlorocinnamic acid (0.261 g, 1 mmol) were stirred in POCl₃ (3 ml) at 80 °C for 30 min. 6-(4-Bromophenyl)-3-methyl-1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazole was obtained along with 4-bromobenzoic acid after pouring the reaction mixture in crushed ice followed by washing with dilute NaOH. Finally, it was crystallized from methanol, affording white crystals.

S3. Refinement

All H atoms were positioned geometrically and were treated as riding on their parent atoms, with O—H = 0.82 Å for OH, C—H = 0.93 Å for aromatic H, C—H = 0.96 Å for methyl H, and with $U_{\text{iso}}(\text{H}_{\text{aryl}}) = 1.2U_{\text{eq}}(\text{C}_{\text{aryl}})$, $U_{\text{iso}}(\text{H}_{\text{methyl}}) = 1.5U_{\text{eq}}(\text{methyl C})$, and $U_{\text{iso}}(\text{H24}) = 1.5U_{\text{eq}}(\text{O24})$.




Figure 1

ORTEP view of the asymmetric unit of the title cocrystal, with thermal ellipsoids drawn at the 40% probability level. H atoms are shown as small spheres of arbitrary radii.




Figure 2

The packing arrangement of molecules viewed down the a axis. The broken lines show the intermolecular O—H \cdots N and C—H \cdots O interactions.

4-Bromobenzoic acid–6-(4-bromophenyl)-3-methyl-1,2,4- triazolo[3,4- b][1,3,4]thiadiazole (1/1)

Crystal data

$C_{10}H_7BrN_4S \cdot C_7H_5BrO_2$

$M_r = 496.19$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.7592$ (3) Å

$b = 8.0634$ (4) Å

$c = 14.9076$ (7) Å

$\alpha = 94.090$ (4)°

$\beta = 92.961$ (3)°

$\gamma = 99.326$ (4)°

$V = 916.13$ (7) Å³

$Z = 2$

$F(000) = 488$

$D_x = 1.799$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2731 reflections

$\theta = 3.4$ – 28.9 °

$\mu = 4.56$ mm⁻¹

$T = 293$ K $0.3 \times 0.2 \times 0.2$ mm
 Block, white

Data collection

Oxford Diffraction Xcalibur Sapphire3 diffractometer	8264 measured reflections
Radiation source: fine-focus sealed tube	3594 independent reflections
Graphite monochromator	2254 reflections with $I > 2\sigma(I)$
Detector resolution: 16.1049 pixels mm ⁻¹	$R_{\text{int}} = 0.036$
ω scans	$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 3.4^\circ$
Absorption correction: multi-scan (<i>CrysAlis PRO RED</i> ; Oxford Diffraction, 2010)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.581$, $T_{\text{max}} = 1.000$	$k = -9 \rightarrow 9$
	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.116$	$w = 1/[\sigma^2(F_o^2) + (0.0424P)^2 + 0.2229P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
3594 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
236 parameters	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.51 \text{ e \AA}^{-3}$
0 constraints	
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.48693 (7)	0.65776 (8)	0.62657 (4)	0.0850 (3)
Br2	0.15874 (8)	0.27859 (8)	0.50571 (4)	0.0967 (3)
N1	-0.4623 (4)	0.3387 (4)	1.0195 (2)	0.0462 (9)
N2	-0.4261 (4)	0.1982 (4)	1.0619 (2)	0.0445 (9)
C3	-0.2777 (5)	0.1581 (5)	1.0362 (3)	0.0397 (10)
N4	-0.2150 (4)	0.2704 (4)	0.9772 (2)	0.0352 (8)
N5	-0.0694 (4)	0.2926 (4)	0.9277 (2)	0.0375 (8)
C6	-0.0778 (5)	0.4209 (5)	0.8811 (3)	0.0340 (9)
S7	-0.25764 (12)	0.52658 (14)	0.89676 (7)	0.0435 (3)
C8	-0.3311 (5)	0.3782 (5)	0.9699 (3)	0.0360 (9)
C9	-0.1926 (5)	0.0180 (5)	1.0658 (3)	0.0554 (13)
H9A	-0.2529	-0.0311	1.1146	0.083*
H9B	-0.0728	0.0603	1.0855	0.083*
H9C	-0.1970	-0.0661	1.0164	0.083*
C10	0.0567 (5)	0.4805 (5)	0.8205 (3)	0.0361 (9)
C11	0.2088 (5)	0.4096 (5)	0.8184 (3)	0.0440 (10)
H11	0.2251	0.3262	0.8565	0.053*
C12	0.3355 (5)	0.4618 (6)	0.7602 (3)	0.0527 (12)
H12	0.4359	0.4128	0.7583	0.063*
C13	0.3124 (5)	0.5858 (6)	0.7056 (3)	0.0480 (11)
C14	0.1622 (6)	0.6563 (6)	0.7054 (3)	0.0657 (14)
H14	0.1465	0.7390	0.6667	0.079*

C15	0.0365 (6)	0.6033 (6)	0.7628 (3)	0.0547 (12)
H15	-0.0648	0.6511	0.7629	0.066*
C16	-0.2498 (6)	0.0900 (5)	0.7076 (3)	0.0474 (11)
C17	-0.2908 (7)	0.1880 (7)	0.6397 (3)	0.0694 (15)
H17	-0.4023	0.2159	0.6342	0.083*
C18	-0.1704 (8)	0.2442 (7)	0.5808 (4)	0.0782 (16)
H18	-0.1989	0.3108	0.5357	0.094*
C19	-0.0075 (7)	0.2015 (6)	0.5886 (3)	0.0596 (13)
C20	0.0382 (6)	0.1051 (6)	0.6543 (3)	0.0614 (13)
H20	0.1500	0.0776	0.6590	0.074*
C21	-0.0852 (6)	0.0486 (6)	0.7140 (3)	0.0557 (12)
H21	-0.0560	-0.0181	0.7590	0.067*
C22	-0.3841 (6)	0.0335 (6)	0.7719 (3)	0.0541 (12)
O23	-0.5243 (5)	0.0796 (5)	0.7726 (3)	0.0950 (13)
O24	-0.3345 (4)	-0.0720 (4)	0.8274 (2)	0.0619 (9)
H24	-0.4124	-0.0992	0.8612	0.093*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0729 (4)	0.1007 (5)	0.0763 (4)	-0.0147 (3)	0.0444 (3)	0.0090 (3)
Br2	0.1079 (5)	0.0963 (5)	0.0782 (5)	-0.0219 (4)	0.0536 (4)	0.0087 (3)
N1	0.0459 (19)	0.044 (2)	0.056 (2)	0.0171 (16)	0.0223 (17)	0.0206 (18)
N2	0.0456 (19)	0.044 (2)	0.049 (2)	0.0128 (16)	0.0183 (16)	0.0164 (17)
C3	0.044 (2)	0.041 (3)	0.039 (2)	0.0135 (19)	0.0142 (19)	0.0119 (19)
N4	0.0386 (17)	0.0350 (19)	0.0361 (19)	0.0122 (15)	0.0118 (14)	0.0096 (15)
N5	0.0347 (17)	0.041 (2)	0.041 (2)	0.0145 (15)	0.0147 (14)	0.0099 (16)
C6	0.036 (2)	0.033 (2)	0.034 (2)	0.0071 (17)	0.0065 (17)	0.0032 (18)
S7	0.0403 (5)	0.0434 (7)	0.0537 (7)	0.0157 (5)	0.0182 (5)	0.0205 (5)
C8	0.037 (2)	0.039 (2)	0.038 (2)	0.0161 (18)	0.0110 (18)	0.0093 (18)
C9	0.058 (3)	0.054 (3)	0.065 (3)	0.026 (2)	0.025 (2)	0.032 (2)
C10	0.034 (2)	0.039 (2)	0.036 (2)	0.0052 (17)	0.0076 (17)	0.0027 (18)
C11	0.043 (2)	0.046 (3)	0.048 (3)	0.011 (2)	0.0139 (19)	0.014 (2)
C12	0.038 (2)	0.065 (3)	0.058 (3)	0.009 (2)	0.014 (2)	0.008 (3)
C13	0.040 (2)	0.056 (3)	0.044 (3)	-0.008 (2)	0.020 (2)	0.000 (2)
C14	0.079 (3)	0.065 (4)	0.062 (3)	0.018 (3)	0.031 (3)	0.032 (3)
C15	0.061 (3)	0.059 (3)	0.054 (3)	0.024 (2)	0.023 (2)	0.026 (2)
C16	0.057 (3)	0.039 (3)	0.046 (3)	0.002 (2)	0.018 (2)	0.004 (2)
C17	0.075 (3)	0.075 (4)	0.068 (4)	0.022 (3)	0.024 (3)	0.034 (3)
C18	0.096 (4)	0.079 (4)	0.065 (4)	0.013 (3)	0.026 (3)	0.036 (3)
C19	0.068 (3)	0.055 (3)	0.050 (3)	-0.011 (3)	0.024 (2)	0.002 (2)
C20	0.058 (3)	0.062 (3)	0.063 (3)	0.002 (2)	0.014 (2)	0.009 (3)
C21	0.057 (3)	0.055 (3)	0.055 (3)	0.004 (2)	0.014 (2)	0.014 (2)
C22	0.059 (3)	0.049 (3)	0.056 (3)	0.006 (2)	0.025 (2)	0.015 (2)
O23	0.092 (3)	0.107 (3)	0.112 (3)	0.055 (2)	0.064 (2)	0.065 (3)
O24	0.0582 (18)	0.075 (2)	0.058 (2)	0.0100 (17)	0.0270 (15)	0.0309 (18)

Geometric parameters (Å, °)

Br1—C13	1.888 (4)	C12—C13	1.362 (6)
Br2—C19	1.895 (4)	C12—H12	0.9300
N1—C8	1.301 (5)	C13—C14	1.377 (6)
N1—N2	1.395 (4)	C14—C15	1.367 (6)
N2—C3	1.313 (5)	C14—H14	0.9300
C3—N4	1.359 (5)	C15—H15	0.9300
C3—C9	1.480 (5)	C16—C21	1.372 (6)
N4—C8	1.355 (4)	C16—C17	1.383 (6)
N4—N5	1.375 (4)	C16—C22	1.491 (6)
N5—C6	1.295 (5)	C17—C18	1.362 (7)
C6—C10	1.460 (5)	C17—H17	0.9300
C6—S7	1.766 (4)	C18—C19	1.364 (7)
S7—C8	1.724 (4)	C18—H18	0.9300
C9—H9A	0.9600	C19—C20	1.360 (7)
C9—H9B	0.9600	C20—C21	1.386 (6)
C9—H9C	0.9600	C20—H20	0.9300
C10—C15	1.380 (6)	C21—H21	0.9300
C10—C11	1.394 (5)	C22—O23	1.205 (5)
C11—C12	1.379 (5)	C22—O24	1.315 (5)
C11—H11	0.9300	O24—H24	0.8200
C8—N1—N2	104.9 (3)	C12—C13—C14	121.1 (4)
C3—N2—N1	110.0 (3)	C12—C13—Br1	119.3 (3)
N2—C3—N4	107.2 (4)	C14—C13—Br1	119.6 (4)
N2—C3—C9	126.9 (4)	C15—C14—C13	119.1 (4)
N4—C3—C9	125.9 (3)	C15—C14—H14	120.4
C8—N4—C3	106.8 (3)	C13—C14—H14	120.4
C8—N4—N5	118.7 (3)	C14—C15—C10	121.5 (4)
C3—N4—N5	134.5 (3)	C14—C15—H15	119.3
C6—N5—N4	107.6 (3)	C10—C15—H15	119.3
N5—C6—C10	122.4 (3)	C21—C16—C17	118.6 (4)
N5—C6—S7	116.7 (3)	C21—C16—C22	121.9 (4)
C10—C6—S7	120.9 (3)	C17—C16—C22	119.5 (4)
C8—S7—C6	87.70 (18)	C18—C17—C16	121.0 (5)
N1—C8—N4	111.2 (3)	C18—C17—H17	119.5
N1—C8—S7	139.6 (3)	C16—C17—H17	119.5
N4—C8—S7	109.2 (3)	C17—C18—C19	119.2 (5)
C3—C9—H9A	109.5	C17—C18—H18	120.4
C3—C9—H9B	109.5	C19—C18—H18	120.4
H9A—C9—H9B	109.5	C20—C19—C18	121.6 (4)
C3—C9—H9C	109.5	C20—C19—Br2	119.4 (4)
H9A—C9—H9C	109.5	C18—C19—Br2	119.0 (4)
H9B—C9—H9C	109.5	C19—C20—C21	118.7 (4)
C15—C10—C11	118.2 (4)	C19—C20—H20	120.6
C15—C10—C6	122.0 (4)	C21—C20—H20	120.6
C11—C10—C6	119.8 (4)	C16—C21—C20	120.7 (5)

C12—C11—C10	120.6 (4)	C16—C21—H21	119.6
C12—C11—H11	119.7	C20—C21—H21	119.6
C10—C11—H11	119.7	O23—C22—O24	123.5 (4)
C13—C12—C11	119.5 (4)	O23—C22—C16	123.2 (5)
C13—C12—H12	120.3	O24—C22—C16	113.2 (4)
C11—C12—H12	120.3	C22—O24—H24	109.5
C8—N1—N2—C3	-0.4 (5)	C15—C10—C11—C12	0.3 (6)
N1—N2—C3—N4	-0.1 (5)	C6—C10—C11—C12	178.6 (4)
N1—N2—C3—C9	179.6 (4)	C10—C11—C12—C13	1.1 (7)
N2—C3—N4—C8	0.5 (5)	C11—C12—C13—C14	-2.0 (7)
C9—C3—N4—C8	-179.2 (4)	C11—C12—C13—Br1	179.6 (3)
N2—C3—N4—N5	-179.2 (4)	C12—C13—C14—C15	1.6 (7)
C9—C3—N4—N5	1.1 (7)	Br1—C13—C14—C15	179.9 (4)
C8—N4—N5—C6	-0.6 (5)	C13—C14—C15—C10	-0.2 (8)
C3—N4—N5—C6	179.1 (4)	C11—C10—C15—C14	-0.7 (7)
N4—N5—C6—C10	180.0 (3)	C6—C10—C15—C14	-179.0 (4)
N4—N5—C6—S7	2.0 (4)	C21—C16—C17—C18	-0.7 (8)
N5—C6—S7—C8	-2.3 (3)	C22—C16—C17—C18	178.9 (5)
C10—C6—S7—C8	179.8 (3)	C16—C17—C18—C19	0.6 (9)
N2—N1—C8—N4	0.8 (5)	C17—C18—C19—C20	-0.4 (8)
N2—N1—C8—S7	-179.3 (4)	C17—C18—C19—Br2	179.5 (4)
C3—N4—C8—N1	-0.8 (5)	C18—C19—C20—C21	0.4 (8)
N5—N4—C8—N1	179.0 (3)	Br2—C19—C20—C21	-179.5 (4)
C3—N4—C8—S7	179.2 (3)	C17—C16—C21—C20	0.6 (7)
N5—N4—C8—S7	-1.0 (4)	C22—C16—C21—C20	-178.9 (4)
C6—S7—C8—N1	-178.3 (5)	C19—C20—C21—C16	-0.5 (7)
C6—S7—C8—N4	1.7 (3)	C21—C16—C22—O23	173.5 (5)
N5—C6—C10—C15	171.8 (4)	C17—C16—C22—O23	-6.0 (8)
S7—C6—C10—C15	-10.4 (6)	C21—C16—C22—O24	-6.7 (6)
N5—C6—C10—C11	-6.5 (6)	C17—C16—C22—O24	173.7 (4)
S7—C6—C10—C11	171.3 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O24—H24 \cdots N2 ⁱ	0.82	1.87	2.674 (4)	169
C9—H9 <i>A</i> \cdots O23 ⁱ	0.96	2.48	3.393 (6)	159

Symmetry code: (i) $-x-1, -y, -z+2$.