

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetramethyl *N,N'*-(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(phosphoramidate)

 Victor A. Trush,^{a*} Kateryna E. Gubina,^a Yaroslav O. Gumeniuk,^b Tetyana Yu. Sliva^a and Irina S. Konovalova^c

^aNational Taras Shevchenko University, Department of Chemistry, Volodymyrska str. 64, 01601 Kyiv, Ukraine, ^bNational University of Life and Environmental Sciences of Ukraine, Heroiv Oboroni str. 15, 03041 Kyiv, Ukraine, and ^cSTC "Institute for Single Crystals", 60 Lenina ave., Khar'kov 61001, Ukraine
Correspondence e-mail: trush@univ.kiev.ua


Received 20 February 2012; accepted 14 March 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.106; data-to-parameter ratio = 20.0.

The molecule of the title compound, $\text{C}_9\text{H}_{14}\text{F}_6\text{N}_2\text{O}_8\text{P}_2$, lies on a twofold rotation axis that passes through the middle C atom of the three-atom fluoromethylene unit. The carbonyl and phosphoryl groups are in an antiperiplanar conformation. In the crystal, $\text{N}-\text{H}\cdots\text{O}=\text{P}$ hydrogen bonds link the molecules into polymeric chains parallel to the c axis.

Related literature

For background to the chemistry of phosphorus–organic compounds, see: Ly & Woollins (1998). For the biological and pharmacological properties of carbacylamidophosphate derivatives, see: Adams *et al.* (2002). For details of the synthesis and properties of phosphoramidate derivatives, see: Kirsanov & Levchenko (1957); For structural analogues of phosphorylated carbacylamides and their properties, see: Trush *et al.* (2005); Gubina *et al.* (2000). For the synthesis and properties of fluorinated compounds, see: Leontieva *et al.* (2002).



Experimental

Crystal data

 $\text{C}_9\text{H}_{14}\text{F}_6\text{N}_2\text{O}_8\text{P}_2$
 $M_r = 454.16$

Monoclinic, $C2/c$
 $a = 19.7862$ (13) Å
 $b = 5.2801$ (4) Å
 $c = 16.9943$ (11) Å
 $\beta = 100.427$ (6)°
 $V = 1746.1$ (2) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.35$ mm⁻¹
 $T = 293$ K
 $0.40 \times 0.20 \times 0.10$ mm

Data collection

Oxford Diffraction Xcalibur3 diffractometer
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)
 $T_{\min} = 0.872$, $T_{\max} = 0.966$
 7084 measured reflections

2504 independent reflections
 1669 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 2 standard reflections every 50 reflections
 intensity decay: 0.3%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.106$
 $S = 0.93$
 2504 reflections

125 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.29$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N1A}\cdots\text{O4}^i$	0.86	1.93	2.7750 (17)	168

 Symmetry code: (i) $-x + 1, -y, -z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2006); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2523).

References

- Adams, L. A., Cox, R. J., Gibson, J. S., Mayo-Martin, M. B., Walter, M. & Whittingham, W. (2002). *Chem. Commun.* pp. 2004–2005.
 Gubina, K. E., Ovchinnikov, V. A., Amirkhanov, V. M., Skopenko, V. V. & Shishkin, O. V. (2000). *Z. Naturforsch. Teil B*, **55**, 495–498.
 Kirsanov, A. & Levchenko, E. (1957). *Zh. Obshch. Khim.* **27**, 2585–2590.
 Leontieva, O. S., Chapurkin, V. V. & Baklanov, A. V. (2002). *Russ. J. Org. Chem.* **38**, 1375–1376.
 Ly, T. Q. & Woollins, J. D. (1998). *Coord. Chem. Rev.* **176**, 451–481.
 Oxford Diffraction (2006). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction, Abingdon, England.
 Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Trush, V. A., Gubina, K. E., Amirkhanov, V. M., Swiatek-Kozłowska, J. & Domasevitch, K. V. (2005). *Polyhedron*, **24**, 1007–1014.

supporting information

Acta Cryst. (2012). E68, o1127 [https://doi.org/10.1107/S1600536812011191]

Tetramethyl *N,N'*-(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis-(phosphoramidate)

Victor A. Trush, Kateryna E. Gubina, Yaroslav O. Gumeniuk, Tetyana Yu. Sliva and Irina S. Konovalova

S1. Comment

Tetramethyl (2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate) is a representative of the carbacylamidophosphates (CAPH), a family of compounds containing C(O)NHP(O) group. The presence of a peptide group in the carbacylamidophosphates causes its diverse biological activity. (Adams *et al.*, 2002). CAPH may be regarded as powerful chelating ligand systems. There has recently been a resurgence of interest in their coordination chemistry as a consequence of the steric control that this ligand system may impart compared to, for example, β -diketonates. The wide range of coordination compounds were synthesized and described in details (Ly & Woollins, 1998).

The crystal structure of tetramethyl (2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate) (**1**) reveals, that the molecule of phosphorylated amide consists of symmetric moiety lying on a twofold rotation axis that passes through the middle atom of the three-atom fluoromethylene unit ($-x, y, -z + 1/2$). CF_2 groups are situated in the retarded conformation to each other with the values of dihedral FCCF angles in the range $57.76 - 68.62^\circ$ (Fig.2). In the molecule, the carbonyl and phosphoryl groups are in antiperiplanar conformation. The frame O4—P1—N1—C3—O1 is almost flat (the values of dihedral O4—P1—N1—C3 and O1—C3—N1—P1 angles are -169.54 and -0.05° , respectively) as it has been observed for the most CAPH (Gubina *et al.*, 2000; Trush *et al.* 2005). The crystal is composed from polymer chains which are built from molecules linked *via* intermolecular hydrogen N—H \cdots O=P bonds (Fig.3). The parameters of the intermolecular hydrogen bond are listed in Table 1.

S2. Experimental

The compound tetramethyl(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate) (**1**) can be synthesized by multistep reaction (Fig.1) starting from dimethyl hexafluoropentanedioate (Leontieva *et al.* 2002). The solution of 26.81 g (0.1 mol) dimethyl hexafluoropentanedioate in 40 ml of methanol was added drop-wise to the well stirred saturated solution of ammonia in methanol (~ 200 ml) under cooling. The obtained mixture was allowed to stand for a weak. Then solvent was removed under reduced pressure to give the crude product. Recrystallization from water gave the white powder of 2,2,3,3,4,4-hexafluoropentanediamide (22.6 g, 95%). Subsequently, the meticulous dried diamide (11.91 g, 0.05 mol) was involved in phosphoroazo-reaction (Kirsanov & Levchenko, 1957) with 20.82 g (0.1 mol) of PCl_5 in 10 ml CCl_4 . The treatment of crude hexachloroanhydride with NaOCH_3 (0.3 mol) in methanol solution leads to obtain hexaester with good yield. Further alkaline hydrolysis and acidification gave the final product - tetramethyl(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate) (yield 13.6 g, 60%). The single crystals of **1** suitable for X-ray analysis were grown from aqueous-methanol solution (1:1).

S3. Refinement

All H atoms were placed at calculated positions and treated as riding on their parent atoms [$C-H = 0.96 \text{ \AA}$, and $U_{iso}(H) = 1.5U_{eq}(C)$, $N-H = 0.86 \text{ \AA}$ and $U_{iso}(H) = 1.2U_{eq}(N)$].




Figure 1

Scheme of synthesis of tetramethyl (2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate).




Figure 2

A view of tetramethyl(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane-1,5-diyl)bis(amidophosphate). Symmetry transformation used to generate equivalent atoms: $i - x + 1, y, -z + 1/2$.




Figure 3

A system of hydrogen bonds as a one-dimensional polymeric ladder chains.

Tetramethyl *N,N'*-(2,2,3,3,4,4-hexafluoro-1,5-dioxopentane- 1,5-diyl)bis(phosphoramidate)*Crystal data*

C₉H₁₄F₆N₂O₈P₂
M_r = 454.16
 Monoclinic, *C*2/*c*
 Hall symbol: -*C* 2yc
a = 19.7862 (13) Å
b = 5.2801 (4) Å
c = 16.9943 (11) Å
 β = 100.427 (6)°
V = 1746.1 (2) Å³
Z = 4

F(000) = 920
D_x = 1.728 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 2504 reflections
 θ = 2.9–30°
 μ = 0.35 mm⁻¹
T = 293 K
 Block, colourless
 0.40 × 0.20 × 0.10 mm

Data collection

Oxford Diffraction Xcalibur3
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Oxford Diffraction, 2010)
T_{min} = 0.872, *T_{max}* = 0.966
 7084 measured reflections

2504 independent reflections
 1669 reflections with *I* > 2σ(*I*)
R_{int} = 0.023
 θ_{\max} = 30.0°, θ_{\min} = 2.9°
 $h = -27 \rightarrow 27$
 $k = -7 \rightarrow 7$
 $l = -23 \rightarrow 22$
 2 standard reflections every 50 reflections
 intensity decay: 0.3%

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.037
wR(*F*²) = 0.106
S = 0.93
 2504 reflections
 125 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0667P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.29 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$

Special details

Experimental. Analysis found: IR (KBr pellet, cm⁻¹): 3095(s, NH), 2925(ns, CH), 1190(s), 1746(s, C=O), 1478(s CN); 1291 (as, CF), 1212 (s, PO), 1141 (s, CF). NMR -¹H (DMSO-d₆): C—H 3.74 (*d*) 12H, ³J_{PH} = 11.6 Hz; NH 11.38 (*d*) 2H; ³¹P (DMSO-d₆): -0.28 (hept) ³J_{HP} = 11.6 Hz; ¹³C (DMSO-d₆): C(O) 159.68, CF 108.6 - 105.9, CH₃ 54.74.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of *F*² against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*², conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*². The threshold expression of *F*² > 2σ(*F*²) is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*² are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
F1	0.54045 (6)	-0.1156 (2)	0.30276 (6)	0.0559 (3)

F2	0.58399 (6)	-0.5823 (2)	0.25478 (7)	0.0568 (3)
F3	0.50776 (6)	-0.5503 (2)	0.14516 (6)	0.0495 (3)
P1	0.61282 (2)	0.06932 (8)	0.04812 (2)	0.03549 (13)
N1	0.57021 (7)	-0.1298 (3)	0.09836 (8)	0.0368 (3)
H1NA	0.5280	-0.1619	0.0784	0.044*
O1	0.65572 (7)	-0.2202 (3)	0.20427 (8)	0.0509 (3)
O2	0.67108 (6)	-0.0844 (2)	0.01929 (7)	0.0470 (3)
O3	0.65216 (6)	0.2593 (2)	0.10883 (7)	0.0458 (3)
O4	0.56309 (6)	0.1876 (3)	-0.01514 (7)	0.0463 (3)
C1	0.5000	-0.2656 (4)	0.2500	0.0349 (5)
C2	0.54710 (9)	-0.4170 (3)	0.20389 (9)	0.0368 (3)
C3	0.59784 (8)	-0.2443 (3)	0.16861 (9)	0.0359 (3)
C4	0.65304 (12)	-0.2569 (4)	-0.04796 (13)	0.0609 (6)
H4C	0.6191	-0.3749	-0.0369	0.091*
H4B	0.6933	-0.3475	-0.0561	0.091*
H4A	0.6349	-0.1621	-0.0953	0.091*
C5	0.72440 (10)	0.2612 (5)	0.14340 (16)	0.0672 (6)
H5C	0.7340	0.4008	0.1799	0.101*
H5B	0.7509	0.2787	0.1017	0.101*
H5A	0.7363	0.1054	0.1716	0.101*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0650 (7)	0.0573 (7)	0.0483 (6)	-0.0249 (6)	0.0182 (6)	-0.0188 (5)
F2	0.0620 (7)	0.0545 (7)	0.0567 (7)	0.0208 (6)	0.0179 (6)	0.0233 (5)
F3	0.0586 (7)	0.0482 (6)	0.0447 (6)	-0.0125 (5)	0.0172 (5)	-0.0133 (5)
P1	0.0276 (2)	0.0451 (2)	0.0324 (2)	-0.00321 (17)	0.00186 (15)	0.00114 (16)
N1	0.0276 (7)	0.0485 (8)	0.0325 (6)	-0.0032 (6)	0.0008 (5)	0.0049 (6)
O1	0.0375 (7)	0.0623 (8)	0.0471 (7)	-0.0022 (6)	-0.0077 (6)	0.0082 (6)
O2	0.0322 (6)	0.0634 (8)	0.0463 (7)	-0.0026 (6)	0.0093 (5)	-0.0114 (6)
O3	0.0347 (6)	0.0518 (7)	0.0479 (7)	-0.0046 (5)	-0.0009 (5)	-0.0094 (5)
O4	0.0367 (6)	0.0573 (7)	0.0413 (7)	-0.0064 (6)	-0.0026 (5)	0.0145 (5)
C1	0.0389 (12)	0.0357 (11)	0.0296 (10)	0.000	0.0045 (9)	0.000
C2	0.0420 (9)	0.0365 (8)	0.0311 (7)	0.0032 (7)	0.0042 (6)	0.0042 (6)
C3	0.0337 (8)	0.0413 (8)	0.0317 (7)	0.0025 (7)	0.0032 (6)	0.0009 (6)
C4	0.0603 (13)	0.0614 (13)	0.0622 (13)	-0.0049 (11)	0.0143 (11)	-0.0208 (10)
C5	0.0381 (10)	0.0707 (14)	0.0843 (15)	-0.0082 (10)	-0.0121 (10)	-0.0189 (12)

Geometric parameters (Å, °)

F1—C1	1.3461 (16)	O3—C5	1.444 (2)
F2—C2	1.3469 (18)	C1—F1 ⁱ	1.3461 (16)
F3—C2	1.3475 (18)	C1—C2 ⁱ	1.545 (2)
P1—O4	1.4598 (12)	C1—C2	1.545 (2)
P1—O3	1.5444 (12)	C2—C3	1.555 (2)
P1—O2	1.5595 (13)	C4—H4C	0.9600
P1—N1	1.6751 (14)	C4—H4B	0.9600

N1—C3	1.361 (2)	C4—H4A	0.9600
N1—H1NA	0.8600	C5—H5C	0.9600
O1—C3	1.202 (2)	C5—H5B	0.9600
O2—C4	1.454 (2)	C5—H5A	0.9600
O4—P1—O3	113.89 (8)	F3—C2—C1	108.98 (12)
O4—P1—O2	115.43 (8)	F2—C2—C3	108.38 (13)
O3—P1—O2	103.65 (7)	F3—C2—C3	110.33 (12)
O4—P1—N1	108.04 (7)	C1—C2—C3	112.52 (14)
O3—P1—N1	107.84 (7)	O1—C3—N1	126.17 (16)
O2—P1—N1	107.59 (7)	O1—C3—C2	119.38 (14)
C3—N1—P1	124.51 (11)	N1—C3—C2	114.45 (13)
C3—N1—H1NA	117.7	O2—C4—H4C	109.5
P1—N1—H1NA	117.7	O2—C4—H4B	109.5
C4—O2—P1	118.85 (13)	H4C—C4—H4B	109.5
C5—O3—P1	128.21 (14)	O2—C4—H4A	109.5
F1—C1—F1 ⁱ	107.92 (19)	H4C—C4—H4A	109.5
F1—C1—C2 ⁱ	107.89 (7)	H4B—C4—H4A	109.5
F1 ⁱ —C1—C2 ⁱ	107.56 (8)	O3—C5—H5C	109.5
F1—C1—C2	107.56 (8)	O3—C5—H5B	109.5
F1 ⁱ —C1—C2	107.89 (7)	H5C—C5—H5B	109.5
C2 ⁱ —C1—C2	117.67 (19)	O3—C5—H5A	109.5
F2—C2—F3	108.08 (13)	H5C—C5—H5A	109.5
F2—C2—C1	108.44 (12)	H5B—C5—H5A	109.5
O4—P1—N1—C3	-169.53 (14)	F1 ⁱ —C1—C2—F3	-57.75 (17)
O3—P1—N1—C3	-46.00 (15)	C2 ⁱ —C1—C2—F3	64.07 (10)
O2—P1—N1—C3	65.23 (15)	F1—C1—C2—C3	-51.23 (15)
O4—P1—O2—C4	-45.46 (17)	F1 ⁱ —C1—C2—C3	64.95 (15)
O3—P1—O2—C4	-170.70 (14)	C2 ⁱ —C1—C2—C3	-173.22 (13)
N1—P1—O2—C4	75.24 (15)	P1—N1—C3—O1	-0.1 (3)
O4—P1—O3—C5	-137.75 (19)	P1—N1—C3—C2	178.97 (11)
O2—P1—O3—C5	-11.5 (2)	F2—C2—C3—O1	-22.1 (2)
N1—P1—O3—C5	102.4 (2)	F3—C2—C3—O1	-140.28 (16)
F1—C1—C2—F2	68.63 (17)	C1—C2—C3—O1	97.78 (17)
F1 ⁱ —C1—C2—F2	-175.18 (13)	F2—C2—C3—N1	158.77 (14)
C2 ⁱ —C1—C2—F2	-53.36 (10)	F3—C2—C3—N1	40.62 (19)
F1—C1—C2—F3	-173.94 (12)	C1—C2—C3—N1	-81.33 (15)

Symmetry code: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1NA \cdots O4 ⁱⁱ	0.86	1.93	2.7750 (17)	168

Symmetry code: (ii) $-x+1, -y, -z$.