

1-(4-Methylphenyl)-3-phenyl-1*H*-pyrazol-5-yl 4-nitrobenzenesulfonate

Solange M. S. V. Wardell,^a Edward R. T. Tiekink^{b*} and James L. Wardell^{c‡}

^aCHEMSOL, 1 Harcourt Road, Aberdeen, AB15 5NY, Scotland, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cCentro de Desenvolvimento Tecnológico em Saúde (CDTS), Fundação Oswaldo Cruz (FIOCRUZ), Casa Amarela, Campus de Manguinhos, Av. Brasil 4365, 21040-900 Rio de Janeiro, RJ, Brazil
Correspondence e-mail: edward.tiekink@gmail.com


Received 8 March 2012; accepted 9 March 2012

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.069; wR factor = 0.179; data-to-parameter ratio = 15.8.

In the title molecule, $\text{C}_{22}\text{H}_{17}\text{N}_3\text{O}_5\text{S}$, the pyrazole ring is planar (r.m.s. deviation = 0.018 Å) and forms dihedral angles of 21.45 (10) and 6.96 (10)° with the N- and C-bound benzene rings, respectively. Supramolecular layers in the *bc* plane are formed in the crystal *via* C–H···O and π – π interactions involving the sulfonamide benzene ring interacting with the N- and C-bound benzene rings [centroid–centroid distances = 3.790 (2) and 3.730 (2) Å, respectively]. The crystal studied was found to be a merohedral twin (twin law 1 0 0.678, 0 $\bar{1}$ 0, 0 0 $\bar{1}$), the fractional contribution of the minor component being approximately 36%.

Related literature

For related structures and background references to pyrazoles, see: Wardell *et al.* (2012); Baddeley *et al.* (2012). For the synthesis, see: Galoyan *et al.* (1969). For the treatment of twinned diffraction data, see: Spek (2009).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{17}\text{N}_3\text{O}_5\text{S}$	$V = 1964.3\text{ (3) \AA}^3$
$M_r = 435.46$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 13.5339\text{ (12) \AA}$	$\mu = 0.21\text{ mm}^{-1}$
$b = 10.4827\text{ (10) \AA}$	$T = 120\text{ K}$
$c = 14.9303\text{ (13) \AA}$	$0.58 \times 0.38 \times 0.04\text{ mm}$
$\beta = 111.975\text{ (3)}^\circ$	

Data collection

Rigaku Saturn724+ diffractometer	4454 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2007)	4454 independent reflections
$T_{\min} = 0.620$, $T_{\max} = 0.746$	3951 reflections with $I > 2\sigma(I)$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$	282 parameters
$wR(F^2) = 0.179$	H-atom parameters constrained
$S = 1.19$	$\Delta\rho_{\max} = 0.59\text{ e \AA}^{-3}$
4454 reflections	$\Delta\rho_{\min} = -0.62\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C5}-\text{H5}\cdots \text{O4}^{\dagger}$	0.95	2.50	3.387 (5)	155
Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$				

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The use of the EPSRC X-ray crystallographic service at the University of Southampton, England, and the valuable assistance of the staff there is gratefully acknowledged. JLW acknowledges support from CAPES (Brazil). Support from the Ministry of Higher Education, Malaysia, High-Impact Research scheme (UM.C/HIR/MOHE/SC/12) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6674).

References

- Baddeley, T. C., Wardell, S. M. S. V., Tiekink, E. R. T. & Wardell, J. L. (2012). *Acta Cryst. E68*, o1016–o1017.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Galoyan, G. A., Agbalyan, S. G. & Esayan, G. T. (1969). *Arm. Khim. Zhur.* **22**, 430–433.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.

‡ Additional correspondence author, e-mail: j.wardell@abdn.ac.uk.

- Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wardell, S. M. S. V., Howie, A. H., Tiekink, E. R. T. & Wardell, J. L. (2012).
Acta Cryst. E **68**, o992–o993.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, o1086–o1087 [https://doi.org/10.1107/S1600536812010598]

1-(4-Methylphenyl)-3-phenyl-1*H*-pyrazol-5-yl 4-nitrobenzenesulfonate

Solange M. S. V. Wardell, Edward R. T. Tiekink and James L. Wardell

S1. Comment

The structure of the title compound is now reported in continuation of related structural studies (Wardell *et al.* 2012; Baddeley *et al.*, 2012).

In the title compound, Fig. 2, the pyrazole ring is planar with a r.m.s. deviation for the fitted atoms of 0.018 Å; the maximum deviations from this plane are 0.015 (1) Å (for the N1 atom) and -0.015 (1) Å (C8). The N– and C-bound benzene rings are inclined to this plane forming dihedral angles of 21.45 (10) and 6.96 (10)°, respectively; the dihedral angle between the benzene rings is 20.42 (10)° consistent with a non-planar molecule.


In the crystal, molecules are assembled into supramolecular layers in the *bc* plane *via* C—H···O, Table 1, and π — π interactions involving the sulfonamide benzene ring interacting with the N– and C-bound benzene rings {ring centroid···ring centroid distances = 3.790 (2) Å [angle of inclination = 0.96 (17)° for symmetry operation 1 - *x*, 1 - *y*, -*z*] and 3.730 (2) Å [angle of inclination = 10.02 (17)° for symmetry operation 1 - *x*, -1/2 + *y*, -1/2 - *z*], respectively}, Fig. 3. Layers stack along the *a* axis with no specific interactions between them, Fig. 4.

S2. Experimental


A solution of 4-MeC₆H₄NHNH₂.HCl (2 mmol) and PhCOCH₂CONHPh (2 mmol) in Me₂CO (20 ml) was refluxed for 1 h. A solution of 4-nitrobenzenesulfonyl chloride (2 mmol) in Me₂CO (10 ml) was added and the reaction mixture was refluxed for 30 min, rotary evaporated and the residue was recrystallized twice from EtOH as yellow plates, *M.pt*: 445–447 K.

S3. Refinement


The C-bound H atoms were geometrically placed (C—H = 0.95–0.98 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2$ –1.5 $U_{\text{eq}}(\text{C})$. Owing to poor agreement three reflections, *i.e.* (−7 0 10), (−7 1 13) and (−12 5 13), were removed from the final cycles of refinement. The sample was a non-merohedral twin (twin law 1 0 0.678, 0 −1 0, 0 0 −1) and the fractional contribution of the minor component refined to 0.362 (2). The twin domains were separated by using the *TwinRotMat* routine in *PLATON* (Spek, 2009).

**Figure 1**


Reaction scheme. For further details see Baddeley *et al.* (2012).

**Figure 2**

The molecular structure of (I) showing displacement ellipsoids at the 50% probability level.

**Figure 3**

Supramolecular layer in the *bc* plane in (I) sustained by C—H···O and π—π interactions shown as orange and purple dashed lines, respectively.

**Figure 4**

A view in projection down the c axis of the crystal packing in (I) showing the stacking of layers. The $\text{C}—\text{H}··\cdot\text{O}$, and $\pi—\pi$ interactions are shown as orange and purple dashed lines, respectively.

1-(4-Methylphenyl)-3-phenyl-1*H*-pyrazol-5-yl 4-nitrobenzenesulfonate

Crystal data


 $M_r = 435.46$

 Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 13.5339 (12)$ Å

 $b = 10.4827 (10)$ Å

 $c = 14.9303 (13)$ Å

 $\beta = 111.975 (3)^\circ$
 $V = 1964.3 (3)$ Å³
 $Z = 4$
 $F(000) = 904$
 $D_x = 1.472 \text{ Mg m}^{-3}$

 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 67707 reflections

 $\theta = 2.9—27.5^\circ$
 $\mu = 0.21 \text{ mm}^{-1}$
 $T = 120 \text{ K}$

Plate, yellow

 $0.58 \times 0.38 \times 0.04$ mm

Data collection

 Rigaku Saturn724+
diffractometer

Radiation source: Rotating Anode

Confocal monochromator

 Detector resolution: 28.5714 pixels mm⁻¹

 profile data from ω -scans

 Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2007)

 $T_{\min} = 0.620$, $T_{\max} = 0.746$

4454 measured reflections

4454 independent reflections

 3951 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.000$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.2^\circ$
 $h = -17 \rightarrow 16$
 $k = 0 \rightarrow 13$
 $l = 0 \rightarrow 19$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.069$$

$$wR(F^2) = 0.179$$

$$S = 1.19$$

4454 reflections

282 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0562P)^2 + 4.737P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.59 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.62 \text{ e \AA}^{-3}$$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.47829 (6)	0.20331 (9)	0.00726 (6)	0.0203 (2)
O1	0.51909 (18)	0.3508 (3)	0.01445 (17)	0.0203 (5)
O2	0.5619 (2)	0.1219 (3)	0.0044 (2)	0.0268 (6)
O3	0.4369 (2)	0.1938 (3)	0.08223 (19)	0.0315 (7)
O4	0.1490 (2)	0.2303 (3)	-0.4513 (2)	0.0360 (7)
O5	0.0359 (2)	0.2650 (3)	-0.3824 (2)	0.0377 (8)
N1	0.7023 (2)	0.4081 (3)	0.0618 (2)	0.0156 (6)
N2	0.7780 (2)	0.4243 (3)	0.0225 (2)	0.0156 (6)
N3	0.1265 (2)	0.2427 (3)	-0.3796 (2)	0.0254 (7)
C1	0.6074 (2)	0.3760 (3)	-0.0095 (2)	0.0169 (7)
C2	0.7290 (2)	0.4006 (3)	-0.0717 (2)	0.0141 (6)
C3	0.6198 (3)	0.3718 (3)	-0.0957 (2)	0.0183 (7)
H3	0.5673	0.3539	-0.1576	0.022*
C4	0.3726 (3)	0.2062 (3)	-0.1065 (3)	0.0185 (7)
C5	0.2715 (3)	0.2430 (4)	-0.1110 (3)	0.0206 (7)
H5	0.2587	0.2609	-0.0539	0.025*
C6	0.1898 (3)	0.2525 (4)	-0.2022 (3)	0.0222 (7)
H6	0.1196	0.2749	-0.2081	0.027*
C7	0.2126 (3)	0.2292 (3)	-0.2831 (3)	0.0199 (7)
C8	0.3132 (3)	0.1929 (4)	-0.2796 (3)	0.0220 (7)
H8	0.3262	0.1777	-0.3370	0.026*
C9	0.3942 (3)	0.1796 (3)	-0.1886 (3)	0.0189 (7)
H9	0.4633	0.1527	-0.1829	0.023*
C10	0.7302 (3)	0.4228 (3)	0.1639 (2)	0.0152 (6)
C11	0.8359 (3)	0.3992 (3)	0.2245 (3)	0.0181 (7)

H11	0.8871	0.3757	0.1980	0.022*
C12	0.8651 (3)	0.4106 (3)	0.3239 (2)	0.0198 (7)
H12	0.9369	0.3951	0.3651	0.024*
C13	0.7909 (3)	0.4445 (3)	0.3644 (3)	0.0210 (7)
C14	0.6856 (3)	0.4686 (4)	0.3023 (3)	0.0211 (7)
H14	0.6343	0.4919	0.3287	0.025*
C15	0.6551 (3)	0.4590 (3)	0.2020 (3)	0.0197 (7)
H15	0.5838	0.4770	0.1604	0.024*
C16	0.8218 (3)	0.4510 (4)	0.4733 (3)	0.0307 (9)
H16A	0.7642	0.4912	0.4879	0.046*
H16B	0.8871	0.5014	0.5020	0.046*
H16C	0.8341	0.3645	0.5002	0.046*
C17	0.7883 (3)	0.4054 (3)	-0.1369 (2)	0.0157 (6)
C18	0.7359 (3)	0.3973 (3)	-0.2370 (2)	0.0166 (6)
H18	0.6609	0.3851	-0.2640	0.020*
C19	0.7925 (3)	0.4068 (3)	-0.2978 (2)	0.0191 (7)
H19	0.7563	0.4007	-0.3658	0.023*
C20	0.9031 (3)	0.4255 (4)	-0.2584 (3)	0.0210 (7)
H20	0.9417	0.4348	-0.2997	0.025*
C21	0.9563 (3)	0.4303 (3)	-0.1586 (3)	0.0194 (7)
H21	1.0315	0.4409	-0.1319	0.023*
C22	0.8997 (3)	0.4196 (3)	-0.0977 (2)	0.0185 (7)
H22	0.9365	0.4218	-0.0297	0.022*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0124 (4)	0.0288 (5)	0.0195 (4)	-0.0052 (3)	0.0057 (3)	0.0020 (4)
O1	0.0110 (10)	0.0319 (14)	0.0212 (12)	-0.0056 (10)	0.0097 (10)	-0.0077 (10)
O2	0.0167 (12)	0.0245 (14)	0.0359 (15)	0.0004 (10)	0.0058 (11)	0.0053 (12)
O3	0.0206 (12)	0.0545 (19)	0.0206 (13)	-0.0108 (13)	0.0090 (11)	0.0067 (13)
O4	0.0335 (15)	0.0501 (19)	0.0210 (14)	-0.0070 (14)	0.0061 (12)	-0.0007 (13)
O5	0.0202 (13)	0.050 (2)	0.0338 (16)	0.0089 (13)	-0.0002 (12)	-0.0092 (14)
N1	0.0106 (12)	0.0238 (15)	0.0136 (13)	-0.0030 (11)	0.0057 (11)	0.0008 (11)
N2	0.0110 (12)	0.0238 (15)	0.0144 (13)	-0.0021 (11)	0.0075 (11)	0.0014 (11)
N3	0.0230 (15)	0.0232 (16)	0.0244 (17)	-0.0028 (13)	0.0024 (13)	-0.0025 (13)
C1	0.0077 (13)	0.0240 (17)	0.0179 (15)	-0.0046 (12)	0.0034 (12)	-0.0022 (14)
C2	0.0113 (14)	0.0175 (16)	0.0127 (15)	-0.0014 (12)	0.0034 (12)	-0.0004 (12)
C3	0.0126 (14)	0.0237 (18)	0.0171 (16)	-0.0034 (13)	0.0037 (12)	-0.0022 (13)
C4	0.0124 (14)	0.0224 (17)	0.0190 (16)	-0.0046 (13)	0.0040 (12)	-0.0001 (14)
C5	0.0161 (15)	0.0254 (18)	0.0228 (18)	-0.0043 (14)	0.0100 (13)	-0.0032 (14)
C6	0.0139 (15)	0.0242 (18)	0.0274 (19)	-0.0010 (14)	0.0066 (14)	-0.0014 (15)
C7	0.0146 (15)	0.0196 (17)	0.0225 (17)	-0.0042 (13)	0.0035 (13)	-0.0020 (14)
C8	0.0190 (16)	0.0241 (18)	0.0227 (17)	-0.0044 (14)	0.0077 (14)	-0.0029 (14)
C9	0.0137 (14)	0.0210 (17)	0.0240 (18)	-0.0029 (13)	0.0093 (13)	-0.0036 (14)
C10	0.0163 (15)	0.0177 (16)	0.0104 (15)	-0.0014 (12)	0.0036 (12)	-0.0001 (12)
C11	0.0141 (15)	0.0231 (17)	0.0172 (16)	0.0001 (13)	0.0060 (13)	0.0017 (13)
C12	0.0174 (16)	0.0220 (18)	0.0172 (17)	-0.0024 (13)	0.0034 (13)	0.0007 (13)

C13	0.0273 (18)	0.0200 (17)	0.0164 (16)	-0.0054 (14)	0.0090 (14)	-0.0015 (14)
C14	0.0214 (17)	0.0235 (18)	0.0225 (18)	-0.0019 (14)	0.0130 (15)	-0.0020 (14)
C15	0.0170 (15)	0.0209 (16)	0.0218 (17)	-0.0019 (13)	0.0080 (14)	-0.0015 (14)
C16	0.037 (2)	0.039 (2)	0.0143 (17)	-0.0007 (18)	0.0082 (16)	0.0005 (16)
C17	0.0159 (15)	0.0178 (16)	0.0145 (15)	-0.0020 (12)	0.0071 (13)	-0.0001 (12)
C18	0.0159 (15)	0.0189 (16)	0.0137 (15)	-0.0007 (12)	0.0040 (12)	-0.0002 (13)
C19	0.0209 (16)	0.0214 (17)	0.0135 (16)	-0.0014 (13)	0.0050 (13)	-0.0006 (13)
C20	0.0217 (17)	0.0263 (19)	0.0220 (18)	-0.0007 (14)	0.0163 (15)	-0.0008 (14)
C21	0.0134 (15)	0.0246 (18)	0.0207 (17)	-0.0030 (13)	0.0071 (13)	-0.0002 (14)
C22	0.0161 (15)	0.0248 (18)	0.0135 (15)	-0.0007 (13)	0.0045 (13)	-0.0001 (13)

Geometric parameters (\AA , $^{\circ}$)

S1—O3	1.430 (3)	C10—C15	1.391 (5)
S1—O2	1.431 (3)	C10—C11	1.399 (4)
S1—O1	1.632 (3)	C11—C12	1.391 (5)
S1—C4	1.765 (3)	C11—H11	0.9500
O1—C1	1.395 (4)	C12—C13	1.397 (5)
O4—N3	1.225 (4)	C12—H12	0.9500
O5—N3	1.233 (4)	C13—C14	1.402 (5)
N1—N2	1.366 (4)	C13—C16	1.522 (5)
N1—C1	1.368 (4)	C14—C15	1.401 (5)
N1—C10	1.435 (4)	C14—H14	0.9500
N2—C2	1.336 (4)	C15—H15	0.9500
N3—C7	1.482 (5)	C16—H16A	0.9800
C1—C3	1.360 (5)	C16—H16B	0.9800
C2—C3	1.417 (4)	C16—H16C	0.9800
C2—C17	1.476 (4)	C17—C18	1.397 (5)
C3—H3	0.9500	C17—C22	1.406 (5)
C4—C9	1.390 (5)	C18—C19	1.393 (5)
C4—C5	1.399 (5)	C18—H18	0.9500
C5—C6	1.399 (5)	C19—C20	1.402 (5)
C5—H5	0.9500	C19—H19	0.9500
C6—C7	1.376 (5)	C20—C21	1.391 (5)
C6—H6	0.9500	C20—H20	0.9500
C7—C8	1.396 (5)	C21—C22	1.396 (5)
C8—C9	1.397 (5)	C21—H21	0.9500
C8—H8	0.9500	C22—H22	0.9500
C9—H9	0.9500		
O3—S1—O2	121.83 (18)	C15—C10—N1	121.5 (3)
O3—S1—O1	103.66 (16)	C11—C10—N1	117.8 (3)
O2—S1—O1	108.38 (14)	C12—C11—C10	119.2 (3)
O3—S1—C4	109.94 (16)	C12—C11—H11	120.4
O2—S1—C4	110.29 (17)	C10—C11—H11	120.4
O1—S1—C4	100.35 (15)	C11—C12—C13	121.4 (3)
C1—O1—S1	117.6 (2)	C11—C12—H12	119.3
N2—N1—C1	109.5 (3)	C13—C12—H12	119.3

N2—N1—C10	120.0 (3)	C12—C13—C14	118.5 (3)
C1—N1—C10	130.5 (3)	C12—C13—C16	121.3 (3)
C2—N2—N1	105.8 (2)	C14—C13—C16	120.2 (3)
O4—N3—O5	124.0 (3)	C15—C14—C13	120.9 (3)
O4—N3—C7	118.5 (3)	C15—C14—H14	119.5
O5—N3—C7	117.4 (3)	C13—C14—H14	119.5
C3—C1—N1	109.4 (3)	C10—C15—C14	119.2 (3)
C3—C1—O1	131.1 (3)	C10—C15—H15	120.4
N1—C1—O1	119.5 (3)	C14—C15—H15	120.4
N2—C2—C3	111.4 (3)	C13—C16—H16A	109.5
N2—C2—C17	120.7 (3)	C13—C16—H16B	109.5
C3—C2—C17	127.9 (3)	H16A—C16—H16B	109.5
C1—C3—C2	103.9 (3)	C13—C16—H16C	109.5
C1—C3—H3	128.0	H16A—C16—H16C	109.5
C2—C3—H3	128.0	H16B—C16—H16C	109.5
C9—C4—C5	122.5 (3)	C18—C17—C22	119.2 (3)
C9—C4—S1	118.9 (3)	C18—C17—C2	121.3 (3)
C5—C4—S1	118.5 (3)	C22—C17—C2	119.5 (3)
C4—C5—C6	118.0 (3)	C19—C18—C17	120.7 (3)
C4—C5—H5	121.0	C19—C18—H18	119.7
C6—C5—H5	121.0	C17—C18—H18	119.7
C7—C6—C5	119.1 (3)	C18—C19—C20	119.8 (3)
C7—C6—H6	120.4	C18—C19—H19	120.1
C5—C6—H6	120.4	C20—C19—H19	120.1
C6—C7—C8	123.5 (3)	C21—C20—C19	119.9 (3)
C6—C7—N3	118.9 (3)	C21—C20—H20	120.1
C8—C7—N3	117.6 (3)	C19—C20—H20	120.1
C7—C8—C9	117.5 (3)	C20—C21—C22	120.3 (3)
C7—C8—H8	121.2	C20—C21—H21	119.9
C9—C8—H8	121.2	C22—C21—H21	119.9
C4—C9—C8	119.4 (3)	C21—C22—C17	120.1 (3)
C4—C9—H9	120.3	C21—C22—H22	119.9
C8—C9—H9	120.3	C17—C22—H22	119.9
C15—C10—C11	120.7 (3)		
O3—S1—O1—C1	147.7 (2)	C6—C7—C8—C9	0.2 (6)
O2—S1—O1—C1	17.0 (3)	N3—C7—C8—C9	-179.7 (3)
C4—S1—O1—C1	-98.6 (2)	C5—C4—C9—C8	1.7 (5)
C1—N1—N2—C2	-0.7 (4)	S1—C4—C9—C8	-174.2 (3)
C10—N1—N2—C2	178.1 (3)	C7—C8—C9—C4	-1.8 (5)
N2—N1—C1—C3	-0.5 (4)	N2—N1—C10—C15	153.3 (3)
C10—N1—C1—C3	-179.1 (3)	C1—N1—C10—C15	-28.3 (6)
N2—N1—C1—O1	178.0 (3)	N2—N1—C10—C11	-26.9 (5)
C10—N1—C1—O1	-0.5 (6)	C1—N1—C10—C11	151.6 (4)
S1—O1—C1—C3	69.1 (5)	C15—C10—C11—C12	0.9 (5)
S1—O1—C1—N1	-109.1 (3)	N1—C10—C11—C12	-179.0 (3)
N1—N2—C2—C3	1.5 (4)	C10—C11—C12—C13	0.4 (5)
N1—N2—C2—C17	-178.1 (3)	C11—C12—C13—C14	-0.8 (5)

N1—C1—C3—C2	1.3 (4)	C11—C12—C13—C16	177.2 (4)
O1—C1—C3—C2	-177.0 (4)	C12—C13—C14—C15	0.1 (5)
N2—C2—C3—C1	-1.8 (4)	C16—C13—C14—C15	-177.9 (4)
C17—C2—C3—C1	177.8 (3)	C11—C10—C15—C14	-1.6 (5)
O3—S1—C4—C9	-162.3 (3)	N1—C10—C15—C14	178.2 (3)
O2—S1—C4—C9	-25.3 (3)	C13—C14—C15—C10	1.1 (5)
O1—S1—C4—C9	88.9 (3)	N2—C2—C17—C18	-171.6 (3)
O3—S1—C4—C5	21.6 (4)	C3—C2—C17—C18	8.8 (5)
O2—S1—C4—C5	158.7 (3)	N2—C2—C17—C22	7.8 (5)
O1—S1—C4—C5	-87.2 (3)	C3—C2—C17—C22	-171.8 (3)
C9—C4—C5—C6	0.2 (5)	C22—C17—C18—C19	-1.9 (5)
S1—C4—C5—C6	176.1 (3)	C2—C17—C18—C19	177.5 (3)
C4—C5—C6—C7	-1.8 (5)	C17—C18—C19—C20	-0.4 (5)
C5—C6—C7—C8	1.6 (6)	C18—C19—C20—C21	2.1 (6)
C5—C6—C7—N3	-178.4 (3)	C19—C20—C21—C22	-1.5 (6)
O4—N3—C7—C6	175.1 (4)	C20—C21—C22—C17	-0.8 (5)
O5—N3—C7—C6	-5.4 (5)	C18—C17—C22—C21	2.5 (5)
O4—N3—C7—C8	-5.0 (5)	C2—C17—C22—C21	-177.0 (3)
O5—N3—C7—C8	174.5 (4)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C5—H5···O4 ⁱ	0.95	2.50	3.387 (5)	155

Symmetry code: (i) $x, -y+1/2, z+1/2$.