

{2-[2-Aminocyclohexyl]iminomethyl}-phenolato}dioxidovanadium(V)

Xin-Zhi Sun

College of Chemistry and Pharmaceutical Sciences, Qingdao Agricultural University, Qingdao 266109, People's Republic of China
Correspondence e-mail: xinzhi_sun@126.com


Received 9 March 2012; accepted 17 March 2012

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C-C}) = 0.015\text{ \AA}$; R factor = 0.099; wR factor = 0.203; data-to-parameter ratio = 14.8.

In the title dioxidovanadium complex, $[\text{V}(\text{C}_{13}\text{H}_{17}\text{N}_2\text{O})\text{O}_2]$, the V^{V} atom is in a square-based pyramidal coordination: the basal plane is defined by the phenolate O, imine N and amine N atoms of the tridentate Schiff base ligand, and by one oxide O atom. The apical position is occupied by the other oxide O atom. In the crystal, molecules are connected by $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots(\text{O},\text{O})$ hydrogen bonds, forming a tetramer.

Related literature

For related structures and their properties, see: Agarwal & Prasad (2006); Chohan & Sumrra (2010); Huo *et al.* (2004); Jing *et al.* (2005); Lodyga-Chruscinska *et al.* (2008); Xie *et al.* (2004); Yuan *et al.* (2009).



Experimental

Crystal data

$[\text{V}(\text{C}_{13}\text{H}_{17}\text{N}_2\text{O})\text{O}_2]$
 $M_r = 300.23$
Tetragonal, $I4_1/a$
 $a = 19.120(9)\text{ \AA}$
 $c = 15.421(3)\text{ \AA}$
 $V = 5638(4)\text{ \AA}^3$

$Z = 16$
Mo $K\alpha$ radiation
 $\mu = 0.71\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.23 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker SMART 1000 CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2000)
 $T_{\min} = 0.854$, $T_{\max} = 0.871$

19404 measured reflections
2537 independent reflections
1487 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.172$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.099$
 $wR(F^2) = 0.203$
 $S = 1.08$
2537 reflections

172 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.44\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

V1—O2	1.607 (5)	V1—N2	2.095 (5)
V1—O3	1.621 (5)	V1—N1	2.137 (6)
V1—O1	1.900 (5)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2B \cdots O2 ⁱ	0.90	2.26	3.099 (7)	155
N2—H2B \cdots O3 ⁱ	0.90	2.55	3.229 (7)	133
N2—H2A \cdots O3 ⁱⁱ	0.90	2.07	2.948 (7)	166

Symmetry codes: (i) $-y + \frac{1}{4}, x + \frac{1}{4}, -z + \frac{5}{4}$; (ii) $y - \frac{1}{4}, -x + \frac{1}{4}, -z + \frac{5}{4}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The author acknowledges the Qingdao Agricultural University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6676).

References

- Agarwal, R. K. & Prasad, S. (2006). *Rev. Inorg. Chem.* **26**, 471–492.
- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chohan, Z. H. & Sumrra, S. H. (2010). *J. Enzyme Inhib. Med. Chem.* **25**, 599–607.
- Huo, L.-H., Gao, S., Liu, J.-W., Zhao, H. & Ng, S. W. (2004). *Acta Cryst. E* **60**, m606–m608.
- Jing, B., Li, L., Wang, D., Xu, T. & Guo, X. (2005). *Acta Cryst. E* **61**, m2244–m2246.
- Lodyga-Chruscinska, E., Sanna, D., Garribba, E. & Micera, G. (2008). *Dalton Trans.* pp. 4903–4916.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xie, M.-J., Ping, Y.-S., Zheng, L.-D., Hui, J.-Z. & Peng, C. (2004). *Acta Cryst. E* **60**, m1382–m1383.
- Yuan, C. X., Lu, L. P., Gao, X. L., Wu, Y. B., Guo, M. L., Li, Y., Fu, X. Q. & Zhu, M. L. (2009). *J. Biol. Inorg. Chem.* **14**, 841–851.

supporting information

Acta Cryst. (2012). E68, m476 [https://doi.org/10.1107/S1600536812011592]

{2-[(2-Aminocyclohexyl)iminomethyl]phenolato}dioxidovanadium(V)

Xin-Zhi Sun

S1. Comment

Oxovanadium complexes with Schiff bases have been received much attention in bioinorganic chemistry (Chohan & Sumra, 2010; Agarwal & Prasad, 2006; Lodyga-Chruscinska *et al.*, 2008; Yuan *et al.*, 2009). In this paper, the title new dioxovanadium(V) complex, (I), is reported.


In the title dioxovanadium complex, Fig. 1, the V atom is in a square pyramidal coordination. The basal plane of the square pyramid is defined by one phenolate O, one imine N, and one amine N atoms of a Schiff base ligand, and by one oxo O atom. The apical position of the square pyramid is occupied by the other oxo O atom. The bond lengths (Table 1) are comparable to those observed in similar oxovanadium complexes (Xie *et al.*, 2004; Jing *et al.*, 2005; Huo *et al.*, 2004). In the crystal, adjacent molecules are linked through N—H···O and N—H···(O,O) hydrogen bonds (Table 2) to form a tetramer (Fig. 2).

S2. Experimental


To a MeOH solution (30 ml) of salicylaldehyde (0.122 g, 1.0 mmol) was added a MeOH solution (20 ml) of cyclohexane-1,2-diamine (0.114 g, 1.0 mmol) with stirring. To the above mixture was added a MeOH solution (10 ml) of VO(acac)₂ (0.265 g, 1.0 mmol) with stirring. The mixture was refluxed for 1 h, affording a clear yellow solution. This was allowed to stand at room temperature for a week and block-shaped single crystals were obtained by slow evaporation.

S3. Refinement

H atoms attached to C and N atoms were placed in geometrically idealized positions with Csp^2 —H = 0.93 Å, Csp^3 —H = 0.97–0.98 Å, N—H = 0.90 Å, and constrained to ride on their parent atoms, with $U_{iso}(\text{H}) = 1.2U_{eq}(\text{C and N})$. Apparent disorder was evident in the cyclohexyl ring.

**Figure 1**

The structure of the title compound, with displacement ellipsoids drawn at the 30% probability level for Non-H atoms.

**Figure 2**

The tetrameric unit in the crystal, with hydrogen bonds shown as double-dashed lines.

{2-[2-Aminocyclohexyl]iminomethyl}phenolato}dioxidovanadium(V)

Crystal data

 $[V(C_{13}H_{17}N_2O)O_2]$ $M_r = 300.23$ Tetragonal, $I4_1/a$ $a = 19.120 (9) \text{ \AA}$ $c = 15.421 (3) \text{ \AA}$ $V = 5638 (4) \text{ \AA}^3$ $Z = 16$ $F(000) = 2496$ $D_x = 1.415 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1125 reflections

 $\theta = 2.7\text{--}24.5^\circ$ $\mu = 0.71 \text{ mm}^{-1}$ $T = 298 \text{ K}$

Block, yellow

 $0.23 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scanAbsorption correction: multi-scan
(*SADABS*; Sheldrick, 2000) $T_{\min} = 0.854$, $T_{\max} = 0.871$

19404 measured reflections

2537 independent reflections

1487 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.172$ $\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 1.7^\circ$ $h = -23 \rightarrow 23$ $k = -23 \rightarrow 23$ $l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.099$ $wR(F^2) = 0.203$ $S = 1.08$

2537 reflections

172 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0708P)^2 + 16.0234P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.44 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
V1	0.04762 (6)	0.08630 (6)	0.66104 (7)	0.0417 (4)
N1	0.1074 (3)	0.0188 (3)	0.5792 (4)	0.0574 (17)
N2	0.0524 (3)	0.1379 (3)	0.5413 (3)	0.0424 (14)
H2A	0.0122	0.1616	0.5337	0.051*
H2B	0.0872	0.1695	0.5436	0.051*
O1	0.0854 (3)	0.0299 (3)	0.7510 (3)	0.0589 (14)

O2	-0.0330 (2)	0.0631 (2)	0.6543 (3)	0.0581 (14)
O3	0.0526 (3)	0.1602 (2)	0.7122 (3)	0.0596 (14)
C1	0.1142 (4)	-0.0770 (4)	0.6803 (5)	0.062 (2)
C2	0.0962 (4)	-0.0379 (4)	0.7536 (5)	0.057 (2)
C3	0.0907 (5)	-0.0727 (5)	0.8315 (6)	0.078 (3)
H3	0.0788	-0.0477	0.8811	0.093*
C4	0.1022 (6)	-0.1422 (6)	0.8374 (7)	0.109 (4)
H4	0.0984	-0.1643	0.8909	0.130*
C5	0.1195 (6)	-0.1808 (5)	0.7646 (8)	0.111 (4)
H5	0.1271	-0.2287	0.7692	0.133*
C6	0.1251 (5)	-0.1488 (5)	0.6875 (7)	0.094 (3)
H6	0.1365	-0.1748	0.6384	0.112*
C7	0.1238 (4)	-0.0435 (5)	0.5984 (5)	0.073 (3)
H7	0.1441	-0.0701	0.5546	0.087*
C8	0.1201 (5)	0.0398 (6)	0.4903 (6)	0.094 (3)
H8	0.1107	-0.0015	0.4546	0.112*
C9	0.0635 (4)	0.0936 (4)	0.4666 (5)	0.063 (2)
H9	0.0202	0.0670	0.4586	0.075*
C10	0.0774 (7)	0.1306 (6)	0.3813 (6)	0.110 (4)
H10A	0.0630	0.0998	0.3346	0.132*
H10B	0.0478	0.1717	0.3788	0.132*
C11	0.1431 (8)	0.1505 (8)	0.3664 (10)	0.167 (7)
H11A	0.1534	0.1903	0.4032	0.200*
H11B	0.1465	0.1661	0.3067	0.200*
C12	0.1999 (6)	0.0933 (7)	0.3823 (7)	0.115 (4)
H12A	0.1972	0.0579	0.3373	0.137*
H12B	0.2461	0.1141	0.3804	0.137*
C13	0.1873 (5)	0.0597 (7)	0.4717 (8)	0.134 (5)
H13A	0.2170	0.0187	0.4764	0.161*
H13B	0.2023	0.0926	0.5158	0.161*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
V1	0.0364 (7)	0.0422 (8)	0.0467 (7)	0.0001 (6)	0.0060 (6)	-0.0070 (6)
N1	0.056 (4)	0.066 (4)	0.050 (4)	0.032 (3)	0.012 (3)	0.003 (3)
N2	0.033 (3)	0.039 (3)	0.055 (4)	-0.001 (3)	0.000 (3)	0.006 (3)
O1	0.072 (4)	0.055 (3)	0.049 (3)	0.009 (3)	-0.008 (3)	-0.004 (3)
O2	0.054 (3)	0.049 (3)	0.071 (4)	-0.002 (2)	0.009 (3)	-0.005 (3)
O3	0.082 (4)	0.047 (3)	0.050 (3)	-0.010 (3)	0.008 (3)	-0.007 (2)
C1	0.076 (6)	0.061 (5)	0.048 (5)	0.010 (4)	0.013 (4)	-0.001 (4)
C2	0.057 (5)	0.064 (6)	0.050 (5)	0.002 (4)	-0.003 (4)	0.005 (4)
C3	0.105 (7)	0.073 (6)	0.055 (6)	0.018 (5)	0.010 (5)	0.002 (5)
C4	0.160 (11)	0.085 (8)	0.080 (7)	0.024 (7)	0.023 (7)	0.025 (7)
C5	0.167 (11)	0.051 (6)	0.115 (10)	0.037 (6)	0.030 (8)	0.003 (6)
C6	0.150 (10)	0.055 (6)	0.076 (7)	0.040 (6)	0.022 (6)	0.015 (5)
C7	0.073 (6)	0.097 (7)	0.048 (5)	0.044 (5)	0.005 (4)	-0.012 (5)
C8	0.095 (8)	0.124 (8)	0.062 (6)	0.045 (6)	0.034 (5)	0.014 (6)

C9	0.083 (6)	0.063 (5)	0.043 (4)	0.028 (5)	0.003 (4)	0.002 (4)
C10	0.161 (12)	0.115 (9)	0.054 (6)	0.037 (8)	0.029 (7)	0.036 (6)
C11	0.142 (13)	0.200 (16)	0.158 (13)	-0.015 (12)	0.024 (10)	0.123 (12)
C12	0.111 (9)	0.135 (10)	0.098 (8)	-0.012 (8)	0.034 (7)	0.026 (7)
C13	0.058 (7)	0.215 (14)	0.129 (10)	0.002 (7)	-0.005 (7)	0.079 (10)

Geometric parameters (\AA , $^\circ$)

V1—O2	1.607 (5)	C5—H5	0.9300
V1—O3	1.621 (5)	C6—H6	0.9300
V1—O1	1.900 (5)	C7—H7	0.9300
V1—N2	2.095 (5)	C8—C13	1.370 (12)
V1—N1	2.137 (6)	C8—C9	1.536 (11)
N1—C7	1.267 (9)	C8—H8	0.9800
N1—C8	1.448 (10)	C9—C10	1.517 (11)
N2—C9	1.447 (8)	C9—H9	0.9800
N2—H2A	0.9000	C10—C11	1.332 (15)
N2—H2B	0.9000	C10—H10A	0.9700
O1—C2	1.314 (8)	C10—H10B	0.9700
C1—C6	1.394 (11)	C11—C12	1.561 (17)
C1—C2	1.398 (10)	C11—H11A	0.9700
C1—C7	1.428 (11)	C11—H11B	0.9700
C2—C3	1.377 (11)	C12—C13	1.541 (13)
C3—C4	1.350 (12)	C12—H12A	0.9700
C3—H3	0.9300	C12—H12B	0.9700
C4—C5	1.383 (13)	C13—H13A	0.9700
C4—H4	0.9300	C13—H13B	0.9700
C5—C6	1.342 (13)		
O2—V1—O3	109.2 (3)	N1—C7—H7	116.7
O2—V1—O1	104.8 (2)	C1—C7—H7	116.7
O3—V1—O1	96.7 (2)	C13—C8—N1	115.6 (9)
O2—V1—N2	96.6 (2)	C13—C8—C9	115.2 (9)
O3—V1—N2	90.9 (2)	N1—C8—C9	107.0 (6)
O1—V1—N2	153.4 (2)	C13—C8—H8	106.1
O2—V1—N1	108.0 (2)	N1—C8—H8	106.1
O3—V1—N1	141.5 (3)	C9—C8—H8	106.1
O1—V1—N1	83.4 (2)	N2—C9—C10	116.3 (7)
N2—V1—N1	75.0 (2)	N2—C9—C8	107.8 (6)
C7—N1—C8	116.1 (7)	C10—C9—C8	113.3 (7)
C7—N1—V1	124.2 (5)	N2—C9—H9	106.2
C8—N1—V1	118.8 (5)	C10—C9—H9	106.2
C9—N2—V1	115.6 (4)	C8—C9—H9	106.2
C9—N2—H2A	108.4	C11—C10—C9	116.6 (11)
V1—N2—H2A	108.4	C11—C10—H10A	108.1
C9—N2—H2B	108.4	C9—C10—H10A	108.1
V1—N2—H2B	108.4	C11—C10—H10B	108.1
H2A—N2—H2B	107.4	C9—C10—H10B	108.1

C2—O1—V1	129.9 (5)	H10A—C10—H10B	107.3
C6—C1—C2	119.9 (8)	C10—C11—C12	115.4 (11)
C6—C1—C7	119.5 (8)	C10—C11—H11A	108.4
C2—C1—C7	120.5 (7)	C12—C11—H11A	108.4
O1—C2—C3	119.4 (7)	C10—C11—H11B	108.4
O1—C2—C1	122.8 (7)	C12—C11—H11B	108.4
C3—C2—C1	117.9 (8)	H11A—C11—H11B	107.5
C4—C3—C2	121.4 (9)	C13—C12—C11	108.9 (10)
C4—C3—H3	119.3	C13—C12—H12A	109.9
C2—C3—H3	119.3	C11—C12—H12A	109.9
C3—C4—C5	120.7 (10)	C13—C12—H12B	109.9
C3—C4—H4	119.7	C11—C12—H12B	109.9
C5—C4—H4	119.7	H12A—C12—H12B	108.3
C6—C5—C4	119.7 (9)	C8—C13—C12	116.7 (9)
C6—C5—H5	120.1	C8—C13—H13A	108.1
C4—C5—H5	120.1	C12—C13—H13A	108.1
C5—C6—C1	120.4 (9)	C8—C13—H13B	108.1
C5—C6—H6	119.8	C12—C13—H13B	108.1
C1—C6—H6	119.8	H13A—C13—H13B	107.3
N1—C7—C1	126.6 (7)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2B \cdots O2 ⁱ	0.90	2.26	3.099 (7)	155
N2—H2B \cdots O3 ⁱ	0.90	2.55	3.229 (7)	133
N2—H2A \cdots O3 ⁱⁱ	0.90	2.07	2.948 (7)	166

Symmetry codes: (i) $-y+1/4, x+1/4, -z+5/4$; (ii) $y-1/4, -x+1/4, -z+5/4$.