

Cinnarizinium 3,5-dinitrosalicylate

Alaloor S. Dayananda,^a Hemmige S. Yathirajan,^a Thomas Gerber,^b Eric Hosten^b and Richard Betz^{b*}

^aUniversity of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570 006, India, and ^bNelson Mandela Metropolitan University, Summerstrand Campus, Department of Chemistry, University Way, Summerstrand, PO Box 77000, Port Elizabeth 6031, South Africa

Correspondence e-mail: richard.betz@webmail.co.za


Received 6 February 2012; accepted 16 March 2012

Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.051; wR factor = 0.148; data-to-parameter ratio = 18.3.

The title compound [systematic name: 4-diphenylmethyl-1-(3-phenylprop-2-en-1-yl)piperazin-1-i um 2-carboxy-4,6-dinitrophenolate], $\text{C}_{26}\text{H}_{29}\text{N}_2^+\cdot\text{C}_7\text{H}_3\text{N}_2\text{O}_7^-$, is the dinitrosalicylate salt of a tertiary amine. Deprotonation of the carboxylic acid group occurred on the phenolic hydroxy group. The diaza-cyclohexane ring adopts a chair conformation. Intramolecular O—H···O and intermolecular C—H···O and N—H···O hydrogen bonds are observed. The N—H···O hydrogen bonds are bifurcated at the H atom and connect the cinnarizinium and 3,5-dinitrosalicylate ions together. Intermolecular C—H···O hydrogen bonds connect the components into layers perpendicular to the crystallographic a axis.

Related literature

For pharmaceutical background to cinnarizine, see: Barrett & Zolov (1960). For related structures, see: Bertolasi *et al.* (1980); Smith *et al.* (2001); Jasinski *et al.* (2011). For puckering analysis, see: Cremer & Pople (1975). For graph-set analysis of hydrogen bonds, see: Etter *et al.* (1990); Bernstein *et al.* (1995).

**Experimental***Crystal data*

$\text{C}_{26}\text{H}_{29}\text{N}_2^+\cdot\text{C}_7\text{H}_3\text{N}_2\text{O}_7^-$
 $M_r = 596.63$
Monoclinic, $P2_1/c$
 $a = 14.5648 (3)$ Å
 $b = 12.9374 (3)$ Å
 $c = 16.1619 (3)$ Å
 $\beta = 103.900 (1)^\circ$

$V = 2956.22 (11)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 200$ K
 $0.51 \times 0.26 \times 0.17$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2008)
 $T_{\min} = 0.932$, $T_{\max} = 1.000$

29552 measured reflections
7344 independent reflections
6023 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.148$
 $S = 1.03$
7344 reflections
401 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.85$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H71···O1 ⁱ	0.89 (2)	1.99 (2)	2.8105 (18)	153.5 (17)
N1—H71···O2 ⁱ	0.89 (2)	2.36 (2)	3.037 (2)	132.4 (16)
O7—H7···O1	0.84	1.75	2.507 (2)	149
C3—H3A···O6 ⁱⁱ	0.99	2.40	3.341 (2)	160
C32—H32···O5	0.95	2.52	3.453 (2)	166

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

ASD thanks University of Mysore for research facilities

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RN2101).

References

- Barrett, R. J. & Zolov, B. (1960). *J. Maine Med. Assoc.* **51**, 454–457.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bertolasi, V., Borea, P. A., Gilli, G. & Sacerdoti, M. (1980). *Acta Cryst. B* **36**, 1975–1977.
- Bruker (2008). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2010). *APEX2* and *SAINT* Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst. B* **46**, 256–262.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Jasinski, J. P., Butcher, R. J., Siddegowda, M. S., Yathirajan, H. S. & Chidan Kumar, C. S. (2011). *Acta Cryst. E* **67**, o500–o501.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.

organic compounds

- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Smith, G., Bott, R. C. & Wermuth, U. D. (2001). *Acta Cryst. E* **57**, o640–o642.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o1165–o1166 [https://doi.org/10.1107/S1600536812011518]

Cinnarizinium 3,5-dinitrosalicylate

Alaloor S. Dayananda, Hemmige S. Yathirajan, Thomas Gerber, Eric Hosten and Richard Betz

S1. Comment

Cinnarizine is an antihistamine which is mainly used for the control of nausea and vomiting due to motion sickness. A clinical evaluation of cinnarizine in various allergic disorders is published (Barrett & Zolov, 1960). The crystal structures of some related compounds such as cyclizine hydrochloride (Bertolasi *et al.*, 1980), guanidinium 3,5-dinitrosalicylate (Smith *et al.*, 2001) and cinnarizine dipicrate (Jasinski *et al.*, 2011) have been reported.

Deprotonation occurred at the phenolic hydroxy group while the carboxyl group remains in its protonated state. Protonation occurred on the nitrogen atom bearing the vinylic substituent. The 1,4-diazacyclohexane moiety adopts a 4C_1 ($^{N2}C_{N1}$) conformation (Cremer & Pople, 1975). The least-squares planes defined by the carbon atoms of the aromatic moieties on the diphenylmethyl substituent enclose an angle of 66.06 (8) $^\circ$ and intersect with the plane defined by the phenyl group bonded to the vinylic substituent at angles of 16.76 (9) $^\circ$ and 72.01 (9) $^\circ$, respectively. The least-squares planes defined by the individual phenyl groups of the cation enclose angles of 37.54 (9) $^\circ$, 53.70 (8) $^\circ$ and 79.26 (8) $^\circ$ with the corresponding plane of the carboxylic acid (Fig. 1).


In the crystal, intramolecular O—H \cdots O and intermolecular C—H \cdots O and N—H \cdots O hydrogen bonds are observed. The N—H \cdots O hydrogen bonds are bifurcated at H7. A graphical representation in terms of donor and acceptor atoms for a selection of these contacts is given in Figure 2. In terms of graph-set analysis, (Etter *et al.*, 1990; Bernstein *et al.*, 1995), the descriptor for the hydrogen bonds is *S*(6)DD on the unary level while the C—H \cdots O contacts necessitate a DD descriptor on the same level. The C—H \cdots O hydrogen bonds connect the components together into layers perpendicular to the crystallographic *a* axis (Fig. 2). The shortest intercentroid distance between two aromatic systems was measured at 4.2848 (11) Å and is apparent between the phenyl group conjugated to the vinylic system and the aromatic ring of the dinitrosalicylate anion.

S2. Experimental


Cinnarizine (3.68 g, 0.01 mol) and 3,5-dinitrosalicylic acid (2.28 g, 0.01 mol) were dissolved in hot acetone and stirred over a heating magnetic stirrer for few minutes (330 K). The resulting solution was allowed to cool slowly at room temperature. The salt formed was filtered and dried in a vacuum desiccator over phosphorous pentoxide. The compound was recrystallized from a mixture of DMSO and acetonitrile (*v*:*v* = 1:1) by slow evaporation (m.p.: 383 K).

S3. Refinement

Carbon-bound H atoms were placed in calculated positions (C—H 0.95 Å for vinylic and aromatic C atoms, C—H 0.99 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U_{eq}(C)$. The nitrogen-bound H atom was located on a difference Fourier map and their coordinates as well as isotropic displacement parameters were refined freely.

**Figure 1**


The molecular structure of the title compound, with atom labels and anisotropic displacement ellipsoids (drawn at 50% probability level).

**Figure 2**

Intermolecular contacts, viewed approximately along [1 1 1]. For clarity, only a selection of intra- and intermolecular contacts is depicted. Yellow dashed lines denote intramolecular hydrogen bonds, blue dashed lines denote intermolecular hydrogen bonds and green dashed lines indicate C–H···O contacts. Symmetry operators: ⁱ -x + 1, -y + 1, -z + 1; ⁱⁱ -x + 1, y + 1/2, -z + 1/2.

4-diphenylmethyl-1-(3-phenylprop-2-en-1-yl)piperazin-1-ium 2-carboxy-4,6-dinitrophenolate

Crystal data



$M_r = 596.63$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.5648 (3) \text{ \AA}$

$b = 12.9374 (3) \text{ \AA}$

$c = 16.1619 (3) \text{ \AA}$

$\beta = 103.900 (1)^\circ$

$V = 2956.22 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 1256$

$D_x = 1.341 \text{ Mg m}^{-3}$

Melting point: 383 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9909 reflections

$\theta = 2.3\text{--}28.3^\circ$

$\mu = 0.10 \text{ mm}^{-1}$

$T = 200 \text{ K}$

Block, yellow

$0.51 \times 0.26 \times 0.17 \text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2008)
 $T_{\min} = 0.932$, $T_{\max} = 1.000$

29552 measured reflections
7344 independent reflections
6023 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$
 $\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.0^\circ$
 $h = -19 \rightarrow 19$
 $k = -17 \rightarrow 17$
 $l = -21 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.148$
 $S = 1.03$
7344 reflections
401 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.070P)^2 + 1.6929P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.25623 (8)	0.71175 (10)	0.31330 (8)	0.0251 (2)
H71	0.2998 (14)	0.6749 (15)	0.3494 (13)	0.034 (5)*
N2	0.10014 (8)	0.56688 (9)	0.26578 (7)	0.0238 (2)
C1	0.29863 (11)	0.95171 (13)	0.41663 (10)	0.0338 (3)
H1A	0.2459	0.9812	0.3774	0.041*
C2	0.33475 (11)	0.86690 (12)	0.39167 (10)	0.0326 (3)
H2	0.3869	0.8355	0.4301	0.039*
C3	0.29836 (12)	0.81760 (12)	0.30658 (10)	0.0341 (3)
H3A	0.3508	0.8110	0.2777	0.041*
H3B	0.2495	0.8628	0.2711	0.041*
C4	0.17100 (10)	0.71680 (12)	0.34936 (10)	0.0287 (3)
H4A	0.1231	0.7629	0.3138	0.034*
H4B	0.1889	0.7459	0.4077	0.034*
C5	0.12940 (10)	0.61023 (12)	0.35209 (9)	0.0279 (3)
H5A	0.1770	0.5644	0.3883	0.034*
H5B	0.0741	0.6143	0.3775	0.034*
C6	0.18554 (10)	0.55568 (12)	0.23334 (9)	0.0267 (3)
H6A	0.1686	0.5236	0.1761	0.032*
H6B	0.2310	0.5095	0.2716	0.032*
C7	0.23147 (11)	0.65942 (12)	0.22794 (9)	0.0285 (3)
H7A	0.2895	0.6497	0.2072	0.034*
H7B	0.1877	0.7039	0.1866	0.034*
C8	0.05330 (9)	0.46543 (11)	0.26749 (9)	0.0234 (3)
H8	0.0999	0.4178	0.3043	0.028*

C11	0.33137 (12)	1.00538 (12)	0.49873 (10)	0.0328 (3)
C12	0.41578 (13)	0.97939 (14)	0.55689 (11)	0.0393 (4)
H12	0.4549	0.9265	0.5432	0.047*
C13	0.44279 (15)	1.03042 (18)	0.63443 (12)	0.0502 (5)
H13	0.4998	1.0115	0.6739	0.060*
C14	0.38744 (17)	1.10827 (18)	0.65452 (13)	0.0555 (5)
H14	0.4068	1.1435	0.7074	0.067*
C15	0.30396 (16)	1.13519 (17)	0.59790 (14)	0.0521 (5)
H15	0.2657	1.1887	0.6119	0.062*
C16	0.27599 (13)	1.08397 (14)	0.52044 (12)	0.0407 (4)
H16	0.2184	1.1027	0.4817	0.049*
C21	-0.03163 (10)	0.47510 (11)	0.30648 (9)	0.0246 (3)
C22	-0.04283 (11)	0.40380 (12)	0.36751 (10)	0.0323 (3)
H22	0.0030	0.3509	0.3849	0.039*
C23	-0.12068 (13)	0.40914 (14)	0.40346 (11)	0.0397 (4)
H23	-0.1275	0.3601	0.4454	0.048*
C24	-0.18792 (12)	0.48540 (14)	0.37840 (12)	0.0388 (4)
H24	-0.2414	0.4885	0.4024	0.047*
C25	-0.17702 (11)	0.55719 (13)	0.31830 (11)	0.0352 (3)
H25	-0.2231	0.6099	0.3011	0.042*
C26	-0.09910 (10)	0.55295 (12)	0.28272 (10)	0.0294 (3)
H26	-0.0918	0.6033	0.2420	0.035*
C31	0.02530 (10)	0.41884 (11)	0.17808 (9)	0.0251 (3)
C32	0.06680 (12)	0.32722 (12)	0.16047 (11)	0.0337 (3)
H32	0.1122	0.2937	0.2044	0.040*
C33	0.04250 (13)	0.28413 (13)	0.07907 (12)	0.0394 (4)
H33	0.0719	0.2220	0.0676	0.047*
C34	-0.02374 (13)	0.33119 (14)	0.01544 (11)	0.0386 (4)
H34	-0.0400	0.3019	-0.0401	0.046*
C35	-0.06687 (13)	0.42141 (15)	0.03242 (10)	0.0390 (4)
H35	-0.1138	0.4532	-0.0112	0.047*
C36	-0.04164 (11)	0.46560 (13)	0.11309 (10)	0.0322 (3)
H36	-0.0705	0.5284	0.1239	0.039*
O1	0.63343 (8)	0.37345 (10)	0.53526 (8)	0.0398 (3)
O2	0.58174 (10)	0.44031 (15)	0.67150 (9)	0.0619 (4)
O3	0.43582 (10)	0.48104 (13)	0.65686 (9)	0.0557 (4)
O4	0.19460 (9)	0.32142 (13)	0.44738 (11)	0.0578 (4)
O5	0.22771 (11)	0.24235 (14)	0.34093 (10)	0.0635 (4)
O6	0.54743 (11)	0.23538 (12)	0.30218 (9)	0.0525 (4)
O7	0.66612 (10)	0.28953 (12)	0.40548 (10)	0.0517 (4)
H7	0.6736	0.3061	0.4569	0.078*
N3	0.49860 (10)	0.43821 (12)	0.63196 (9)	0.0388 (3)
N4	0.25126 (10)	0.29137 (13)	0.40744 (10)	0.0436 (4)
C9	0.57578 (13)	0.27360 (14)	0.37134 (12)	0.0402 (4)
C41	0.54463 (10)	0.35631 (12)	0.50730 (10)	0.0305 (3)
C42	0.50991 (11)	0.30575 (12)	0.42633 (10)	0.0321 (3)
C43	0.41582 (12)	0.28489 (13)	0.39505 (10)	0.0332 (3)
H43	0.3951	0.2501	0.3421	0.040*

C44	0.35102 (11)	0.31446 (12)	0.44044 (10)	0.0318 (3)
C45	0.37853 (11)	0.36483 (12)	0.51759 (10)	0.0310 (3)
H45	0.3329	0.3856	0.5476	0.037*
C46	0.47329 (11)	0.38454 (12)	0.55055 (10)	0.0297 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0223 (6)	0.0289 (6)	0.0240 (6)	-0.0037 (5)	0.0053 (5)	-0.0023 (5)
N2	0.0211 (5)	0.0270 (6)	0.0237 (5)	-0.0032 (4)	0.0064 (4)	-0.0032 (4)
C1	0.0304 (7)	0.0359 (8)	0.0317 (8)	-0.0039 (6)	0.0011 (6)	-0.0002 (6)
C2	0.0314 (7)	0.0309 (7)	0.0331 (8)	-0.0073 (6)	0.0030 (6)	0.0005 (6)
C3	0.0400 (8)	0.0311 (8)	0.0322 (8)	-0.0112 (6)	0.0105 (6)	-0.0016 (6)
C4	0.0231 (6)	0.0334 (7)	0.0307 (7)	-0.0026 (5)	0.0087 (6)	-0.0078 (6)
C5	0.0251 (7)	0.0345 (7)	0.0254 (7)	-0.0053 (6)	0.0083 (5)	-0.0053 (6)
C6	0.0240 (6)	0.0307 (7)	0.0272 (7)	-0.0035 (5)	0.0095 (5)	-0.0048 (5)
C7	0.0287 (7)	0.0342 (7)	0.0235 (6)	-0.0061 (6)	0.0080 (5)	-0.0042 (6)
C8	0.0205 (6)	0.0252 (6)	0.0246 (6)	0.0003 (5)	0.0058 (5)	0.0008 (5)
C11	0.0354 (8)	0.0313 (7)	0.0310 (7)	-0.0078 (6)	0.0064 (6)	-0.0001 (6)
C12	0.0396 (9)	0.0414 (9)	0.0344 (8)	-0.0059 (7)	0.0040 (7)	0.0005 (7)
C13	0.0517 (11)	0.0614 (12)	0.0325 (9)	-0.0141 (9)	0.0000 (8)	0.0007 (8)
C14	0.0685 (14)	0.0629 (13)	0.0365 (9)	-0.0230 (11)	0.0154 (9)	-0.0156 (9)
C15	0.0600 (12)	0.0464 (11)	0.0552 (12)	-0.0103 (9)	0.0244 (10)	-0.0147 (9)
C16	0.0402 (9)	0.0359 (8)	0.0456 (9)	-0.0052 (7)	0.0095 (7)	-0.0041 (7)
C21	0.0217 (6)	0.0264 (6)	0.0253 (6)	-0.0019 (5)	0.0052 (5)	-0.0023 (5)
C22	0.0320 (7)	0.0331 (8)	0.0340 (8)	0.0033 (6)	0.0120 (6)	0.0052 (6)
C23	0.0436 (9)	0.0399 (9)	0.0423 (9)	0.0008 (7)	0.0236 (8)	0.0059 (7)
C24	0.0347 (8)	0.0390 (9)	0.0497 (10)	-0.0022 (7)	0.0241 (7)	-0.0077 (7)
C25	0.0279 (7)	0.0339 (8)	0.0446 (9)	0.0044 (6)	0.0105 (6)	-0.0057 (7)
C26	0.0281 (7)	0.0280 (7)	0.0325 (7)	0.0014 (6)	0.0084 (6)	0.0003 (6)
C31	0.0231 (6)	0.0258 (6)	0.0279 (7)	-0.0047 (5)	0.0089 (5)	-0.0018 (5)
C32	0.0337 (8)	0.0301 (7)	0.0373 (8)	0.0019 (6)	0.0084 (6)	-0.0025 (6)
C33	0.0452 (9)	0.0304 (8)	0.0451 (9)	-0.0014 (7)	0.0156 (8)	-0.0096 (7)
C34	0.0467 (9)	0.0383 (9)	0.0322 (8)	-0.0107 (7)	0.0122 (7)	-0.0100 (7)
C35	0.0401 (9)	0.0457 (9)	0.0285 (8)	-0.0005 (7)	0.0031 (7)	-0.0022 (7)
C36	0.0321 (7)	0.0341 (8)	0.0300 (7)	0.0026 (6)	0.0066 (6)	-0.0019 (6)
O1	0.0252 (5)	0.0475 (7)	0.0449 (7)	-0.0019 (5)	0.0047 (5)	0.0070 (5)
O2	0.0369 (7)	0.0965 (13)	0.0456 (8)	0.0019 (7)	-0.0034 (6)	-0.0231 (8)
O3	0.0473 (8)	0.0700 (10)	0.0462 (8)	0.0148 (7)	0.0041 (6)	-0.0177 (7)
O4	0.0247 (6)	0.0743 (10)	0.0719 (10)	0.0057 (6)	0.0065 (6)	0.0004 (8)
O5	0.0467 (8)	0.0779 (11)	0.0535 (9)	-0.0096 (8)	-0.0122 (7)	-0.0138 (8)
O6	0.0609 (9)	0.0615 (9)	0.0405 (7)	0.0067 (7)	0.0230 (7)	-0.0021 (6)
O7	0.0400 (7)	0.0633 (9)	0.0579 (8)	0.0010 (6)	0.0236 (6)	-0.0020 (7)
N3	0.0367 (7)	0.0409 (8)	0.0361 (7)	0.0042 (6)	0.0037 (6)	-0.0025 (6)
N4	0.0309 (7)	0.0474 (9)	0.0452 (8)	-0.0006 (6)	-0.0051 (6)	0.0038 (7)
C9	0.0398 (9)	0.0360 (8)	0.0489 (10)	0.0046 (7)	0.0188 (8)	0.0090 (7)
C41	0.0246 (7)	0.0280 (7)	0.0373 (8)	0.0023 (5)	0.0045 (6)	0.0099 (6)
C42	0.0336 (8)	0.0300 (7)	0.0354 (8)	0.0049 (6)	0.0135 (6)	0.0076 (6)

C43	0.0378 (8)	0.0321 (8)	0.0287 (7)	0.0010 (6)	0.0059 (6)	0.0027 (6)
C44	0.0245 (7)	0.0338 (8)	0.0338 (8)	0.0008 (6)	0.0003 (6)	0.0030 (6)
C45	0.0261 (7)	0.0327 (7)	0.0341 (8)	0.0058 (6)	0.0069 (6)	0.0022 (6)
C46	0.0292 (7)	0.0287 (7)	0.0286 (7)	0.0025 (6)	0.0018 (6)	0.0018 (6)

Geometric parameters (\AA , $\text{^{\circ}}$)

N1—C4	1.4940 (18)	C22—C23	1.395 (2)
N1—C7	1.5009 (18)	C22—H22	0.9500
N1—C3	1.5151 (19)	C23—C24	1.381 (3)
N1—H71	0.89 (2)	C23—H23	0.9500
N2—C6	1.4683 (17)	C24—C25	1.380 (3)
N2—C5	1.4685 (17)	C24—H24	0.9500
N2—C8	1.4826 (17)	C25—C26	1.391 (2)
C1—C2	1.321 (2)	C25—H25	0.9500
C1—C11	1.472 (2)	C26—H26	0.9500
C1—H1A	0.9500	C31—C36	1.388 (2)
C2—C3	1.493 (2)	C31—C32	1.391 (2)
C2—H2	0.9500	C32—C33	1.394 (2)
C3—H3A	0.9900	C32—H32	0.9500
C3—H3B	0.9900	C33—C34	1.372 (3)
C4—C5	1.511 (2)	C33—H33	0.9500
C4—H4A	0.9900	C34—C35	1.384 (3)
C4—H4B	0.9900	C34—H34	0.9500
C5—H5A	0.9900	C35—C36	1.390 (2)
C5—H5B	0.9900	C35—H35	0.9500
C6—C7	1.511 (2)	C36—H36	0.9500
C6—H6A	0.9900	O1—C41	1.2826 (19)
C6—H6B	0.9900	O2—N3	1.226 (2)
C7—H7A	0.9900	O3—N3	1.217 (2)
C7—H7B	0.9900	O4—N4	1.227 (2)
C8—C21	1.5224 (18)	O5—N4	1.224 (2)
C8—C31	1.5279 (19)	O6—C9	1.201 (2)
C8—H8	1.0000	O7—C9	1.315 (2)
C11—C16	1.394 (2)	O7—H7	0.8399
C11—C12	1.397 (2)	N3—C46	1.455 (2)
C12—C13	1.387 (3)	N4—C44	1.453 (2)
C12—H12	0.9500	C9—C42	1.514 (2)
C13—C14	1.377 (3)	C41—C46	1.432 (2)
C13—H13	0.9500	C41—C42	1.442 (2)
C14—C15	1.379 (3)	C42—C43	1.369 (2)
C14—H14	0.9500	C43—C44	1.381 (2)
C15—C16	1.388 (3)	C43—H43	0.9500
C15—H15	0.9500	C44—C45	1.378 (2)
C16—H16	0.9500	C45—C46	1.379 (2)
C21—C22	1.388 (2)	C45—H45	0.9500
C21—C26	1.395 (2)		

C4—N1—C7	109.85 (11)	C15—C16—H16	119.6
C4—N1—C3	112.16 (12)	C11—C16—H16	119.6
C7—N1—C3	110.73 (11)	C22—C21—C26	118.69 (13)
C4—N1—H71	107.2 (12)	C22—C21—C8	119.03 (13)
C7—N1—H71	109.6 (12)	C26—C21—C8	122.28 (13)
C3—N1—H71	107.1 (13)	C21—C22—C23	120.60 (15)
C6—N2—C5	107.43 (11)	C21—C22—H22	119.7
C6—N2—C8	110.63 (11)	C23—C22—H22	119.7
C5—N2—C8	110.50 (11)	C24—C23—C22	120.24 (15)
C2—C1—C11	126.91 (15)	C24—C23—H23	119.9
C2—C1—H1A	116.5	C22—C23—H23	119.9
C11—C1—H1A	116.5	C25—C24—C23	119.60 (14)
C1—C2—C3	123.84 (15)	C25—C24—H24	120.2
C1—C2—H2	118.1	C23—C24—H24	120.2
C3—C2—H2	118.1	C24—C25—C26	120.50 (15)
C2—C3—N1	112.36 (12)	C24—C25—H25	119.8
C2—C3—H3A	109.1	C26—C25—H25	119.8
N1—C3—H3A	109.1	C25—C26—C21	120.35 (14)
C2—C3—H3B	109.1	C25—C26—H26	119.8
N1—C3—H3B	109.1	C21—C26—H26	119.8
H3A—C3—H3B	107.9	C36—C31—C32	118.51 (14)
N1—C4—C5	110.38 (12)	C36—C31—C8	121.53 (13)
N1—C4—H4A	109.6	C32—C31—C8	119.96 (13)
C5—C4—H4A	109.6	C31—C32—C33	120.64 (15)
N1—C4—H4B	109.6	C31—C32—H32	119.7
C5—C4—H4B	109.6	C33—C32—H32	119.7
H4A—C4—H4B	108.1	C34—C33—C32	120.22 (16)
N2—C5—C4	110.34 (12)	C34—C33—H33	119.9
N2—C5—H5A	109.6	C32—C33—H33	119.9
C4—C5—H5A	109.6	C33—C34—C35	119.77 (15)
N2—C5—H5B	109.6	C33—C34—H34	120.1
C4—C5—H5B	109.6	C35—C34—H34	120.1
H5A—C5—H5B	108.1	C34—C35—C36	120.18 (16)
N2—C6—C7	110.95 (12)	C34—C35—H35	119.9
N2—C6—H6A	109.4	C36—C35—H35	119.9
C7—C6—H6A	109.4	C31—C36—C35	120.65 (15)
N2—C6—H6B	109.4	C31—C36—H36	119.7
C7—C6—H6B	109.4	C35—C36—H36	119.7
H6A—C6—H6B	108.0	C9—O7—H7	109.5
N1—C7—C6	111.07 (11)	O3—N3—O2	122.94 (16)
N1—C7—H7A	109.4	O3—N3—C46	117.98 (14)
C6—C7—H7A	109.4	O2—N3—C46	119.07 (14)
N1—C7—H7B	109.4	O5—N4—O4	122.97 (16)
C6—C7—H7B	109.4	O5—N4—C44	118.25 (16)
H7A—C7—H7B	108.0	O4—N4—C44	118.78 (15)
N2—C8—C21	110.99 (11)	O6—C9—O7	122.73 (17)
N2—C8—C31	110.55 (11)	O6—C9—C42	122.32 (17)
C21—C8—C31	111.42 (11)	O7—C9—C42	114.95 (16)

N2—C8—H8	107.9	O1—C41—C46	124.92 (15)
C21—C8—H8	107.9	O1—C41—C42	120.16 (15)
C31—C8—H8	107.9	C46—C41—C42	114.92 (13)
C16—C11—C12	118.36 (16)	C43—C42—C41	121.77 (14)
C16—C11—C1	119.29 (15)	C43—C42—C9	116.64 (15)
C12—C11—C1	122.35 (16)	C41—C42—C9	121.59 (15)
C13—C12—C11	120.39 (18)	C42—C43—C44	120.03 (15)
C13—C12—H12	119.8	C42—C43—H43	120.0
C11—C12—H12	119.8	C44—C43—H43	120.0
C14—C13—C12	120.41 (19)	C45—C44—C43	121.62 (14)
C14—C13—H13	119.8	C45—C44—N4	118.47 (15)
C12—C13—H13	119.8	C43—C44—N4	119.91 (15)
C13—C14—C15	120.06 (18)	C44—C45—C46	118.90 (14)
C13—C14—H14	120.0	C44—C45—H45	120.5
C15—C14—H14	120.0	C46—C45—H45	120.5
C14—C15—C16	119.9 (2)	C45—C46—C41	122.74 (14)
C14—C15—H15	120.0	C45—C46—N3	116.77 (14)
C16—C15—H15	120.0	C41—C46—N3	120.48 (14)
C15—C16—C11	120.85 (18)		
C11—C1—C2—C3	-179.12 (15)	N2—C8—C31—C36	-64.73 (17)
C1—C2—C3—N1	-114.54 (17)	C21—C8—C31—C36	59.19 (17)
C4—N1—C3—C2	63.88 (17)	N2—C8—C31—C32	115.63 (14)
C7—N1—C3—C2	-173.02 (13)	C21—C8—C31—C32	-120.45 (14)
C7—N1—C4—C5	54.46 (16)	C36—C31—C32—C33	0.8 (2)
C3—N1—C4—C5	178.06 (12)	C8—C31—C32—C33	-179.55 (14)
C6—N2—C5—C4	63.12 (15)	C31—C32—C33—C34	-0.8 (3)
C8—N2—C5—C4	-176.12 (11)	C32—C33—C34—C35	-0.4 (3)
N1—C4—C5—N2	-60.74 (15)	C33—C34—C35—C36	1.5 (3)
C5—N2—C6—C7	-61.67 (15)	C32—C31—C36—C35	0.3 (2)
C8—N2—C6—C7	177.64 (11)	C8—C31—C36—C35	-179.31 (14)
C4—N1—C7—C6	-53.14 (16)	C34—C35—C36—C31	-1.5 (3)
C3—N1—C7—C6	-177.57 (12)	O1—C41—C42—C43	179.06 (15)
N2—C6—C7—N1	57.82 (16)	C46—C41—C42—C43	-1.4 (2)
C6—N2—C8—C21	176.26 (11)	O1—C41—C42—C9	-1.3 (2)
C5—N2—C8—C21	57.42 (14)	C46—C41—C42—C9	178.26 (14)
C6—N2—C8—C31	-59.58 (14)	O6—C9—C42—C43	2.9 (2)
C5—N2—C8—C31	-178.42 (11)	O7—C9—C42—C43	-176.34 (15)
C2—C1—C11—C16	-169.51 (17)	O6—C9—C42—C41	-176.79 (16)
C2—C1—C11—C12	9.5 (3)	O7—C9—C42—C41	4.0 (2)
C16—C11—C12—C13	0.5 (3)	C41—C42—C43—C44	1.4 (2)
C1—C11—C12—C13	-178.46 (16)	C9—C42—C43—C44	-178.21 (14)
C11—C12—C13—C14	-0.9 (3)	C42—C43—C44—C45	-0.2 (2)
C12—C13—C14—C15	0.8 (3)	C42—C43—C44—N4	-179.44 (15)
C13—C14—C15—C16	-0.3 (3)	O5—N4—C44—C45	-176.60 (17)
C14—C15—C16—C11	-0.1 (3)	O4—N4—C44—C45	3.2 (2)
C12—C11—C16—C15	0.0 (3)	O5—N4—C44—C43	2.7 (2)
C1—C11—C16—C15	179.02 (17)	O4—N4—C44—C43	-177.57 (16)

N2—C8—C21—C22	−133.38 (14)	C43—C44—C45—C46	−1.0 (2)
C31—C8—C21—C22	102.95 (15)	N4—C44—C45—C46	178.23 (14)
N2—C8—C21—C26	46.96 (17)	C44—C45—C46—C41	1.0 (2)
C31—C8—C21—C26	−76.71 (17)	C44—C45—C46—N3	179.78 (14)
C26—C21—C22—C23	0.8 (2)	O1—C41—C46—C45	179.67 (15)
C8—C21—C22—C23	−178.82 (15)	C42—C41—C46—C45	0.1 (2)
C21—C22—C23—C24	0.3 (3)	O1—C41—C46—N3	1.0 (2)
C22—C23—C24—C25	−0.8 (3)	C42—C41—C46—N3	−178.57 (13)
C23—C24—C25—C26	0.2 (3)	O3—N3—C46—C45	−15.1 (2)
C24—C25—C26—C21	0.9 (2)	O2—N3—C46—C45	166.34 (17)
C22—C21—C26—C25	−1.4 (2)	O3—N3—C46—C41	163.64 (16)
C8—C21—C26—C25	178.21 (14)	O2—N3—C46—C41	−14.9 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H71···O1 ⁱ	0.89 (2)	1.99 (2)	2.8105 (18)	153.5 (17)
N1—H71···O2 ⁱ	0.89 (2)	2.36 (2)	3.037 (2)	132.4 (16)
O7—H7···O1	0.84	1.75	2.507 (2)	149
C3—H3A···O6 ⁱⁱ	0.99	2.40	3.341 (2)	160
C32—H32···O5	0.95	2.52	3.453 (2)	166

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, y+1/2, -z+1/2$.