

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Pentaaqua(4,6-dihydroxybenzene-1,3-disulfonato-κO¹)zinc pentahydrate

Zhi-Biao Zhu,^a Shan Gao,^{a†} Seik Weng Ng^{b,c} and Edward R. T. Tiekink^{b*}

^aKey Laboratory of Functional Inorganic Material Chemistry, Ministry of Education, Heilongjiang University, Harbin 150080, People's Republic of China, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: edward.tiekink@gmail.com


Received 12 March 2012; accepted 21 March 2012

Key indicators: single-crystal X-ray study; *T* = 293 K; mean $\sigma(C-C)$ = 0.003 Å; *R* factor = 0.032; *wR* factor = 0.091; data-to-parameter ratio = 13.7.

The Zn^{II} atom in the title complex, [Zn(C₆H₄O₈S₂)(H₂O)₅].5H₂O, is coordinated by five water molecules and an O atom of a 4,6-dihydroxybenzene-1,3-disulfonate dianion. The coordination geometry is distorted octahedral, with the Zn–O_{sulfonate} bond relatively long compared to the Zn–O_{water} bonds. The coordinated and lattice water molecules interact with each other and with the hydroxy groups and sulfonate ligand through O–H...O hydrogen bonds, generating a tightly held three-dimensional network.

Related literature

For related structures, see: Xie *et al.* (2010); Bakirci *et al.* (2006).



Experimental

Crystal data

[Zn(C₆H₄O₈S₂)(H₂O)₅].5H₂O
M_r = 513.74
 Triclinic, *P* $\bar{1}$
a = 7.1479 (3) Å
b = 11.8929 (5) Å
c = 12.2044 (6) Å
 α = 109.368 (1)°
 β = 104.690 (1)°

γ = 92.953 (1)°
V = 936.27 (7) Å³
Z = 2
 Mo *K* α radiation
 μ = 1.62 mm⁻¹
T = 293 K
 0.21 × 0.17 × 0.14 mm

† Additional correspondence author, e-mail: shangao67@yahoo.com.

Data collection

Rigaku R-AXIS RAPID IP diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
*T*_{min} = 0.747, *T*_{max} = 0.820

9270 measured reflections
 4257 independent reflections
 3883 reflections with *I* > 2σ(*I*)
*R*_{int} = 0.019

Refinement

R[*F*² > 2σ(*F*²)] = 0.032
wR(*F*²) = 0.091
S = 1.06
 4257 reflections
 310 parameters
 32 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}}$ = 0.67 e Å⁻³
 $\Delta\rho_{\text{min}}$ = -0.35 e Å⁻³

Table 1 Selected bond lengths (Å).

Zn–O1	2.5415 (15)	Zn–O3 _w	1.9671 (17)
Zn–O1 _w	1.9950 (15)	Zn–O4 _w	1.9588 (17)
Zn–O2 _w	1.9460 (18)	Zn–O5 _w	2.2355 (17)

Table 2 Hydrogen-bond geometry (Å, °).

<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
O7–H7...O9 ^w ⁱ	0.83 (1)	1.79 (1)	2.612 (2)	170 (3)
O8–H8...O8 ^w ⁱⁱ	0.83 (1)	1.84 (1)	2.664 (2)	172 (3)
O1 _w –H11...O2	0.84 (1)	2.09 (2)	2.852 (2)	151 (3)
O1 _w –H12...O7 ⁱⁱⁱ	0.83 (1)	2.09 (1)	2.923 (2)	175 (3)
O2 _w –H21...O6 ^{iv}	0.83 (1)	2.22 (1)	3.021 (3)	162 (3)
O2 _w –H22...O7 ^w ⁱⁱⁱ	0.83 (1)	1.88 (1)	2.710 (3)	171 (4)
O3 _w –H31...O6 ^w ^{iv}	0.84 (1)	2.01 (2)	2.765 (2)	149 (3)
O3 _w –H32...O10 ^w ^v	0.83 (1)	1.79 (1)	2.618 (3)	178 (5)
O4 _w –H41...O6 ^w ^{vi}	0.83 (1)	1.98 (1)	2.774 (2)	160 (3)
O4 _w –H42...O7 ^w ^{vii}	0.84 (1)	1.99 (1)	2.802 (3)	164 (3)
O5 _w –H51...O4 ^v	0.83 (1)	2.00 (1)	2.833 (3)	176 (4)
O5 _w –H52...O6 ^{viii}	0.84 (1)	1.97 (1)	2.802 (3)	170 (3)
O6 _w –H62...O1	0.84 (1)	2.08 (1)	2.903 (2)	168 (3)
O6 _w –H61...O4 ^{iv}	0.83 (1)	1.99 (1)	2.821 (2)	176 (3)
O7 _w –H71...O2	0.84 (1)	2.05 (1)	2.874 (2)	170 (3)
O7 _w –H72...O5 ⁱⁱ	0.83 (1)	2.33 (2)	3.034 (2)	143 (3)
O8 _w –H81...O2	0.83 (1)	2.07 (2)	2.854 (2)	157 (3)
O8 _w –H82...O3 ^{vi}	0.84 (1)	1.96 (1)	2.799 (2)	176 (3)
O9 _w –H91...O5 ^{ix}	0.83 (1)	1.95 (1)	2.747 (2)	162 (3)
O9 _w –H92...O6	0.83 (1)	1.97 (1)	2.795 (2)	173 (3)
O10 _w –H101...O8 _w	0.83 (1)	2.07 (2)	2.887 (3)	167 (4)
O10 _w –H102...O9 ^w ^{vi}	0.83 (1)	2.03 (1)	2.851 (3)	171 (5)

Symmetry codes: (i) $-x + 2, -y + 1, -z + 2$; (ii) $-x + 1, -y + 1, -z + 2$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x + 1, -y, -z + 1$; (v) $-x, -y, -z + 1$; (vi) $x - 1, y, z$; (vii) $-x, -y + 1, -z + 1$; (viii) $x - 1, y, z - 1$; (ix) $x + 1, y, z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSK and Rigaku, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work was supported by the Key Project of the Natural Science Foundation of Heilongjiang Province (grant No. ZD200903), the Key Project of the Education Bureau of Heilongjiang Province (grant Nos 12511z023 and 2011CJHB006), the Innovation Team of the Education Bureau of Heilongjiang Province (grant No. 2010 t d03),

Heilongjiang University (Hdtd2010-04), and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/3).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5482).

References

- Bakirci, H., Koner, A. L., Dickman, M. H., Kortz, U. & Nau, W. M. (2006). *Angew. Chem. Int. Ed.* **45**, 7400–7404.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSD and Rigaku (2002). *CrystalClear*. Rigaku/MSD Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Xie, B.-Y., Huang, W., Zhang, Y., Yang, R.-Q. & Xie, Y.-R. (2010). *Acta Cryst.* **E66**, m341.

supporting information

Acta Cryst. (2012). E68, m492–m493 [https://doi.org/10.1107/S1600536812012317]

Pentaaqua(4,6-dihydroxybenzene-1,3-disulfonato- κO^1)zinc pentahydrate**Zhi-Biao Zhu, Shan Gao, Seik Weng Ng and Edward R. T. Tiekink****S1. Comment**

In earlier studies (Xie *et al.*, 2010), the crystal structure determination of $[Zn(CH_3CN)(H_2O)_5](C_6H_4O_8S_2) \cdot 3H_2O$ showed that the Zn atom is octahedrally coordinated by one acetonitrile-N atom and five water molecules. The 4,6-dihydroxybenzene-1,3-disulfonate anion did not interact directly with the metal atom, instead forming hydrogen bonds to the coordinated water molecules (Xie *et al.*, 2010). When the synthesis was repeated in the absence of acetonitrile, the title compound was obtained in which the 4,6-dihydroxybenzene-1,3-disulfonate anion is now bonded to the zinc atom, Fig. 1. The covalent Zn—O_{sulfonate} bond is relatively long compared to the Zn—O_{water} bonds, Table 1. The observed coordination geometry resembles that seen in a related pentaquozinc/sulphonate structure (Bakirci *et al.*, 2006).

The coordinated and lattice water molecules interact with each other and with the sulphonate ligand through O—H \cdots O hydrogen bonds to generate a tightly-held three-dimensional network, Fig. 2 and Table 2.

S2. Experimental

Zinc nitrate (1 mmol) and 2,4-dihydroxyl-1,5-benzenedisulfonic acid (1 mmol) were dissolved in water (10 ml). The solution was filtered and then set aside for the formation of crystals. Colourless crystals were obtained after a week.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H = 0.93 Å) and were included in the refinement in the riding model approximation, with $U(H) = 1.2U_{eq}(C)$. The hydroxyl H and water H atoms were located in a difference Fourier map, and were refined with the distance restraints O—H = 0.84 ± 0.01 Å and H \cdots H = 1.37 ± 0.01 Å; $U(H) = 1.5U_{eq}(O)$.




Figure 1

The asymmetric unit of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.




Figure 2

A view in projection down the a axis of the unit-cell contents of (I). The O—H...O interactions are shown as orange dashed lines.

Pentaaqua(4,6-dihydroxybenzene-1,3-disulfonato- κO^1)zinc pentahydrate

Crystal data

$[\text{Zn}(\text{C}_6\text{H}_4\text{O}_8\text{S}_2)(\text{H}_2\text{O})_5] \cdot 5\text{H}_2\text{O}$

$M_r = 513.74$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.1479$ (3) Å

$b = 11.8929$ (5) Å

$c = 12.2044$ (6) Å

$\alpha = 109.368$ (1)°

$\beta = 104.690$ (1)°

$\gamma = 92.953$ (1)°

$V = 936.27$ (7) Å³

$Z = 2$

$F(000) = 532$

$D_x = 1.822$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8162 reflections

$\theta = 3.0\text{--}27.5^\circ$

$\mu = 1.62$ mm⁻¹

$T = 293$ K

Prism, colourless

$0.21 \times 0.17 \times 0.14$ mm

Data collection

Rigaku R-AXIS RAPID IP
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator
 ω scan

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.747$, $T_{\max} = 0.820$

9270 measured reflections

4257 independent reflections

3883 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$

$h = -8 \rightarrow 9$

$k = -15 \rightarrow 15$

$l = -15 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.091$

$S = 1.06$

4257 reflections

310 parameters

32 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0505P)^2 + 0.7681P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.67 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn	0.13086 (4)	0.18054 (2)	0.34895 (2)	0.02620 (9)
S1	0.51381 (7)	0.36541 (4)	0.60727 (4)	0.01787 (11)
S2	0.58913 (7)	0.13241 (4)	0.92703 (4)	0.01968 (12)
O1	0.4285 (2)	0.24063 (13)	0.53364 (13)	0.0256 (3)
O2	0.3656 (2)	0.44586 (14)	0.60175 (14)	0.0260 (3)
O3	0.6847 (2)	0.40513 (15)	0.57826 (14)	0.0272 (3)
O4	0.4610 (3)	0.04682 (14)	0.81323 (15)	0.0335 (4)
O5	0.5033 (2)	0.15452 (15)	1.02630 (14)	0.0277 (3)
O6	0.7809 (2)	0.09405 (16)	0.95478 (17)	0.0331 (4)
O7	0.7208 (3)	0.57857 (13)	0.82673 (14)	0.0294 (4)
H7	0.777 (4)	0.637 (2)	0.8893 (18)	0.044*
O8	0.7840 (3)	0.37304 (15)	1.10942 (14)	0.0329 (4)
H8	0.828 (5)	0.4422 (15)	1.160 (2)	0.049*
O1w	0.1481 (3)	0.34947 (14)	0.35174 (15)	0.0289 (3)
H11	0.205 (4)	0.401 (2)	0.4201 (12)	0.043*
H12	0.178 (4)	0.369 (2)	0.2982 (17)	0.043*
O2w	0.3028 (3)	0.15031 (17)	0.24487 (19)	0.0411 (4)
H21	0.300 (5)	0.0797 (13)	0.201 (3)	0.062*

H22	0.412 (3)	0.191 (2)	0.263 (3)	0.062*
O3w	0.1534 (3)	0.03068 (16)	0.38332 (19)	0.0500 (6)
H31	0.231 (5)	0.030 (3)	0.447 (2)	0.075*
H32	0.081 (5)	-0.0356 (19)	0.347 (3)	0.075*
O4w	-0.0565 (2)	0.22860 (16)	0.44222 (16)	0.0313 (4)
H41	-0.117 (4)	0.172 (2)	0.450 (3)	0.047*
H42	-0.136 (3)	0.263 (2)	0.406 (3)	0.047*
O5w	-0.1099 (3)	0.07189 (19)	0.18311 (16)	0.0389 (4)
H51	-0.214 (3)	0.041 (3)	0.186 (3)	0.058*
H52	-0.130 (4)	0.084 (3)	0.1172 (19)	0.058*
O6w	0.6621 (3)	0.05893 (17)	0.44204 (16)	0.0330 (4)
H61	0.623 (4)	0.025 (3)	0.3672 (10)	0.050*
H62	0.582 (4)	0.103 (3)	0.467 (2)	0.050*
O7w	0.3618 (3)	0.70069 (16)	0.70377 (16)	0.0320 (4)
H71	0.369 (5)	0.6284 (12)	0.668 (2)	0.048*
H72	0.365 (5)	0.712 (2)	0.7754 (12)	0.048*
O8w	0.0699 (3)	0.41514 (16)	0.71292 (15)	0.0320 (4)
H81	0.142 (3)	0.404 (3)	0.668 (2)	0.048*
H82	-0.0465 (17)	0.409 (3)	0.673 (2)	0.048*
O9w	1.1390 (3)	0.22383 (15)	0.98465 (17)	0.0345 (4)
H91	1.236 (3)	0.190 (2)	0.998 (3)	0.052*
H92	1.037 (2)	0.180 (2)	0.975 (3)	0.052*
O10w	0.0671 (5)	0.1796 (2)	0.7321 (2)	0.0718 (8)
H101	0.066 (8)	0.241 (2)	0.714 (4)	0.108*
H102	0.100 (8)	0.197 (4)	0.8071 (11)	0.108*
C1	0.5924 (3)	0.37148 (18)	0.75822 (17)	0.0186 (4)
C2	0.6887 (3)	0.47868 (18)	0.85336 (19)	0.0207 (4)
C3	0.7494 (3)	0.48013 (19)	0.97151 (18)	0.0231 (4)
H3	0.8104	0.5517	1.0346	0.028*
C4	0.7200 (3)	0.37540 (19)	0.99630 (18)	0.0218 (4)
C5	0.6239 (3)	0.26820 (18)	0.90129 (18)	0.0195 (4)
C6	0.5610 (3)	0.26795 (18)	0.78383 (18)	0.0200 (4)
H6	0.4965	0.1969	0.7210	0.024*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn	0.02933 (16)	0.02089 (14)	0.02952 (15)	0.00324 (11)	0.00921 (11)	0.00995 (11)
S1	0.0216 (2)	0.0159 (2)	0.0145 (2)	0.00053 (18)	0.00261 (17)	0.00571 (17)
S2	0.0248 (3)	0.0179 (2)	0.0185 (2)	0.00282 (19)	0.00699 (19)	0.00866 (18)
O1	0.0329 (8)	0.0185 (7)	0.0188 (7)	-0.0016 (6)	0.0006 (6)	0.0041 (6)
O2	0.0291 (8)	0.0240 (7)	0.0240 (8)	0.0080 (6)	0.0039 (6)	0.0096 (6)
O3	0.0283 (8)	0.0317 (8)	0.0239 (8)	-0.0009 (7)	0.0094 (6)	0.0123 (6)
O4	0.0508 (11)	0.0224 (8)	0.0213 (8)	-0.0085 (7)	0.0031 (7)	0.0076 (6)
O5	0.0330 (8)	0.0307 (8)	0.0260 (8)	0.0064 (7)	0.0148 (7)	0.0137 (7)
O6	0.0310 (9)	0.0364 (9)	0.0463 (10)	0.0153 (7)	0.0180 (8)	0.0262 (8)
O7	0.0471 (10)	0.0156 (7)	0.0209 (8)	-0.0056 (7)	0.0042 (7)	0.0064 (6)
O8	0.0491 (10)	0.0268 (8)	0.0151 (7)	-0.0062 (8)	-0.0021 (7)	0.0079 (6)

O1w	0.0389 (9)	0.0205 (7)	0.0278 (8)	0.0024 (7)	0.0081 (7)	0.0106 (6)
O2w	0.0415 (10)	0.0296 (9)	0.0476 (11)	0.0019 (8)	0.0255 (9)	-0.0010 (8)
O3w	0.0634 (14)	0.0237 (9)	0.0437 (11)	-0.0093 (9)	-0.0222 (10)	0.0183 (8)
O4w	0.0299 (8)	0.0332 (9)	0.0338 (9)	0.0037 (7)	0.0129 (7)	0.0133 (7)
O5w	0.0309 (9)	0.0531 (11)	0.0280 (9)	-0.0082 (8)	-0.0014 (7)	0.0181 (8)
O6w	0.0352 (9)	0.0363 (9)	0.0273 (8)	0.0055 (8)	0.0054 (7)	0.0137 (7)
O7w	0.0383 (9)	0.0302 (8)	0.0309 (9)	0.0087 (7)	0.0149 (7)	0.0109 (7)
O8w	0.0312 (9)	0.0354 (9)	0.0251 (8)	0.0032 (7)	0.0056 (7)	0.0073 (7)
O9w	0.0304 (9)	0.0253 (8)	0.0401 (10)	0.0006 (7)	0.0062 (8)	0.0053 (7)
O10w	0.103 (2)	0.0377 (12)	0.0521 (14)	-0.0216 (13)	-0.0148 (14)	0.0197 (11)
C1	0.0211 (9)	0.0187 (9)	0.0151 (9)	0.0011 (8)	0.0032 (7)	0.0065 (7)
C2	0.0241 (10)	0.0163 (9)	0.0209 (10)	0.0006 (8)	0.0053 (8)	0.0070 (8)
C3	0.0282 (11)	0.0184 (9)	0.0171 (9)	-0.0014 (8)	0.0024 (8)	0.0030 (7)
C4	0.0248 (10)	0.0231 (10)	0.0164 (9)	0.0014 (8)	0.0033 (8)	0.0079 (8)
C5	0.0233 (10)	0.0172 (9)	0.0178 (9)	0.0011 (8)	0.0045 (7)	0.0073 (7)
C6	0.0238 (10)	0.0153 (9)	0.0175 (9)	-0.0010 (8)	0.0025 (7)	0.0047 (7)

Geometric parameters (Å, °)

Zn—O1	2.5415 (15)	O3w—H32	0.833 (10)
Zn—O1w	1.9950 (15)	O4w—H41	0.833 (10)
Zn—O2w	1.9460 (18)	O4w—H42	0.838 (10)
Zn—O3w	1.9671 (17)	O5w—H51	0.831 (10)
Zn—O4w	1.9588 (17)	O5w—H52	0.840 (10)
Zn—O5w	2.2355 (17)	O6w—H61	0.831 (10)
S1—O3	1.4528 (16)	O6w—H62	0.836 (10)
S1—O1	1.4581 (15)	O7w—H71	0.837 (10)
S1—O2	1.4675 (16)	O7w—H72	0.834 (10)
S1—C1	1.760 (2)	O8w—H81	0.832 (10)
S2—O5	1.4475 (16)	O8w—H82	0.838 (10)
S2—O4	1.4577 (17)	O9w—H91	0.828 (10)
S2—O6	1.4607 (17)	O9w—H92	0.834 (10)
S2—C5	1.761 (2)	O10w—H101	0.832 (10)
O7—C2	1.355 (2)	O10w—H102	0.834 (10)
O7—H7	0.832 (10)	C1—C6	1.388 (3)
O8—C4	1.350 (2)	C1—C2	1.404 (3)
O8—H8	0.834 (10)	C2—C3	1.390 (3)
O1w—H11	0.836 (10)	C3—C4	1.394 (3)
O1w—H12	0.832 (10)	C3—H3	0.9300
O2w—H21	0.830 (10)	C4—C5	1.403 (3)
O2w—H22	0.834 (10)	C5—C6	1.388 (3)
O3w—H31	0.836 (10)	C6—H6	0.9300
O2w—Zn—O4w	171.28 (8)	Zn—O2w—H22	124 (2)
O2w—Zn—O3w	95.37 (10)	H21—O2w—H22	110.9 (17)
O4w—Zn—O3w	92.91 (10)	Zn—O3w—H31	121 (2)
O2w—Zn—O1w	87.93 (8)	Zn—O3w—H32	128 (2)
O4w—Zn—O1w	84.59 (7)	H31—O3w—H32	109.9 (17)

O3w—Zn—O1w	167.14 (8)	Zn—O4w—H41	114 (2)
O2w—Zn—O5w	86.55 (8)	Zn—O4w—H42	107 (2)
O4w—Zn—O5w	91.25 (7)	H41—O4w—H42	109.4 (16)
O3w—Zn—O5w	86.37 (8)	Zn—O5w—H51	123 (2)
O1w—Zn—O5w	106.26 (7)	Zn—O5w—H52	123 (2)
O2w—Zn—O1	89.53 (8)	H51—O5w—H52	109.2 (16)
O4w—Zn—O1	95.00 (7)	H61—O6w—H62	110.4 (16)
O3w—Zn—O1	77.08 (6)	H71—O7w—H72	109.6 (16)
O1w—Zn—O1	90.55 (6)	H81—O8w—H82	110.1 (16)
O5w—Zn—O1	162.56 (7)	H91—O9w—H92	110.9 (17)
O3—S1—O1	113.06 (10)	H101—O10w—H102	111.0 (18)
O3—S1—O2	112.15 (10)	C6—C1—C2	119.17 (18)
O1—S1—O2	110.90 (9)	C6—C1—S1	119.49 (15)
O3—S1—C1	107.07 (9)	C2—C1—S1	121.33 (15)
O1—S1—C1	106.27 (9)	O7—C2—C3	121.69 (18)
O2—S1—C1	106.94 (9)	O7—C2—C1	118.50 (18)
O5—S2—O4	112.66 (10)	C3—C2—C1	119.81 (18)
O5—S2—O6	112.17 (10)	C2—C3—C4	120.69 (19)
O4—S2—O6	110.50 (11)	C2—C3—H3	119.7
O5—S2—C5	108.45 (10)	C4—C3—H3	119.7
O4—S2—C5	105.67 (10)	O8—C4—C3	121.99 (19)
O6—S2—C5	107.00 (10)	O8—C4—C5	118.46 (18)
S1—O1—Zn	123.46 (9)	C3—C4—C5	119.53 (18)
C2—O7—H7	111 (2)	C6—C5—C4	119.40 (18)
C4—O8—H8	111 (2)	C6—C5—S2	119.21 (15)
Zn—O1w—H11	114 (2)	C4—C5—S2	121.34 (15)
Zn—O1w—H12	123 (2)	C1—C6—C5	121.38 (18)
H11—O1w—H12	110.0 (16)	C1—C6—H6	119.3
Zn—O2w—H21	118 (2)	C5—C6—H6	119.3
O3—S1—O1—Zn	-99.48 (12)	O7—C2—C3—C4	-178.1 (2)
O2—S1—O1—Zn	27.48 (13)	C1—C2—C3—C4	1.6 (3)
C1—S1—O1—Zn	143.35 (10)	C2—C3—C4—O8	177.4 (2)
O2w—Zn—O1—S1	99.04 (12)	C2—C3—C4—C5	-1.5 (3)
O4w—Zn—O1—S1	-73.50 (12)	O8—C4—C5—C6	-178.5 (2)
O3w—Zn—O1—S1	-165.34 (14)	C3—C4—C5—C6	0.5 (3)
O1w—Zn—O1—S1	11.11 (12)	O8—C4—C5—S2	-0.8 (3)
O5w—Zn—O1—S1	175.93 (17)	C3—C4—C5—S2	178.15 (17)
O3—S1—C1—C6	-122.53 (17)	O5—S2—C5—C6	-131.59 (17)
O1—S1—C1—C6	-1.4 (2)	O4—S2—C5—C6	-10.6 (2)
O2—S1—C1—C6	117.08 (18)	O6—S2—C5—C6	107.21 (18)
O3—S1—C1—C2	56.62 (19)	O5—S2—C5—C4	50.8 (2)
O1—S1—C1—C2	177.71 (17)	O4—S2—C5—C4	171.81 (18)
O2—S1—C1—C2	-63.77 (19)	O6—S2—C5—C4	-70.4 (2)
C6—C1—C2—O7	179.0 (2)	C2—C1—C6—C5	-0.4 (3)
S1—C1—C2—O7	-0.1 (3)	S1—C1—C6—C5	178.79 (16)
C6—C1—C2—C3	-0.6 (3)	C4—C5—C6—C1	0.4 (3)
S1—C1—C2—C3	-179.75 (17)	S2—C5—C6—C1	-177.26 (16)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O7—H7 \cdots O9 ^w ⁱ	0.83 (1)	1.79 (1)	2.612 (2)	170 (3)
O8—H8 \cdots O8 ^w ⁱⁱ	0.83 (1)	1.84 (1)	2.664 (2)	172 (3)
O1 ^w —H11 \cdots O2	0.84 (1)	2.09 (2)	2.852 (2)	151 (3)
O1 ^w —H12 \cdots O7 ^w ⁱⁱⁱ	0.83 (1)	2.09 (1)	2.923 (2)	175 (3)
O2 ^w —H21 \cdots O6 ^{iv}	0.83 (1)	2.22 (1)	3.021 (3)	162 (3)
O2 ^w —H22 \cdots O7 ^w ⁱⁱⁱ	0.83 (1)	1.88 (1)	2.710 (3)	171 (4)
O3 ^w —H31 \cdots O6 ^w ^{iv}	0.84 (1)	2.01 (2)	2.765 (2)	149 (3)
O3 ^w —H32 \cdots O10 ^w ^v	0.83 (1)	1.79 (1)	2.618 (3)	178 (5)
O4 ^w —H41 \cdots O6 ^w ^{vi}	0.83 (1)	1.98 (1)	2.774 (2)	160 (3)
O4 ^w —H42 \cdots O7 ^w ^{vii}	0.84 (1)	1.99 (1)	2.802 (3)	164 (3)
O5 ^w —H51 \cdots O4 ^v	0.83 (1)	2.00 (1)	2.833 (3)	176 (4)
O5 ^w —H52 \cdots O6 ^w ^{viii}	0.84 (1)	1.97 (1)	2.802 (3)	170 (3)
O6 ^w —H62 \cdots O1	0.84 (1)	2.08 (1)	2.903 (2)	168 (3)
O6 ^w —H61 \cdots O4 ^{iv}	0.83 (1)	1.99 (1)	2.821 (2)	176 (3)
O7 ^w —H71 \cdots O2	0.84 (1)	2.05 (1)	2.874 (2)	170 (3)
O7 ^w —H72 \cdots O5 ⁱⁱ	0.83 (1)	2.33 (2)	3.034 (2)	143 (3)
O8 ^w —H81 \cdots O2	0.83 (1)	2.07 (2)	2.854 (2)	157 (3)
O8 ^w —H82 \cdots O3 ^{vi}	0.84 (1)	1.96 (1)	2.799 (2)	176 (3)
O9 ^w —H91 \cdots O5 ^{ix}	0.83 (1)	1.95 (1)	2.747 (2)	162 (3)
O9 ^w —H92 \cdots O6	0.83 (1)	1.97 (1)	2.795 (2)	173 (3)
O10 ^w —H101 \cdots O8 ^w	0.83 (1)	2.07 (2)	2.887 (3)	167 (4)
O10 ^w —H102 \cdots O9 ^w ^{vi}	0.83 (1)	2.03 (1)	2.851 (3)	171 (5)

Symmetry codes: (i) $-x+2, -y+1, -z+2$; (ii) $-x+1, -y+1, -z+2$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x+1, -y, -z+1$; (v) $-x, -y, -z+1$; (vi) $x-1, y, z$; (vii) $-x, -y+1, -z+1$; (viii) $x-1, y, z-1$; (ix) $x+1, y, z$.