

## Di- $\mu$ -methanolato- $\kappa^4$ O:O-bis[bis(3-methyl-5-phenyl-1H-pyrazole- $\kappa N^2$ )-(nitrato- $\kappa O$ )copper(II)]

Xu-Guang Li,<sup>a</sup> Meng-Meng Gao<sup>b</sup> and Seik Weng Ng<sup>c\*</sup>

<sup>a</sup>Fundamental Laboratory of Life Science, Medical School of Tibet University for Nationalities, Xianyang 712082, People's Republic of China, <sup>b</sup>School of Materials Science and Engineering, Beijing Institute of Technology, Beijing 100081, People's Republic of China, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and Chemistry Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 16 March 2012; accepted 22 March 2012

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.005$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.116; data-to-parameter ratio = 16.9.

Copper nitrate in methanol solution cleaves the N—C<sub>methanol</sub> bond when reacted with 3-methyl-5-phenylpyrazole-1-methanol to yield the centrosymmetric dinuclear title compound,  $[Cu_2(CH_3O)_2(NO_3)_2(C_{10}H_{10}N_2)_4]$ , in which the Cu<sup>II</sup> atom is linked to a nitrate ion, two methanolate ions and two pyrazole ligands in a distorted square-pyramidal environment. The O atom of the nitrate anion occupies the apical site. The crystal structure features intramolecular N—H···O hydrogen bonds.

### Related literature

For a related structure, see: He & Sykes (2007). For the synthesis of 3-methyl-5-phenylpyrazole-1-methanol, see: Zhu *et al.* (2004).



### Experimental

#### Crystal data

$[Cu_2(CH_3O)_2(NO_3)_2(C_{10}H_{10}N_2)_4]$	$\gamma = 95.853 (2)^\circ$
$M_r = 945.97$	$V = 1106.85 (18) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 8.3896 (8) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 11.2569 (11) \text{ \AA}$	$\mu = 1.02 \text{ mm}^{-1}$
$c = 12.7200 (12) \text{ \AA}$	$T = 293 \text{ K}$
$\alpha = 106.120 (2)^\circ$	$0.12 \times 0.11 \times 0.10 \text{ mm}$
$\beta = 103.025 (2)^\circ$	

#### Data collection

Bruker SMART-1000 diffractometer	6768 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	4898 independent reflections
$T_{\min} = 0.887$ , $T_{\max} = 0.905$	3461 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.015$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.116$	$\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$
$S = 1.02$	$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$
4898 reflections	
290 parameters	
2 restraints	

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N2—H2···O2 <sup>i</sup>	0.87 (1)	2.26 (2)	3.012 (3)	144 (3)
N4—H4···O3	0.87 (1)	2.17 (2)	3.022 (4)	166 (3)

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors acknowledge support from the Scientific Research Project of Higher Education of Inner Mongolia (grant No. NJ09204) and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5488).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2001). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- He, H. & Sykes, A. G. (2007). *Acta Cryst. E* **63**, m2448.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhu, W.-R., Hu, P.-Z. & Li, M.-Y. (2004). *Chin. J. Synth. Chem.* **12**, 28–30.

# supporting information

*Acta Cryst.* (2012). E68, m503 [https://doi.org/10.1107/S160053681201241X]

## Di- $\mu$ -methanolato- $\kappa^4$ O:O-bis[bis(3-methyl-5-phenyl-1*H*-pyrazole- $\kappa N^2$ )(nitrato- $\kappa O$ )copper(II)]

Xu-Guang Li, Meng-Meng Gao and Seik Weng Ng

### S1. Comment

Copper nitrate in methanol solution cleaves the  $N$ —C<sub>methanol</sub> bond when reacted with 3-methyl-5-phenylpyrazole-1-methanol to yield the dinuclear title compound (Scheme I, Fig. 1). The molecule lies on a center-of-inversion; the Cu<sub>II</sub> atom is linked to a nitrate ion, two methanolate ions and two of the pyrazole ligands in a square-pyramidal environment. In the perchlorate analog, [Cu(OCH<sub>3</sub>)(C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>)<sub>2</sub>]<sub>2</sub>(ClO<sub>4</sub>)<sub>2</sub>, the counterion is not connected to the copper atom, whose geometry is a square pyramid. The compound was synthesized by directly reacting 3-methyl-5-phenylpyrazole with copper perchlorate in methanol medium (He & Sykes, 2007).


### S2. Experimental

3-Methyl-5-phenylpyrazole-1-methanol was synthesized by using a literature procedure (Zhu *et al.*, 2004). The ligand (0.065 g, 0.4 mmol) was dissolved in dichloromethane (10 mol) and this was mixed with a methanol solution (10 ml) of copper nitrate trihydrate (0.024 g, 0.1 mmol). The clear blue solution was filtered and then set aside for the growth of deep blue crystals. CH&N elemental analysis. Calc. for C<sub>42</sub>H<sub>46</sub>Cu<sub>2</sub>N<sub>10</sub>O<sub>8</sub>: C 53.33, H 4.90; N 14.81%. Found: C 56.36, H 5.18, N 15.02%.

### S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.96 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to 1.2 to 1.5 $U(C)$ .

The amino H-atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H 0.88±0.01 Å; their temperature factors were freely refined.

**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of centrosymmetric  $[\text{Cu}(\text{OCH}_3)(\text{NO}_3)(\text{C}_{10}\text{H}_{10}\text{N}_2)_2]_2$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Inversion-related atoms are not labeled.

### Di- $\mu$ -methanolato- $\kappa^4\text{O:O}$ -bis[3-methyl-5-phenyl-1*H*-pyrazole- $\kappa\text{N}^2$ ](nitrato- $\kappa\text{O}$ )copper(II)]

#### Crystal data

$[\text{Cu}_2(\text{CH}_3\text{O})_2(\text{NO}_3)_2(\text{C}_{10}\text{H}_{10}\text{N}_2)_4]$   
 $M_r = 945.97$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 8.3896 (8)$  Å  
 $b = 11.2569 (11)$  Å  
 $c = 12.7200 (12)$  Å  
 $\alpha = 106.120 (2)^\circ$   
 $\beta = 103.025 (2)^\circ$   
 $\gamma = 95.853 (2)^\circ$   
 $V = 1106.85 (18)$  Å<sup>3</sup>

$Z = 1$   
 $F(000) = 490$   
 $D_x = 1.419 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 2170 reflections  
 $\theta = 2.5\text{--}24.5^\circ$   
 $\mu = 1.02 \text{ mm}^{-1}$   
 $T = 293$  K  
Prism, blue  
 $0.12 \times 0.11 \times 0.10$  mm

#### Data collection

Bruker SMART-1000  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.887$ ,  $T_{\max} = 0.905$

6768 measured reflections  
4898 independent reflections  
3461 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.015$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 1.9^\circ$   
 $h = -10 \rightarrow 10$   
 $k = -12 \rightarrow 14$   
 $l = -16 \rightarrow 10$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.116$   
 $S = 1.02$   
4898 reflections  
290 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0516P)^2 + 0.4958P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.54841 (4)	0.37278 (3)	0.47584 (3)	0.04912 (14)
O1	0.7401 (3)	0.4733 (3)	0.3606 (2)	0.0769 (7)
O2	0.6684 (3)	0.6571 (2)	0.3847 (2)	0.0768 (7)
O3	0.5511 (3)	0.5059 (2)	0.23096 (17)	0.0703 (6)
O4	0.3871 (2)	0.46286 (18)	0.41469 (14)	0.0464 (5)
N1	0.7022 (3)	0.3017 (2)	0.57615 (18)	0.0508 (6)
N2	0.6938 (3)	0.3252 (2)	0.68576 (19)	0.0507 (6)
N3	0.4722 (3)	0.2251 (2)	0.3377 (2)	0.0576 (6)
N4	0.4708 (3)	0.2394 (3)	0.2345 (2)	0.0573 (6)
N5	0.6566 (3)	0.5464 (3)	0.3264 (2)	0.0535 (6)
C1	0.9035 (5)	0.2356 (4)	0.4677 (3)	0.0793 (11)
H1A	0.8149	0.2370	0.4056	0.119*
H1B	0.9330	0.1535	0.4525	0.119*
H1C	0.9983	0.2965	0.4764	0.119*
C2	0.8479 (4)	0.2663 (3)	0.5748 (2)	0.0519 (7)
C3	0.9314 (4)	0.2661 (3)	0.6822 (2)	0.0545 (7)
H3	1.0350	0.2440	0.7028	0.065*
C4	0.8307 (3)	0.3052 (3)	0.7523 (2)	0.0469 (6)
C5	0.8519 (3)	0.3269 (3)	0.8744 (2)	0.0493 (7)
C6	0.7739 (4)	0.4139 (3)	0.9353 (3)	0.0603 (8)
H6	0.7085	0.4603	0.8990	0.072*
C7	0.7932 (4)	0.4319 (4)	1.0499 (3)	0.0730 (10)
H7	0.7408	0.4906	1.0901	0.088*
C8	0.8892 (4)	0.3639 (4)	1.1049 (3)	0.0753 (11)
H8	0.8993	0.3748	1.1815	0.090*
C9	0.9694 (4)	0.2801 (4)	1.0461 (3)	0.0717 (10)
H9	1.0362	0.2353	1.0835	0.086*
C10	0.9524 (4)	0.2610 (3)	0.9315 (3)	0.0579 (8)
H10	1.0082	0.2040	0.8925	0.069*
C11	0.3473 (6)	0.0666 (4)	0.4104 (3)	0.0996 (15)
H11A	0.4399	0.1010	0.4763	0.149*
H11B	0.3330	-0.0234	0.3870	0.149*
H11C	0.2484	0.0925	0.4281	0.149*
C12	0.3794 (5)	0.1132 (3)	0.3157 (3)	0.0659 (9)
C13	0.3201 (5)	0.0571 (3)	0.1989 (3)	0.0731 (10)
H13	0.2534	-0.0214	0.1618	0.088*
C14	0.3786 (4)	0.1393 (3)	0.1488 (3)	0.0582 (8)
C15	0.3536 (4)	0.1299 (3)	0.0279 (3)	0.0616 (8)

C16	0.4062 (5)	0.2286 (4)	-0.0078 (3)	0.0764 (10)
H16	0.4597	0.3052	0.0456	0.092*
C17	0.3799 (5)	0.2146 (5)	-0.1229 (3)	0.0884 (12)
H17	0.4166	0.2815	-0.1461	0.106*
C18	0.3000 (6)	0.1023 (5)	-0.2022 (3)	0.0935 (14)
H18	0.2828	0.0925	-0.2792	0.112*
C19	0.2461 (7)	0.0050 (5)	-0.1673 (3)	0.1131 (18)
H19	0.1906	-0.0708	-0.2212	0.136*
C20	0.2722 (6)	0.0169 (4)	-0.0533 (3)	0.0944 (14)
H20	0.2353	-0.0507	-0.0310	0.113*
C21	0.2225 (4)	0.4090 (3)	0.3499 (3)	0.0620 (8)
H21A	0.1667	0.4727	0.3290	0.093*
H21B	0.1647	0.3730	0.3940	0.093*
H21C	0.2241	0.3446	0.2825	0.093*
H2	0.602 (2)	0.349 (3)	0.698 (3)	0.061 (10)*
H4	0.510 (4)	0.3130 (18)	0.231 (3)	0.079 (12)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.0482 (2)	0.0578 (2)	0.03230 (18)	0.00578 (16)	-0.00269 (13)	0.01163 (15)
O1	0.0655 (15)	0.0950 (19)	0.0817 (17)	0.0317 (14)	0.0191 (13)	0.0398 (15)
O2	0.0714 (16)	0.0622 (15)	0.0842 (17)	0.0014 (13)	0.0201 (13)	0.0070 (14)
O3	0.0657 (14)	0.0978 (18)	0.0423 (12)	0.0083 (13)	0.0082 (10)	0.0205 (12)
O4	0.0393 (10)	0.0574 (12)	0.0334 (9)	0.0009 (9)	-0.0025 (8)	0.0123 (9)
N1	0.0506 (14)	0.0663 (16)	0.0343 (11)	0.0126 (12)	0.0056 (10)	0.0174 (11)
N2	0.0439 (14)	0.0703 (17)	0.0409 (12)	0.0157 (13)	0.0089 (11)	0.0217 (12)
N3	0.0634 (16)	0.0554 (16)	0.0418 (13)	0.0061 (13)	-0.0030 (11)	0.0109 (12)
N4	0.0658 (17)	0.0528 (16)	0.0400 (13)	0.0032 (13)	0.0029 (12)	0.0047 (12)
N5	0.0435 (14)	0.0709 (18)	0.0501 (14)	0.0076 (13)	0.0165 (11)	0.0225 (14)
C1	0.074 (2)	0.114 (3)	0.0451 (18)	0.029 (2)	0.0173 (17)	0.011 (2)
C2	0.0500 (17)	0.0613 (19)	0.0406 (15)	0.0102 (14)	0.0093 (13)	0.0118 (14)
C3	0.0439 (16)	0.068 (2)	0.0475 (16)	0.0148 (15)	0.0051 (13)	0.0148 (15)
C4	0.0457 (15)	0.0516 (17)	0.0411 (14)	0.0053 (13)	0.0041 (12)	0.0180 (13)
C5	0.0423 (15)	0.0620 (19)	0.0402 (14)	0.0003 (13)	0.0032 (12)	0.0199 (14)
C6	0.0504 (18)	0.081 (2)	0.0470 (16)	0.0099 (16)	0.0088 (14)	0.0196 (16)
C7	0.053 (2)	0.107 (3)	0.0500 (18)	0.0045 (19)	0.0137 (15)	0.0134 (19)
C8	0.056 (2)	0.119 (3)	0.0397 (16)	-0.012 (2)	0.0029 (15)	0.025 (2)
C9	0.060 (2)	0.095 (3)	0.0545 (19)	0.0006 (19)	-0.0066 (16)	0.037 (2)
C10	0.0499 (17)	0.073 (2)	0.0492 (16)	0.0066 (15)	0.0039 (13)	0.0259 (16)
C11	0.140 (4)	0.077 (3)	0.071 (2)	-0.007 (3)	0.002 (3)	0.035 (2)
C12	0.078 (2)	0.0510 (19)	0.0567 (19)	0.0079 (17)	-0.0037 (17)	0.0158 (16)
C13	0.085 (3)	0.0482 (19)	0.062 (2)	-0.0004 (18)	-0.0088 (18)	0.0060 (16)
C14	0.0610 (19)	0.0508 (18)	0.0482 (17)	0.0117 (15)	-0.0006 (14)	0.0035 (14)
C15	0.061 (2)	0.064 (2)	0.0430 (16)	0.0170 (16)	-0.0007 (14)	-0.0006 (15)
C16	0.073 (2)	0.088 (3)	0.0516 (19)	-0.003 (2)	0.0066 (17)	0.0086 (19)
C17	0.078 (3)	0.122 (4)	0.059 (2)	0.008 (3)	0.017 (2)	0.024 (2)
C18	0.098 (3)	0.125 (4)	0.045 (2)	0.033 (3)	0.015 (2)	0.006 (2)

C19	0.162 (5)	0.090 (3)	0.048 (2)	0.022 (3)	0.000 (3)	-0.017 (2)
C20	0.138 (4)	0.065 (2)	0.052 (2)	0.007 (2)	0.003 (2)	-0.0041 (18)
C21	0.0456 (17)	0.071 (2)	0.0514 (17)	-0.0038 (15)	-0.0093 (13)	0.0134 (16)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

Cu1—O4 <sup>i</sup>	1.9185 (19)	C7—C8	1.377 (5)
Cu1—O4	1.9256 (18)	C7—H7	0.9300
Cu1—N3	1.979 (2)	C8—C9	1.366 (5)
Cu1—N1	1.992 (2)	C8—H8	0.9300
Cu1—Cu1 <sup>i</sup>	2.9939 (8)	C9—C10	1.385 (4)
O1—N5	1.239 (3)	C9—H9	0.9300
O2—N5	1.242 (3)	C10—H10	0.9300
O3—N5	1.261 (3)	C11—C12	1.505 (5)
O4—C21	1.412 (3)	C11—H11A	0.9600
O4—Cu1 <sup>i</sup>	1.9185 (19)	C11—H11B	0.9600
N1—C2	1.326 (4)	C11—H11C	0.9600
N1—N2	1.364 (3)	C12—C13	1.391 (4)
N2—C4	1.344 (3)	C13—C14	1.370 (5)
N2—H2	0.873 (10)	C13—H13	0.9300
N3—C12	1.333 (4)	C14—C15	1.478 (4)
N3—N4	1.363 (3)	C15—C16	1.380 (5)
N4—C14	1.348 (4)	C15—C20	1.388 (5)
N4—H4	0.874 (10)	C16—C17	1.392 (5)
C1—C2	1.501 (4)	C16—H16	0.9300
C1—H1A	0.9600	C17—C18	1.370 (6)
C1—H1B	0.9600	C17—H17	0.9300
C1—H1C	0.9600	C18—C19	1.365 (6)
C2—C3	1.388 (4)	C18—H18	0.9300
C3—C4	1.379 (4)	C19—C20	1.383 (6)
C3—H3	0.9300	C19—H19	0.9300
C4—C5	1.469 (4)	C20—H20	0.9300
C5—C6	1.387 (4)	C21—H21A	0.9600
C5—C10	1.393 (4)	C21—H21B	0.9600
C6—C7	1.383 (4)	C21—H21C	0.9600
C6—H6	0.9300		
O4 <sup>i</sup> —Cu1—O4	77.69 (8)	C8—C7—H7	119.6
O4 <sup>i</sup> —Cu1—N3	166.83 (9)	C6—C7—H7	119.6
O4—Cu1—N3	91.96 (9)	C9—C8—C7	119.4 (3)
O4 <sup>i</sup> —Cu1—N1	91.67 (9)	C9—C8—H8	120.3
O4—Cu1—N1	165.30 (9)	C7—C8—H8	120.3
N3—Cu1—N1	99.88 (10)	C8—C9—C10	120.8 (3)
O4 <sup>i</sup> —Cu1—Cu1 <sup>i</sup>	38.93 (5)	C8—C9—H9	119.6
O4—Cu1—Cu1 <sup>i</sup>	38.76 (5)	C10—C9—H9	119.6
N3—Cu1—Cu1 <sup>i</sup>	130.23 (7)	C5—C10—C9	120.1 (3)
N1—Cu1—Cu1 <sup>i</sup>	129.86 (7)	C5—C10—H10	119.9
C21—O4—Cu1 <sup>i</sup>	124.42 (19)	C9—C10—H10	119.9

C21—O4—Cu1	125.07 (19)	C12—C11—H11A	109.5
Cu1 <sup>i</sup> —O4—Cu1	102.31 (8)	C12—C11—H11B	109.5
C2—N1—N2	105.4 (2)	H11A—C11—H11B	109.5
C2—N1—Cu1	133.5 (2)	C12—C11—H11C	109.5
N2—N1—Cu1	117.77 (18)	H11A—C11—H11C	109.5
C4—N2—N1	111.9 (2)	H11B—C11—H11C	109.5
C4—N2—H2	133 (2)	N3—C12—C13	109.7 (3)
N1—N2—H2	115 (2)	N3—C12—C11	120.9 (3)
C12—N3—N4	105.7 (2)	C13—C12—C11	129.3 (3)
C12—N3—Cu1	132.4 (2)	C14—C13—C12	107.1 (3)
N4—N3—Cu1	119.6 (2)	C14—C13—H13	126.5
C14—N4—N3	111.5 (3)	C12—C13—H13	126.5
C14—N4—H4	128 (2)	N4—C14—C13	106.0 (3)
N3—N4—H4	119 (2)	N4—C14—C15	123.3 (3)
O1—N5—O2	122.3 (3)	C13—C14—C15	130.7 (3)
O1—N5—O3	119.0 (3)	C16—C15—C20	118.7 (3)
O2—N5—O3	118.6 (3)	C16—C15—C14	123.0 (3)
C2—C1—H1A	109.5	C20—C15—C14	118.3 (3)
C2—C1—H1B	109.5	C15—C16—C17	120.6 (4)
H1A—C1—H1B	109.5	C15—C16—H16	119.7
C2—C1—H1C	109.5	C17—C16—H16	119.7
H1A—C1—H1C	109.5	C18—C17—C16	120.0 (4)
H1B—C1—H1C	109.5	C18—C17—H17	120.0
N1—C2—C3	110.3 (3)	C16—C17—H17	120.0
N1—C2—C1	120.8 (3)	C19—C18—C17	119.5 (4)
C3—C2—C1	128.8 (3)	C19—C18—H18	120.3
C4—C3—C2	106.6 (3)	C17—C18—H18	120.3
C4—C3—H3	126.7	C18—C19—C20	121.3 (4)
C2—C3—H3	126.7	C18—C19—H19	119.4
N2—C4—C3	105.7 (2)	C20—C19—H19	119.4
N2—C4—C5	121.9 (3)	C19—C20—C15	119.8 (4)
C3—C4—C5	132.4 (3)	C19—C20—H20	120.1
C6—C5—C10	118.7 (3)	C15—C20—H20	120.1
C6—C5—C4	120.9 (3)	O4—C21—H21A	109.5
C10—C5—C4	120.4 (3)	O4—C21—H21B	109.5
C7—C6—C5	120.2 (3)	H21A—C21—H21B	109.5
C7—C6—H6	119.9	O4—C21—H21C	109.5
C5—C6—H6	119.9	H21A—C21—H21C	109.5
C8—C7—C6	120.7 (4)	H21B—C21—H21C	109.5
O4 <sup>i</sup> —Cu1—O4—C21	-149.3 (3)	C2—C3—C4—C5	-178.2 (3)
N3—Cu1—O4—C21	38.9 (2)	N2—C4—C5—C6	-27.0 (4)
N1—Cu1—O4—C21	-104.9 (4)	C3—C4—C5—C6	151.7 (3)
Cu1 <sup>i</sup> —Cu1—O4—C21	-149.3 (3)	N2—C4—C5—C10	153.5 (3)
O4 <sup>i</sup> —Cu1—O4—Cu1 <sup>i</sup>	0.0	C3—C4—C5—C10	-27.7 (5)
N3—Cu1—O4—Cu1 <sup>i</sup>	-171.78 (10)	C10—C5—C6—C7	-1.5 (5)
N1—Cu1—O4—Cu1 <sup>i</sup>	44.5 (4)	C4—C5—C6—C7	179.0 (3)
O4 <sup>i</sup> —Cu1—N1—C2	-104.9 (3)	C5—C6—C7—C8	-0.2 (5)

O4—Cu1—N1—C2	−148.1 (3)	C6—C7—C8—C9	1.7 (5)
N3—Cu1—N1—C2	68.7 (3)	C7—C8—C9—C10	−1.4 (5)
Cu1 <sup>i</sup> —Cu1—N1—C2	−113.3 (3)	C6—C5—C10—C9	1.8 (5)
O4 <sup>i</sup> —Cu1—N1—N2	50.9 (2)	C4—C5—C10—C9	−178.7 (3)
O4—Cu1—N1—N2	7.7 (5)	C8—C9—C10—C5	−0.4 (5)
N3—Cu1—N1—N2	−135.5 (2)	N4—N3—C12—C13	0.2 (4)
Cu1 <sup>i</sup> —Cu1—N1—N2	42.5 (2)	Cu1—N3—C12—C13	162.2 (3)
C2—N1—N2—C4	−0.1 (3)	N4—N3—C12—C11	−177.8 (3)
Cu1—N1—N2—C4	−162.1 (2)	Cu1—N3—C12—C11	−15.7 (5)
O4 <sup>i</sup> —Cu1—N3—C12	−141.7 (4)	N3—C12—C13—C14	−0.5 (4)
O4—Cu1—N3—C12	−103.9 (3)	C11—C12—C13—C14	177.2 (4)
N1—Cu1—N3—C12	67.3 (3)	N3—N4—C14—C13	−0.6 (4)
Cu1 <sup>i</sup> —Cu1—N3—C12	−110.6 (3)	N3—N4—C14—C15	179.3 (3)
O4 <sup>i</sup> —Cu1—N3—N4	18.4 (6)	C12—C13—C14—N4	0.7 (4)
O4—Cu1—N3—N4	56.2 (2)	C12—C13—C14—C15	−179.2 (3)
N1—Cu1—N3—N4	−132.6 (2)	N4—C14—C15—C16	−7.8 (5)
Cu1 <sup>i</sup> —Cu1—N3—N4	49.4 (3)	C13—C14—C15—C16	172.1 (4)
C12—N3—N4—C14	0.3 (4)	N4—C14—C15—C20	172.8 (3)
Cu1—N3—N4—C14	−164.5 (2)	C13—C14—C15—C20	−7.3 (6)
N2—N1—C2—C3	0.5 (3)	C20—C15—C16—C17	−0.8 (6)
Cu1—N1—C2—C3	158.4 (2)	C14—C15—C16—C17	179.8 (3)
N2—N1—C2—C1	−178.3 (3)	C15—C16—C17—C18	0.5 (6)
Cu1—N1—C2—C1	−20.5 (5)	C16—C17—C18—C19	0.3 (7)
N1—C2—C3—C4	−0.8 (4)	C17—C18—C19—C20	−0.8 (8)
C1—C2—C3—C4	178.0 (3)	C18—C19—C20—C15	0.5 (8)
N1—N2—C4—C3	−0.4 (3)	C16—C15—C20—C19	0.3 (7)
N1—N2—C4—C5	178.6 (3)	C14—C15—C20—C19	179.7 (4)
C2—C3—C4—N2	0.7 (3)		

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2 $\cdots$ O2 <sup>i</sup>	0.87 (1)	2.26 (2)	3.012 (3)	144 (3)
N4—H4 $\cdots$ O3	0.87 (1)	2.17 (2)	3.022 (4)	166 (3)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .