

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(4-Chlorobutanoyl)-3-(2-chlorophenyl)thiourea

M. Sukeri M. Yusof,^a Nur Farhana Embong,^a Bohari M. Yamin^b and Nurziana Ngah^{c*}^aDepartment of Chemical Sciences, Faculty of Science and Technology, Universiti Malaysia Terengganu, Menggabang Telipot, 21030 Kuala Terengganu, Malaysia,^bSchool of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43600 Bangi Selangor, Malaysia, and ^cKulliyah of Science, International Islamic University Malaysia, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia

Correspondence e-mail: nurziana@iiium.edu.my


Received 13 April 2012; accepted 20 April 2012

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.039; wR factor = 0.109; data-to-parameter ratio = 15.4.

The asymmetric unit of the title compound, $\text{C}_{11}\text{H}_{12}\text{Cl}_2\text{N}_2\text{OS}$, contains two crystallographically independent molecules with different conformations: the benzene ring and the thiourea fragment form dihedral angles of 74.32 (11) and 89.39 (11)°. One amino group in each molecule is involved in intramolecular $\text{N}-\text{H}\cdots\text{O}$ and intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonding: the latter links pairs of independent molecules into dimers. In the crystal, weak $\text{N}-\text{H}\cdots\text{S}$ interactions link these dimers into chains propagating along the c axis.

Related literature

For a related structure, see: Yusof *et al.* (2011). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{11}\text{H}_{12}\text{Cl}_2\text{N}_2\text{OS}$
 $M_r = 291.19$ Monoclinic, $P2_1/c$
 $a = 14.396$ (3) Å $b = 10.941$ (2) Å
 $c = 18.093$ (4) Å
 $\beta = 109.399$ (4)°
 $V = 2688.0$ (9) Å³
 $Z = 8$ Mo $K\alpha$ radiation
 $\mu = 0.62$ mm⁻¹
 $T = 298$ K
 $0.42 \times 0.41 \times 0.39$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.780$, $T_{\max} = 0.793$ 14457 measured reflections
4727 independent reflections
3802 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.109$
 $S = 1.03$
4727 reflections307 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.40$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.33$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O1}$	0.86	2.01	2.675 (3)	133
$\text{N3}-\text{H3A}\cdots\text{O2}$	0.86	1.98	2.652 (3)	134
$\text{N1}-\text{H1A}\cdots\text{O2}$	0.86	2.37	3.070 (3)	138
$\text{N3}-\text{H3A}\cdots\text{O1}$	0.86	2.37	3.073 (3)	139
$\text{N2}-\text{H2A}\cdots\text{S2}^{\text{i}}$	0.86	2.58	3.404 (2)	160
$\text{N4}-\text{H4A}\cdots\text{S1}^{\text{ii}}$	0.86	2.58	3.433 (2)	174

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL, PARST (Nardelli, 1995) and PLATON (Spek, 2009).

MSMY, BMY and NN acknowledge Universiti Malaysia Terengganu, Universiti Kebangsaan Malaysia and the International Islamic University Malaysia for facilities and funding to support the synthetic part of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5287).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Yusof, M. S. M., Embong, N. F., Othman, E. A. & Yamin, B. M. (2011). *Acta Cryst.* **E67**, o1849.

supporting information

Acta Cryst. (2012). E68, o1536 [doi:10.1107/S160053681201759X]

1-(4-Chlorobutanoyl)-3-(2-chlorophenyl)thiourea

M. Sukeri M. Yusof, Nur Farhana Embong, Bohari M. Yamin and Nurziana Ngah

S1. Comment


The asymmetric unit of the title compound, (I), contains two crystallographically independent molecules with different conformations. The title molecule is similar to the previously reported *N*-(4-chlorobutanoyl)-*N'*-(2-fluorophenyl)thiourea (Yusof *et al.*, 2011) except the fluorine atom at *ortho* position of benzene ring is substituted by the chlorine atom at the same position. The bond lengths and angles are in normal ranges (Allen *et al.*, 1987). The benzene [(C1—C6) & (C12—C17)] and thiourea [(N1/N2/C7/S1/C6) & (N3/N4/C18/S2/C17)] fragments are each planar with maximum deviation is 0.047 (2) Å for atom N4 from the mean plane. In each independent molecule, the benzene and thiourea fragments make dihedral angles of 74.32 (11)° and 89.39 (11)°, respectively and comparable to those reported by Yusof *et al.*, (2011). One amino group in each molecule, N1—H1A and N3—H3A, respectively, is involved in bifurcated N—H···O hydrogen-bonding (Table 1) - intra and intermolecular ones, respectively, and the latter ones link two independent molecules into dimer. In the crystal, weak N—H···S interactions (Table 1) link further these dimers into chains propagated along *c* axis.

S2. Experimental

30 ml Acetone solution of 2-chloroaniline (1.81 g, 14.29 mmol) were added into a round-bottom flask containing a solution of 4-chlorobutanoylchloride (2.00 g, 14.29 mmol) and ammonium thiocyanate (1.10 g, 14.29 mmol). The solution mixture was refluxed for 2.5 h then filtered off and left to evaporate at room temperature. The yellowish precipitate obtained was washed with water and cold ethanol. The yellowish crystals suitable for X-ray analysis were obtained by recrystallization of the precipitate in DMF.

S3. Refinement

All H atoms were positioned geometrically [C—H 0.93–0.97 Å; N—H 0.86 Å], and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$.

**Figure 1**

The content of asymmetric unit of the title compound showing the atomic numbering and 40% probability displacement ellipsoids. Dashed lines denote hydrogen bonds.




Figure 2

The molecular packing viewed down the *b* axis. Dashed lines denote N—H···S interactions.

1-(4-Chlorobutanoyl)-3-(2-chlorophenyl)thiourea

Crystal data

$C_{11}H_{12}Cl_2N_2OS$

$M_r = 291.19$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 14.396\ (3)\ \text{\AA}$

$b = 10.941\ (2)\ \text{\AA}$

$c = 18.093\ (4)\ \text{\AA}$

$\beta = 109.399\ (4)^\circ$

$V = 2688.0\ (9)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1200$

$D_x = 1.439\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 6026 reflections

$\theta = 1.5\text{--}25.0^\circ$

$\mu = 0.62\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, yellow

$0.42 \times 0.41 \times 0.39\ \text{mm}$

Data collection

Bruker SMART APEX CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 83.66 pixels mm⁻¹
 ω scan
Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)
 $T_{\min} = 0.780$, $T_{\max} = 0.793$

14457 measured reflections
4727 independent reflections
3802 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -17 \rightarrow 17$
 $k = -13 \rightarrow 12$
 $l = -16 \rightarrow 21$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.109$
 $S = 1.03$
4727 reflections
307 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0514P)^2 + 1.5167P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.33 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.02869 (4)	-0.00674 (6)	0.33632 (3)	0.04643 (17)
S2	0.23770 (4)	0.49698 (6)	0.02320 (4)	0.04616 (17)
Cl1	-0.08277 (6)	-0.17066 (6)	0.15579 (5)	0.0684 (2)
Cl2	0.52142 (7)	0.18550 (11)	0.47501 (6)	0.1086 (4)
Cl3	0.38361 (6)	0.18324 (8)	0.09743 (5)	0.0738 (2)
Cl4	-0.23405 (6)	0.18796 (9)	-0.20368 (5)	0.0792 (3)
O1	0.20202 (12)	0.12434 (18)	0.23359 (10)	0.0555 (5)
O2	0.05946 (12)	0.18451 (16)	0.07274 (10)	0.0543 (5)
N1	0.01342 (13)	0.07137 (17)	0.21195 (10)	0.0385 (4)
H1A	0.0562	0.0916	0.1904	0.046*
N2	0.14546 (13)	0.04033 (17)	0.32535 (11)	0.0400 (4)
H2A	0.1648	0.0114	0.3722	0.048*
N3	0.21846 (13)	0.32495 (18)	0.11973 (11)	0.0430 (5)
H3A	0.1852	0.2670	0.1311	0.052*
N4	0.08539 (13)	0.35124 (17)	0.00731 (11)	0.0382 (4)
H4A	0.0585	0.3957	-0.0334	0.046*

C1	-0.14018 (18)	-0.0300 (2)	0.13660 (14)	0.0450 (6)
C2	-0.2387 (2)	-0.0240 (3)	0.09106 (16)	0.0620 (8)
H2	-0.2740	-0.0952	0.0724	0.074*
C3	-0.2839 (2)	0.0878 (3)	0.07364 (18)	0.0734 (9)
H3	-0.3500	0.0921	0.0431	0.088*
C4	-0.2326 (2)	0.1928 (3)	0.10078 (19)	0.0683 (8)
H4	-0.2636	0.2682	0.0878	0.082*
C5	-0.13441 (19)	0.1870 (2)	0.14771 (16)	0.0540 (6)
H5	-0.0997	0.2584	0.1669	0.065*
C6	-0.08857 (16)	0.0750 (2)	0.16572 (13)	0.0394 (5)
C7	0.04504 (15)	0.03795 (19)	0.28697 (13)	0.0358 (5)
C8	0.21855 (16)	0.0824 (2)	0.29900 (14)	0.0437 (5)
C9	0.31985 (18)	0.0696 (3)	0.35824 (16)	0.0608 (7)
H9A	0.3219	0.1128	0.4056	0.073*
H9B	0.3322	-0.0161	0.3715	0.073*
C10	0.39933 (19)	0.1170 (3)	0.33112 (17)	0.0644 (8)
H10A	0.3885	0.2036	0.3200	0.077*
H10B	0.3955	0.0764	0.2825	0.077*
C11	0.5010 (2)	0.0994 (4)	0.38907 (18)	0.0743 (9)
H11A	0.5491	0.1226	0.3647	0.089*
H11B	0.5107	0.0135	0.4027	0.089*
C12	0.39626 (18)	0.2933 (2)	0.16893 (14)	0.0487 (6)
C13	0.48814 (19)	0.3193 (3)	0.22162 (17)	0.0610 (7)
H13	0.5437	0.2795	0.2183	0.073*
C14	0.4967 (2)	0.4046 (3)	0.27902 (18)	0.0690 (8)
H14	0.5584	0.4223	0.3149	0.083*
C15	0.4150 (2)	0.4644 (3)	0.28422 (18)	0.0700 (8)
H15	0.4217	0.5219	0.3235	0.084*
C16	0.32314 (19)	0.4390 (3)	0.23107 (16)	0.0562 (7)
H16	0.2679	0.4798	0.2343	0.067*
C17	0.31345 (16)	0.3533 (2)	0.17331 (13)	0.0435 (6)
C18	0.17954 (15)	0.3847 (2)	0.05298 (12)	0.0355 (5)
C19	0.02925 (16)	0.2559 (2)	0.01862 (13)	0.0398 (5)
C20	-0.07275 (16)	0.2507 (2)	-0.03992 (15)	0.0455 (6)
H20A	-0.0723	0.2884	-0.0883	0.055*
H20B	-0.1165	0.2979	-0.0202	0.055*
C21	-0.11259 (18)	0.1228 (2)	-0.05729 (15)	0.0527 (6)
H21A	-0.1075	0.0822	-0.0084	0.063*
H21B	-0.0726	0.0776	-0.0817	0.063*
C22	-0.2184 (2)	0.1201 (3)	-0.11039 (17)	0.0689 (8)
H22A	-0.2588	0.1636	-0.0854	0.083*
H22B	-0.2410	0.0360	-0.1180	0.083*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0392 (3)	0.0612 (4)	0.0382 (3)	-0.0085 (3)	0.0118 (3)	-0.0020 (3)
S2	0.0413 (3)	0.0525 (4)	0.0396 (3)	-0.0108 (3)	0.0067 (3)	0.0051 (3)

C11	0.0886 (5)	0.0397 (4)	0.0721 (5)	-0.0045 (3)	0.0203 (4)	-0.0071 (3)
C12	0.0732 (6)	0.1360 (9)	0.0916 (7)	-0.0178 (6)	-0.0060 (5)	-0.0378 (6)
C13	0.0698 (5)	0.0801 (5)	0.0665 (5)	0.0138 (4)	0.0159 (4)	-0.0065 (4)
C14	0.0651 (5)	0.0973 (6)	0.0554 (4)	-0.0125 (4)	-0.0066 (4)	0.0048 (4)
O1	0.0387 (9)	0.0765 (12)	0.0466 (10)	-0.0065 (9)	0.0076 (8)	0.0167 (9)
O2	0.0442 (9)	0.0552 (10)	0.0505 (10)	-0.0112 (8)	-0.0018 (8)	0.0148 (9)
N1	0.0307 (9)	0.0449 (11)	0.0362 (10)	-0.0047 (8)	0.0062 (8)	0.0032 (8)
N2	0.0336 (10)	0.0486 (11)	0.0332 (10)	-0.0001 (8)	0.0050 (8)	0.0053 (8)
N3	0.0321 (10)	0.0524 (12)	0.0382 (10)	-0.0086 (8)	0.0031 (8)	0.0081 (9)
N4	0.0333 (9)	0.0435 (10)	0.0326 (9)	-0.0016 (8)	0.0039 (8)	0.0036 (8)
C1	0.0467 (13)	0.0463 (13)	0.0381 (13)	-0.0081 (11)	0.0089 (11)	-0.0007 (10)
C2	0.0508 (15)	0.077 (2)	0.0480 (15)	-0.0241 (15)	0.0033 (13)	-0.0025 (14)
C3	0.0405 (15)	0.104 (3)	0.0628 (19)	-0.0010 (17)	-0.0006 (13)	0.0162 (18)
C4	0.0504 (16)	0.073 (2)	0.072 (2)	0.0178 (15)	0.0067 (15)	0.0119 (16)
C5	0.0520 (15)	0.0462 (14)	0.0574 (16)	0.0027 (12)	0.0095 (13)	0.0021 (12)
C6	0.0350 (11)	0.0460 (13)	0.0337 (11)	-0.0014 (10)	0.0066 (9)	0.0044 (10)
C7	0.0345 (11)	0.0316 (11)	0.0374 (12)	-0.0011 (9)	0.0067 (9)	-0.0026 (9)
C8	0.0360 (12)	0.0510 (14)	0.0399 (13)	-0.0020 (10)	0.0071 (10)	0.0040 (11)
C9	0.0411 (14)	0.087 (2)	0.0479 (15)	-0.0073 (14)	0.0059 (12)	0.0124 (14)
C10	0.0419 (14)	0.093 (2)	0.0562 (17)	0.0013 (14)	0.0129 (13)	0.0058 (16)
C11	0.0378 (14)	0.113 (3)	0.0657 (19)	0.0026 (16)	0.0092 (13)	-0.0008 (18)
C12	0.0416 (13)	0.0552 (15)	0.0441 (14)	-0.0034 (11)	0.0074 (11)	0.0109 (11)
C13	0.0343 (13)	0.0751 (19)	0.0650 (18)	0.0009 (13)	0.0049 (12)	0.0169 (16)
C14	0.0412 (15)	0.084 (2)	0.0635 (19)	-0.0182 (15)	-0.0068 (13)	0.0083 (17)
C15	0.0652 (19)	0.075 (2)	0.0554 (18)	-0.0198 (16)	0.0008 (15)	-0.0108 (15)
C16	0.0448 (14)	0.0633 (17)	0.0530 (16)	-0.0035 (12)	0.0065 (12)	-0.0016 (13)
C17	0.0339 (12)	0.0532 (14)	0.0364 (12)	-0.0063 (10)	0.0024 (10)	0.0096 (11)
C18	0.0302 (10)	0.0407 (12)	0.0327 (11)	0.0019 (9)	0.0066 (9)	-0.0030 (9)
C19	0.0346 (11)	0.0408 (12)	0.0403 (13)	-0.0014 (10)	0.0073 (10)	-0.0017 (10)
C20	0.0321 (11)	0.0481 (14)	0.0492 (14)	-0.0023 (10)	0.0041 (10)	0.0025 (11)
C21	0.0456 (14)	0.0565 (15)	0.0486 (15)	-0.0126 (12)	0.0058 (12)	0.0036 (12)
C22	0.0501 (16)	0.089 (2)	0.0587 (17)	-0.0276 (15)	0.0064 (13)	-0.0009 (16)

Geometric parameters (Å, °)

S1—C7	1.671 (2)	C5—H5	0.9300
S2—C18	1.673 (2)	C8—C9	1.502 (3)
C11—C1	1.727 (3)	C9—C10	1.481 (4)
C12—C11	1.758 (3)	C9—H9A	0.9700
C13—C12	1.732 (3)	C9—H9B	0.9700
C14—C22	1.788 (3)	C10—C11	1.504 (4)
O1—C8	1.217 (3)	C10—H10A	0.9700
O2—C19	1.215 (3)	C10—H10B	0.9700
N1—C7	1.332 (3)	C11—H11A	0.9700
N1—C6	1.429 (3)	C11—H11B	0.9700
N1—H1A	0.8600	C12—C13	1.379 (4)
N2—C8	1.371 (3)	C12—C17	1.386 (3)
N2—C7	1.381 (3)	C13—C14	1.371 (4)

N2—H2A	0.8600	C13—H13	0.9300
N3—C18	1.323 (3)	C14—C15	1.375 (5)
N3—C17	1.423 (3)	C14—H14	0.9300
N3—H3A	0.8600	C15—C16	1.381 (4)
N4—C19	1.376 (3)	C15—H15	0.9300
N4—C18	1.382 (3)	C16—C17	1.377 (4)
N4—H4A	0.8600	C16—H16	0.9300
C1—C6	1.374 (3)	C19—C20	1.499 (3)
C1—C2	1.385 (4)	C20—C21	1.505 (4)
C2—C3	1.371 (5)	C20—H20A	0.9700
C2—H2	0.9300	C20—H20B	0.9700
C3—C4	1.365 (5)	C21—C22	1.507 (4)
C3—H3	0.9300	C21—H21A	0.9700
C4—C5	1.388 (4)	C21—H21B	0.9700
C4—H4	0.9300	C22—H22A	0.9700
C5—C6	1.379 (3)	C22—H22B	0.9700
C7—N1—C6	122.78 (18)	C10—C11—C12	112.7 (2)
C7—N1—H1A	118.6	C10—C11—H11A	109.1
C6—N1—H1A	118.6	C12—C11—H11A	109.1
C8—N2—C7	128.89 (19)	C10—C11—H11B	109.1
C8—N2—H2A	115.6	C12—C11—H11B	109.1
C7—N2—H2A	115.6	H11A—C11—H11B	107.8
C18—N3—C17	122.27 (19)	C13—C12—C17	120.6 (3)
C18—N3—H3A	118.9	C13—C12—C13	119.9 (2)
C17—N3—H3A	118.9	C17—C12—C13	119.53 (19)
C19—N4—C18	128.30 (19)	C14—C13—C12	119.2 (3)
C19—N4—H4A	115.8	C14—C13—H13	120.4
C18—N4—H4A	115.8	C12—C13—H13	120.4
C6—C1—C2	120.3 (2)	C13—C14—C15	120.8 (3)
C6—C1—C11	120.41 (18)	C13—C14—H14	119.6
C2—C1—C11	119.3 (2)	C15—C14—H14	119.6
C3—C2—C1	119.5 (3)	C14—C15—C16	120.0 (3)
C3—C2—H2	120.3	C14—C15—H15	120.0
C1—C2—H2	120.3	C16—C15—H15	120.0
C4—C3—C2	120.6 (3)	C17—C16—C15	119.9 (3)
C4—C3—H3	119.7	C17—C16—H16	120.1
C2—C3—H3	119.7	C15—C16—H16	120.1
C3—C4—C5	120.1 (3)	C16—C17—C12	119.6 (2)
C3—C4—H4	120.0	C16—C17—N3	119.9 (2)
C5—C4—H4	120.0	C12—C17—N3	120.5 (2)
C6—C5—C4	119.7 (3)	N3—C18—N4	116.67 (19)
C6—C5—H5	120.2	N3—C18—S2	123.45 (16)
C4—C5—H5	120.2	N4—C18—S2	119.88 (16)
C1—C6—C5	119.8 (2)	O2—C19—N4	122.5 (2)
C1—C6—N1	121.4 (2)	O2—C19—C20	123.3 (2)
C5—C6—N1	118.8 (2)	N4—C19—C20	114.20 (19)
N1—C7—N2	116.99 (19)	C19—C20—C21	113.5 (2)

N1—C7—S1	124.22 (16)	C19—C20—H20A	108.9
N2—C7—S1	118.80 (16)	C21—C20—H20A	108.9
O1—C8—N2	122.7 (2)	C19—C20—H20B	108.9
O1—C8—C9	124.0 (2)	C21—C20—H20B	108.9
N2—C8—C9	113.3 (2)	H20A—C20—H20B	107.7
C10—C9—C8	113.9 (2)	C20—C21—C22	112.7 (2)
C10—C9—H9A	108.8	C20—C21—H21A	109.1
C8—C9—H9A	108.8	C22—C21—H21A	109.1
C10—C9—H9B	108.8	C20—C21—H21B	109.1
C8—C9—H9B	108.8	C22—C21—H21B	109.1
H9A—C9—H9B	107.7	H21A—C21—H21B	107.8
C9—C10—C11	113.9 (2)	C21—C22—C14	112.16 (19)
C9—C10—H10A	108.8	C21—C22—H22A	109.2
C11—C10—H10A	108.8	C14—C22—H22A	109.2
C9—C10—H10B	108.8	C21—C22—H22B	109.2
C11—C10—H10B	108.8	C14—C22—H22B	109.2
H10A—C10—H10B	107.7	H22A—C22—H22B	107.9
C6—C1—C2—C3	1.5 (4)	C17—C12—C13—C14	0.6 (4)
C11—C1—C2—C3	-178.0 (2)	C13—C12—C13—C14	-178.5 (2)
C1—C2—C3—C4	0.0 (5)	C12—C13—C14—C15	-0.3 (4)
C2—C3—C4—C5	-1.2 (5)	C13—C14—C15—C16	-0.2 (5)
C3—C4—C5—C6	1.0 (5)	C14—C15—C16—C17	0.5 (5)
C2—C1—C6—C5	-1.8 (4)	C15—C16—C17—C12	-0.2 (4)
C11—C1—C6—C5	177.8 (2)	C15—C16—C17—N3	178.4 (2)
C2—C1—C6—N1	-179.2 (2)	C13—C12—C17—C16	-0.4 (4)
C11—C1—C6—N1	0.4 (3)	C13—C12—C17—C16	178.7 (2)
C4—C5—C6—C1	0.5 (4)	C13—C12—C17—N3	-179.0 (2)
C4—C5—C6—N1	178.0 (2)	C13—C12—C17—N3	0.1 (3)
C7—N1—C6—C1	-76.9 (3)	C18—N3—C17—C16	89.7 (3)
C7—N1—C6—C5	105.7 (3)	C18—N3—C17—C12	-91.7 (3)
C6—N1—C7—N2	-178.95 (19)	C17—N3—C18—N4	-177.8 (2)
C6—N1—C7—S1	1.4 (3)	C17—N3—C18—S2	1.4 (3)
C8—N2—C7—N1	6.1 (3)	C19—N4—C18—N3	-6.3 (3)
C8—N2—C7—S1	-174.26 (19)	C19—N4—C18—S2	174.38 (18)
C7—N2—C8—O1	-0.9 (4)	C18—N4—C19—O2	-1.8 (4)
C7—N2—C8—C9	180.0 (2)	C18—N4—C19—C20	176.4 (2)
O1—C8—C9—C10	2.1 (4)	O2—C19—C20—C21	-33.5 (3)
N2—C8—C9—C10	-178.8 (3)	N4—C19—C20—C21	148.3 (2)
C8—C9—C10—C11	-177.6 (3)	C19—C20—C21—C22	174.7 (2)
C9—C10—C11—C12	-66.2 (4)	C20—C21—C22—C14	61.0 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1A...O1	0.86	2.01	2.675 (3)	133
N3—H3A...O2	0.86	1.98	2.652 (3)	134
N1—H1A...O2	0.86	2.37	3.070 (3)	138

N3—H3A···O1	0.86	2.37	3.073 (3)	139
N2—H2A···S2 ⁱ	0.86	2.58	3.404 (2)	160
N4—H4A···S1 ⁱⁱ	0.86	2.58	3.433 (2)	174

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $x, -y+1/2, z-1/2$.