

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-N'-[3-(4-Chlorobenzoyloxy)benzylidene]pyridine-4-carbohydrazide acetic acid monosolvate monohydrate

Chun-Hua Diao and Zhi Fan*

College of Sciences, Tianjin University of Science and Technology, Tianjin 300457, People's Republic of China

Correspondence e-mail: zhifan@tust.edu.cn


Received 10 April 2012; accepted 19 April 2012

 Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.043; wR factor = 0.119; data-to-parameter ratio = 13.2.

In the Schiff base molecule of the title compound, $\text{C}_{20}\text{H}_{14}\text{ClN}_3\text{O}_3 \cdot \text{CH}_3\text{COOH} \cdot \text{H}_2\text{O}$, the central benzene ring makes dihedral angles of 36.26 (7) and 27.59 (8)°, respectively, with the terminal chlorophenyl and pyridine rings. In the crystal, the three components are linked by $\text{O}-\text{H} \cdots \text{O}$, $\text{N}-\text{H} \cdots \text{O}$, $\text{O}-\text{H} \cdots \text{N}$ and $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds into a double-tape structure along the a axis.

Related literature

For general background to the use of Schiff base derivatives in the development of protein and enzyme mimics, see: Santos *et al.* (2001). For closely related crystal structures, see: Diao *et al.* (2007); Peralta *et al.* (2007); de Souza *et al.* (2007); Wardell *et al.* (2005). For reference bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{14}\text{ClN}_3\text{O}_3 \cdot \text{C}_2\text{H}_4\text{O}_2 \cdot \text{H}_2\text{O}$
 $M_r = 457.86$
 Triclinic, $P\bar{1}$
 $a = 6.6666$ (15) Å
 $b = 7.5437$ (17) Å

$c = 24.781$ (6) Å
 $\alpha = 81.526$ (4)°
 $\beta = 82.969$ (4)°
 $\gamma = 66.632$ (4)°
 $V = 1128.7$ (5) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.21$ mm⁻¹

$T = 294$ K
 $0.18 \times 0.16 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.928$, $T_{\text{max}} = 0.979$

5755 measured reflections
 3936 independent reflections
 2501 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.119$
 $S = 1.03$
 3936 reflections
 298 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O6}-\text{H6A} \cdots \text{O3}$	0.86 (3)	2.00 (3)	2.843 (2)	166 (2)
$\text{O6}-\text{H6B} \cdots \text{O4}^i$	0.86 (2)	1.98 (2)	2.812 (3)	165 (3)
$\text{O5}-\text{H5A} \cdots \text{N3}^{ii}$	0.82	1.85	2.646 (3)	164
$\text{N2}-\text{H2} \cdots \text{O6}^{iii}$	0.86	2.04	2.879 (2)	164
$\text{C14}-\text{H14} \cdots \text{O6}^{iii}$	0.93	2.59	3.367 (4)	141
$\text{C17}-\text{H17} \cdots \text{O6}^{iii}$	0.93	2.50	3.331 (3)	149

Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $x, y, z - 1$; (iii) $x + 1, y, z$.

Data collection: SMART (Bruker, 1999); cell refinement: SAINT (Bruker, 1999); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5118).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (1999). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Diao, C.-H., Fan, Z. & Chen, X. (2007). *Acta Cryst.* **C63**, o717–o719.
- Peralta, M. A., de Souza, M. N. V., Wardell, S. M. S. V., Wardell, J. L., Low, J. N. & Glidewell, C. (2007). *Acta Cryst.* **C63**, o68–o72.
- Santos, M. L. P., Bagatin, I. A., Pereira, E. M. & Ferreira, A. M. D. C. (2001). *J. Chem. Soc. Dalton Trans.* pp. 838–844.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Souza, M. V. N. de, Wardell, S. M. S. V., Wardell, J. L., Low, J. N. & Glidewell, C. (2007). *Acta Cryst.* **C63**, o166–o168.
- Wardell, S. M. S. V., de Souza, M. V. N., Wardell, J. L., Low, J. N. & Glidewell, C. (2005). *Acta Cryst.* **C61**, o683–o689.

supporting information

Acta Cryst. (2012). E68, o1517 [doi:10.1107/S1600536812017369]

(*E*)-*N'*-[3-(4-Chlorobenzoyloxy)benzylidene]pyridine-4-carbohydrazide acetic acid monosolvate monohydrate

Chun-Hua Diao and Zhi Fan

S1. Comment

Many Schiff base derivatives have been synthesized and found to exhibit important pharmacological properties, such as antibacterial, antitumor and antitoxic activities (Santos *et al.*, 2001). Among the large number of the compounds, isonicotinohydrazide forms a variety of Schiff bases with aldehydes, and the synthesis and crystal structures of some of them have been reported (Wardell *et al.*, 2005; Peralta *et al.*, 2007; de Souza *et al.*, 2007).

In order to obtain more detailed information on the structural conformation of the molecule that may be of value in structure-activity analysis, we report here the synthesis and structure of the title compound, (I), as part of our study of isonicotinoylhydrazones (Diao *et al.*, 2007).

In (I) (Fig. 1), the central benzene ring (C8–C14/O2) is nearly planar, with an r.m.s. deviation for fitted atoms of 0.0255 Å. This plane makes dihedral angles of 27.59 (8) and 36.26 (7)° with the pyridine ring (C16–C20/N3) and the terminal benzene ring (C1–C6), respectively. The dihedral angle between the pyridine ring and the benzene ring is 26.03 (7)°. All bond lengths are within normal ranges (Allen *et al.*, 1987).

The acetic acid molecule and the water molecule are effectively tethered to the Schiff base component by a combination of independent hydrogen bonds, two O—H···O, one N—H···O, one O—H···N and one C—H···O types (Fig. 2 and Table 2).

S2. Experimental

An anhydrous ethanol solution (50 ml) of 3-formylphenyl 4-chlorobenzoate (2.61 g, 10 mmol) was added to an anhydrous ethanol solution (50 ml) of isonicotinohydrazide (1.37 g, 10 mmol) and the mixture stirred at 350 K for 5 h under N₂, giving a white precipitate. The product was isolated, recrystallized from ethanol, and then dried in a vacuum to give pure compound (I) in 72% yield. Colorless single crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of a solution of ethanol and acetic acid (80:20 v/v).

S3. Refinement

The H atoms of the water molecules were located in a difference Fourier map and refined, with distance restraints of O—H = 0.85 (1) and H···H = 1.45 (1) Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. Other H atoms were included in calculated positions and refined using a riding model approximation. Constrained C—H and N—H bond lengths and $U_{\text{iso}}(\text{H})$ values: 0.93 Å and $1.2U_{\text{eq}}(\text{C})$ for Csp^2 —H; 0.97 Å and $1.2U_{\text{eq}}(\text{C})$ for methylene C—H; 0.82 Å and $1.5U_{\text{eq}}(\text{O})$ for hydroxyl O—H; 0.86 Å and $1.2U_{\text{eq}}(\text{N})$ for imino N—H.




Figure 1

The structure of the title compound with displacement ellipsoids for non-H atoms drawn at the 50% probability level.




Figure 2

A packing diagram for the title compound with the hydrogen bonds drawn as dashed lines. H atoms not involved in the hydrogen bonds have been omitted.

(E)-N'-[3-(4-Chlorobenzoyloxy)benzylidene]pyridine-4- carbohydrazide acetic acid monosolvate monohydrate

Crystal data

$C_{20}H_{14}ClN_3O_3 \cdot C_2H_4O_2 \cdot H_2O$

$M_r = 457.86$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.6666$ (15) Å

$b = 7.5437$ (17) Å

$c = 24.781$ (6) Å

$\alpha = 81.526$ (4)°

$\beta = 82.969$ (4)°

$\gamma = 66.632$ (4)°

$V = 1128.7$ (5) Å³

$Z = 2$

$F(000) = 476$

$D_x = 1.347$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 1984 reflections
 $\theta = 2.5\text{--}25.2^\circ$
 $\mu = 0.21 \text{ mm}^{-1}$

$T = 294 \text{ K}$
 Block, colorless
 $0.18 \times 0.16 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.928$, $T_{\max} = 0.979$

5755 measured reflections
 3936 independent reflections
 2501 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -6 \rightarrow 7$
 $k = -8 \rightarrow 8$
 $l = -27 \rightarrow 29$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.119$
 $S = 1.03$
 3936 reflections
 298 parameters
 3 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0473P)^2 + 0.2041P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.15 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0070 (17)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.33537 (13)	0.76087 (12)	0.41389 (3)	0.0905 (3)
O1	0.5002 (3)	0.8179 (3)	0.59922 (8)	0.0951 (7)
O2	0.7889 (3)	0.6503 (3)	0.64925 (7)	0.0686 (5)
O3	0.3270 (2)	0.7186 (2)	0.98573 (7)	0.0608 (5)
N1	0.5560 (3)	0.6798 (3)	0.88899 (8)	0.0524 (5)
N2	0.6335 (3)	0.7116 (3)	0.93397 (7)	0.0534 (5)
H2	0.7581	0.7221	0.9319	0.064*
N3	0.7533 (4)	0.8006 (3)	1.12485 (8)	0.0626 (6)
C1	0.7913 (5)	0.7979 (5)	0.50434 (12)	0.0820 (9)
H1	0.6443	0.8341	0.4987	0.098*
C2	0.9364 (5)	0.8039 (4)	0.46046 (11)	0.0809 (9)

H2A	0.8882	0.8445	0.4254	0.097*
C3	1.1507 (4)	0.7501 (4)	0.46886 (10)	0.0628 (7)
C4	1.2231 (5)	0.6916 (5)	0.52037 (11)	0.0822 (9)
H4	1.3702	0.6559	0.5258	0.099*
C5	1.0767 (4)	0.6859 (4)	0.56408 (11)	0.0742 (8)
H5	1.1258	0.6454	0.5991	0.089*
C6	0.8596 (4)	0.7392 (3)	0.55674 (10)	0.0563 (6)
C7	0.6938 (4)	0.7422 (4)	0.60230 (11)	0.0643 (7)
C8	0.6574 (4)	0.6336 (4)	0.69716 (10)	0.0573 (6)
C9	0.4946 (4)	0.5648 (4)	0.69816 (11)	0.0669 (7)
H9	0.4609	0.5348	0.6662	0.080*
C10	0.3821 (4)	0.5412 (4)	0.74740 (11)	0.0659 (7)
H10	0.2705	0.4958	0.7486	0.079*
C11	0.4328 (4)	0.5838 (3)	0.79477 (10)	0.0580 (6)
H11	0.3540	0.5691	0.8276	0.070*
C12	0.6014 (4)	0.6488 (3)	0.79379 (9)	0.0511 (6)
C13	0.7122 (4)	0.6747 (4)	0.74412 (10)	0.0573 (6)
H13	0.8239	0.7201	0.7426	0.069*
C14	0.6656 (4)	0.6871 (4)	0.84357 (10)	0.0581 (6)
H14	0.7887	0.7173	0.8421	0.070*
C15	0.5077 (4)	0.7258 (3)	0.98161 (9)	0.0474 (6)
C16	0.6030 (3)	0.7529 (3)	1.03005 (9)	0.0450 (5)
C17	0.8176 (4)	0.7278 (4)	1.03257 (10)	0.0626 (7)
H17	0.9165	0.6951	1.0022	0.075*
C18	0.8846 (4)	0.7516 (4)	1.08031 (11)	0.0713 (8)
H18	1.0309	0.7321	1.0814	0.086*
C19	0.5476 (4)	0.8251 (4)	1.12229 (10)	0.0636 (7)
H19	0.4521	0.8590	1.1532	0.076*
C20	0.4667 (4)	0.8032 (4)	1.07641 (10)	0.0594 (7)
H20	0.3199	0.8225	1.0767	0.071*
O4	1.0439 (3)	0.9861 (3)	0.19590 (9)	0.0936 (7)
O5	0.8367 (4)	0.8213 (3)	0.22488 (7)	0.0878 (6)
H5A	0.8297	0.8234	0.1920	0.132*
C21	0.9595 (5)	0.9109 (4)	0.23256 (11)	0.0660 (7)
C22	0.9809 (6)	0.9134 (5)	0.29172 (12)	0.1010 (11)
H22A	1.1128	0.9314	0.2958	0.151*
H22B	0.9867	0.7924	0.3116	0.151*
H22C	0.8570	1.0181	0.3058	0.151*
O6	0.0290 (3)	0.7824 (3)	0.90528 (8)	0.0777 (6)
H6A	0.134 (4)	0.765 (4)	0.9249 (10)	0.117*
H6B	0.018 (5)	0.864 (4)	0.8768 (8)	0.117*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0937 (6)	0.1030 (6)	0.0582 (5)	-0.0225 (5)	0.0095 (4)	-0.0146 (4)
O1	0.0563 (12)	0.1292 (18)	0.0847 (15)	-0.0278 (12)	-0.0173 (10)	0.0215 (12)
O2	0.0556 (10)	0.0987 (13)	0.0469 (10)	-0.0248 (10)	-0.0071 (8)	-0.0053 (9)

O3	0.0446 (9)	0.0839 (12)	0.0618 (11)	-0.0325 (9)	-0.0085 (8)	-0.0052 (9)
N1	0.0542 (12)	0.0602 (12)	0.0502 (12)	-0.0289 (10)	-0.0117 (10)	-0.0023 (9)
N2	0.0512 (11)	0.0751 (13)	0.0467 (12)	-0.0376 (10)	-0.0089 (9)	-0.0021 (9)
N3	0.0690 (14)	0.0746 (14)	0.0544 (13)	-0.0382 (12)	-0.0109 (11)	-0.0031 (10)
C1	0.0660 (18)	0.117 (3)	0.0624 (19)	-0.0361 (17)	-0.0226 (15)	0.0082 (16)
C2	0.083 (2)	0.104 (2)	0.0537 (18)	-0.0337 (18)	-0.0217 (16)	0.0064 (15)
C3	0.0722 (17)	0.0602 (15)	0.0519 (15)	-0.0195 (14)	-0.0023 (13)	-0.0129 (12)
C4	0.0588 (17)	0.123 (3)	0.0561 (18)	-0.0247 (17)	-0.0066 (14)	-0.0115 (16)
C5	0.0655 (17)	0.107 (2)	0.0449 (15)	-0.0256 (16)	-0.0122 (13)	-0.0066 (14)
C6	0.0581 (15)	0.0588 (15)	0.0527 (15)	-0.0211 (12)	-0.0122 (12)	-0.0056 (11)
C7	0.0613 (17)	0.0715 (17)	0.0618 (17)	-0.0257 (14)	-0.0178 (14)	-0.0013 (13)
C8	0.0514 (14)	0.0650 (16)	0.0524 (15)	-0.0184 (13)	-0.0058 (12)	-0.0071 (12)
C9	0.0721 (17)	0.0750 (18)	0.0613 (17)	-0.0312 (15)	-0.0091 (14)	-0.0189 (13)
C10	0.0687 (17)	0.0780 (18)	0.0680 (18)	-0.0430 (15)	-0.0068 (14)	-0.0153 (14)
C11	0.0581 (15)	0.0636 (16)	0.0588 (16)	-0.0295 (13)	-0.0036 (12)	-0.0091 (12)
C12	0.0498 (13)	0.0557 (14)	0.0495 (14)	-0.0215 (11)	-0.0086 (11)	-0.0039 (11)
C13	0.0500 (14)	0.0720 (17)	0.0562 (16)	-0.0297 (13)	-0.0100 (12)	-0.0034 (12)
C14	0.0566 (15)	0.0740 (17)	0.0550 (16)	-0.0373 (13)	-0.0093 (12)	-0.0025 (12)
C15	0.0464 (13)	0.0474 (13)	0.0507 (14)	-0.0217 (11)	-0.0097 (11)	0.0034 (10)
C16	0.0433 (12)	0.0474 (13)	0.0475 (13)	-0.0231 (10)	-0.0060 (10)	0.0033 (10)
C17	0.0501 (14)	0.0955 (19)	0.0517 (15)	-0.0379 (14)	-0.0007 (11)	-0.0106 (13)
C18	0.0582 (16)	0.108 (2)	0.0625 (18)	-0.0458 (16)	-0.0087 (14)	-0.0108 (16)
C19	0.0685 (17)	0.0755 (18)	0.0487 (15)	-0.0307 (15)	0.0001 (12)	-0.0076 (12)
C20	0.0481 (14)	0.0770 (17)	0.0574 (16)	-0.0305 (13)	-0.0008 (12)	-0.0045 (13)
O4	0.0878 (14)	0.1262 (18)	0.0788 (14)	-0.0620 (14)	-0.0170 (11)	0.0218 (12)
O5	0.1197 (17)	0.1169 (16)	0.0548 (12)	-0.0778 (15)	-0.0107 (12)	0.0026 (12)
C21	0.0703 (18)	0.0653 (17)	0.0599 (17)	-0.0235 (15)	-0.0166 (14)	0.0033 (13)
C22	0.146 (3)	0.096 (2)	0.068 (2)	-0.049 (2)	-0.040 (2)	0.0006 (17)
O6	0.0587 (11)	0.1268 (17)	0.0607 (12)	-0.0549 (12)	-0.0152 (9)	0.0147 (11)

Geometric parameters (Å, °)

C11—C3	1.736 (3)	C10—C11	1.375 (3)
O1—C7	1.194 (3)	C10—H10	0.9300
O2—C7	1.353 (3)	C11—C12	1.389 (3)
O2—C8	1.408 (3)	C11—H11	0.9300
O3—C15	1.218 (2)	C12—C13	1.385 (3)
N1—C14	1.271 (3)	C12—C14	1.456 (3)
N1—N2	1.373 (2)	C13—H13	0.9300
N2—C15	1.353 (3)	C14—H14	0.9300
N2—H2	0.8600	C15—C16	1.499 (3)
N3—C19	1.317 (3)	C16—C20	1.374 (3)
N3—C18	1.322 (3)	C16—C17	1.375 (3)
C1—C2	1.372 (4)	C17—C18	1.374 (3)
C1—C6	1.380 (3)	C17—H17	0.9300
C1—H1	0.9300	C18—H18	0.9300
C2—C3	1.356 (4)	C19—C20	1.373 (3)
C2—H2A	0.9300	C19—H19	0.9300

C3—C4	1.368 (3)	C20—H20	0.9300
C4—C5	1.374 (4)	O4—C21	1.200 (3)
C4—H4	0.9300	O5—C21	1.295 (3)
C5—C6	1.368 (3)	O5—H5A	0.8200
C5—H5	0.9300	C21—C22	1.494 (4)
C6—C7	1.476 (4)	C22—H22A	0.9600
C8—C9	1.372 (3)	C22—H22B	0.9600
C8—C13	1.373 (3)	C22—H22C	0.9600
C9—C10	1.379 (3)	O6—H6A	0.854 (10)
C9—H9	0.9300	O6—H6B	0.852 (10)
C7—O2—C8	119.8 (2)	C13—C12—C11	118.7 (2)
C14—N1—N2	116.74 (19)	C13—C12—C14	119.7 (2)
C15—N2—N1	117.92 (19)	C11—C12—C14	121.6 (2)
C15—N2—H2	121.0	C8—C13—C12	120.2 (2)
N1—N2—H2	121.0	C8—C13—H13	119.9
C19—N3—C18	116.7 (2)	C12—C13—H13	119.9
C2—C1—C6	121.2 (3)	N1—C14—C12	121.0 (2)
C2—C1—H1	119.4	N1—C14—H14	119.5
C6—C1—H1	119.4	C12—C14—H14	119.5
C3—C2—C1	119.2 (3)	O3—C15—N2	123.0 (2)
C3—C2—H2A	120.4	O3—C15—C16	121.0 (2)
C1—C2—H2A	120.4	N2—C15—C16	116.0 (2)
C2—C3—C4	120.9 (3)	C20—C16—C17	116.8 (2)
C2—C3—C11	119.7 (2)	C20—C16—C15	117.7 (2)
C4—C3—C11	119.4 (2)	C17—C16—C15	125.5 (2)
C3—C4—C5	119.5 (3)	C18—C17—C16	119.4 (2)
C3—C4—H4	120.2	C18—C17—H17	120.3
C5—C4—H4	120.2	C16—C17—H17	120.3
C6—C5—C4	120.8 (2)	N3—C18—C17	123.8 (2)
C6—C5—H5	119.6	N3—C18—H18	118.1
C4—C5—H5	119.6	C17—C18—H18	118.1
C5—C6—C1	118.4 (3)	N3—C19—C20	123.4 (2)
C5—C6—C7	123.2 (2)	N3—C19—H19	118.3
C1—C6—C7	118.4 (2)	C20—C19—H19	118.3
O1—C7—O2	123.4 (3)	C19—C20—C16	119.9 (2)
O1—C7—C6	125.3 (2)	C19—C20—H20	120.0
O2—C7—C6	111.3 (2)	C16—C20—H20	120.0
C9—C8—C13	121.3 (2)	C21—O5—H5A	109.5
C9—C8—O2	122.2 (2)	O4—C21—O5	123.3 (3)
C13—C8—O2	116.3 (2)	O4—C21—C22	123.9 (3)
C8—C9—C10	118.7 (2)	O5—C21—C22	112.8 (3)
C8—C9—H9	120.6	C21—C22—H22A	109.5
C10—C9—H9	120.6	C21—C22—H22B	109.5
C11—C10—C9	120.8 (2)	H22A—C22—H22B	109.5
C11—C10—H10	119.6	C21—C22—H22C	109.5
C9—C10—H10	119.6	H22A—C22—H22C	109.5
C10—C11—C12	120.3 (2)	H22B—C22—H22C	109.5

C10—C11—H11	119.9	H6A—O6—H6B	115.3 (17)
C12—C11—H11	119.9		
C14—N1—N2—C15	172.9 (2)	C10—C11—C12—C13	1.8 (3)
C6—C1—C2—C3	0.4 (5)	C10—C11—C12—C14	-177.3 (2)
C1—C2—C3—C4	-0.5 (5)	C9—C8—C13—C12	-0.6 (4)
C1—C2—C3—C11	-178.9 (2)	O2—C8—C13—C12	-175.4 (2)
C2—C3—C4—C5	0.5 (5)	C11—C12—C13—C8	-1.0 (3)
C11—C3—C4—C5	178.9 (2)	C14—C12—C13—C8	178.0 (2)
C3—C4—C5—C6	-0.4 (5)	N2—N1—C14—C12	177.86 (19)
C4—C5—C6—C1	0.3 (4)	C13—C12—C14—N1	174.2 (2)
C4—C5—C6—C7	-177.9 (3)	C11—C12—C14—N1	-6.7 (4)
C2—C1—C6—C5	-0.3 (5)	N1—N2—C15—O3	-2.7 (3)
C2—C1—C6—C7	178.1 (3)	N1—N2—C15—C16	177.73 (18)
C8—O2—C7—O1	2.7 (4)	O3—C15—C16—C20	-11.2 (3)
C8—O2—C7—C6	-178.4 (2)	N2—C15—C16—C20	168.4 (2)
C5—C6—C7—O1	166.5 (3)	O3—C15—C16—C17	167.4 (2)
C1—C6—C7—O1	-11.8 (4)	N2—C15—C16—C17	-13.0 (3)
C5—C6—C7—O2	-12.4 (4)	C20—C16—C17—C18	0.7 (4)
C1—C6—C7—O2	169.4 (2)	C15—C16—C17—C18	-178.0 (2)
C7—O2—C8—C9	50.5 (3)	C19—N3—C18—C17	0.7 (4)
C7—O2—C8—C13	-134.6 (2)	C16—C17—C18—N3	-1.0 (4)
C13—C8—C9—C10	1.4 (4)	C18—N3—C19—C20	-0.3 (4)
O2—C8—C9—C10	176.0 (2)	N3—C19—C20—C16	0.1 (4)
C8—C9—C10—C11	-0.6 (4)	C17—C16—C20—C19	-0.3 (3)
C9—C10—C11—C12	-1.0 (4)	C15—C16—C20—C19	178.5 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O6—H6A \cdots O3	0.86 (3)	2.00 (3)	2.843 (2)	166 (2)
O6—H6B \cdots O4 ⁱ	0.86 (2)	1.98 (2)	2.812 (3)	165 (3)
O5—H5A \cdots N3 ⁱⁱ	0.82	1.85	2.646 (3)	164
N2—H2 \cdots O6 ⁱⁱⁱ	0.86	2.04	2.879 (2)	164
C14—H14 \cdots O6 ⁱⁱⁱ	0.93	2.59	3.367 (4)	141
C17—H17 \cdots O6 ⁱⁱⁱ	0.93	2.50	3.331 (3)	149

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x, y, z-1$; (iii) $x+1, y, z$.