

2-Amino-3-carboxypyrazin-1-i um perchlorate bis(2-aminopyrazin-1-i um-3-carboxylate) monohydrate

Fadila Berrah,^{a,*} Sofiane Bouacida,^b Ahlem Bouhraoua^a and Thierry Roisnel^c

^aLaboratoire de Chimie Appliquée et Technologie des Matériaux (LCATM), Université d'Oum El Bouaghi 04000, Algeria, ^bUnité de Recherche de Chimie de l'Environnement et Moléculaire Structurale (CHEMS), Faculté des Sciences Exactes, Université Mentouri Constantine 25000, Algeria, and ^cCentre de Diffraction X, UMR 6226 CNRS Unité Sciences Chimiques de Rennes, Université de Rennes I, 263 Avenue du Général Leclerc, 35042 Rennes, France
Correspondence e-mail: fadilaber@yahoo.fr


Received 30 April 2012; accepted 9 May 2012

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.038; wR factor = 0.100; data-to-parameter ratio = 14.2.

The asymmetric unit of the title compound, $\text{C}_5\text{H}_6\text{N}_3\text{O}_2^+\cdot\text{ClO}_4^-\cdot 2\text{C}_5\text{H}_5\text{N}_3\text{O}_2\cdot\text{H}_2\text{O}$, comprises two symmetry-independent zwitterions, one cation, one perchlorate anion and one water molecule. In the crystal, the three different types of organic entities are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds, forming undulating sheets parallel to $(\bar{1}\bar{1}0)$. These sheets are in turn connected by $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds involving perchlorate anions and water molecules, forming a three-dimensional network. Intramolecular $\text{N}-\text{H}\cdots\text{O}$ and weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds are also present.

Related literature

For crystal structures of hybrid compounds obtained from 3-amino-pyrazine 2-carboxylic acid, see: Berrah *et al.* (2011*a,b,c*). For related perchlorate compounds, see: Bendjedou *et al.* (2003); Berrah *et al.* (2012); Toumi Akriche *et al.* (2010).



Experimental

Crystal data

$\text{C}_5\text{H}_6\text{N}_3\text{O}_2^+\cdot\text{ClO}_4^-\cdot 2\text{C}_5\text{H}_5\text{N}_3\text{O}_2\cdot\text{H}_2\text{O}$	$\beta = 108.148(8)^\circ$
$M_r = 535.83$	$\gamma = 102.416(8)^\circ$
Triclinic, $P\bar{1}$	$V = 1039.8(3)\text{ \AA}^3$
$a = 8.1332(14)\text{ \AA}$	$Z = 2$
$b = 11.816(2)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 11.850(2)\text{ \AA}$	$\mu = 0.27\text{ mm}^{-1}$
$\alpha = 95.696(9)^\circ$	$T = 150\text{ K}$
	$0.46 \times 0.27 \times 0.17\text{ mm}$

Data collection

Bruker APEXII diffractometer	15575 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2002)	4705 independent reflections
$T_{\min} = 0.855$, $T_{\max} = 0.955$	4165 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.1$	$\Delta\rho_{\text{max}} = 0.39\text{ e \AA}^{-3}$
$S = 1.04$	$\Delta\rho_{\text{min}} = -0.48\text{ e \AA}^{-3}$
4705 reflections	
332 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1A-H1A \cdots O1W	0.82	1.71	2.5258 (18)	172
O1W-H1W \cdots O1B ⁱ	0.88 (2)	1.92 (2)	2.7873 (19)	169 (2)
O1W-H2W \cdots O2	0.84 (2)	1.99 (2)	2.8176 (19)	172 (2)
O1W-H1W \cdots N3B ⁱ	0.88 (2)	2.56 (2)	3.052 (2)	115.8 (17)
N2B-H2B \cdots O1C	0.86	1.92	2.6935 (18)	149
N1C-H11C \cdots O2B ⁱⁱ	0.86	2.11	2.958 (2)	170
N2C-H2C \cdots O1B ⁱⁱ	0.86	1.76	2.6156 (19)	171
N2A-H2A \cdots O2C ⁱⁱⁱ	0.86	1.80	2.6536 (17)	175
N1A-H11A \cdots O1C ⁱⁱⁱ	0.86	2.14	2.9340 (19)	153
N1B-H11B \cdots O1A	0.86	2.26	2.916 (2)	133
N1B-H11B \cdots O1C	0.86	2.44	3.087 (2)	133
N1B-H12B \cdots O2B	0.86	2.21	2.814 (2)	127
N1A-H12A \cdots O2A	0.86	2.09	2.7038 (19)	128
N1C-H12C \cdots O2C	0.86	2.06	2.6734 (19)	128
N2B-H2B \cdots N3C	0.86	2.41	3.058 (2)	132
N1B-H12B \cdots N3A	0.86	2.41	3.035 (2)	130
N1A-H12A \cdots N3B ^{iv}	0.86	2.44	3.152 (2)	140
C4B-H4B \cdots O1 ^v	0.93	2.41	3.267 (2)	153
C4C-H4C \cdots O3 ^{vi}	0.93	2.45	3.350 (2)	162
CSA-H5A \cdots O2B	0.93	2.58	3.336 (2)	138
CSB-H5B \cdots O2 ^v	0.93	2.48	3.145 (2)	129
CSB-H5B \cdots O2A ^v	0.93	2.48	3.164 (2)	130

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x + 1, y + 1, z$; (iii) $-x, -y, -z$; (iv) $x, y, z - 1$; (v) $x, y, z + 1$; (vi) $-x + 2, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.* 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to the LCATM laboratory, Université Larbi Ben M'Hidi, Oum El Bouaghi, Algeria, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5470).

References

- Bendjeddou, L., Cherouana, A., Berrah, F. & Benali-Cherif, N. (2003). *Acta Cryst. E* **59**, o574–o576.
- Berrah, F., Bouacida, S., Anana, H. & Roisnel, T. (2012). *Acta Cryst. E* **68**, o1601–o1602.
- Berrah, F., Bouacida, S. & Roisnel, T. (2011c). *Acta Cryst. E* **67**, o1409–o1410.
- Berrah, F., Ouakkaf, A., Bouacida, S. & Roisnel, T. (2011a). *Acta Cryst. E* **67**, o525–o526.
- Berrah, F., Ouakkaf, A., Bouacida, S. & Roisnel, T. (2011b). *Acta Cryst. E* **67**, o677–o678.
- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Sheldrick, G. M. (2002). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Toumi Akriche, S., Rzaigui, M., Al-Hokbany, N. & Mahfouz, R. M. (2010). *Acta Cryst. E* **66**, o300.

supporting information

Acta Cryst. (2012). E68, o1714–o1715 [doi:10.1107/S1600536812021071]

2-Amino-3-carboxypyrazin-1-i um perchlorate bis(2-aminopyrazin-1-i um-3-carboxylate) monohydrate

Fadila Berrah, Sofiane Bouacida, Ahlem Bouhraoua and Thierry Roisnel

S1. Comment

N-heterocyclic compounds such as pyrazine and its derivatives encompass a variety of potential hydrogen donors and acceptors which makes them interesting units to built new edifices involving original hydrogen-bonding schemes. As a continuation of our systematic studies concerning the synthesis and structural characterization of organic-inorganic hybrids and in attempts to establish a relationship between the nature of the anion used and hydrogen-bonding pattern encountered in these structures, we report herein the crystal structure of the compound 2-Amino-3-carboxypyrazin-1-i um perchlorate bis(2-aminopyrazin-1-i um 3-carboxylate) monohydrate. Related compounds obtained with nitrate, sulfate and dihydrogen phosphate anions have been reported (Berrah *et al.* 2011*a,b,c*).

The asymmetric unit, shown in Fig.1, comprises two symmetry independent zwitterions (B and C), one cation (A), one perchlorate anion and one water molecule. Bonds distances and angles in the three organic entities are comparable to that encountered in similar structures (Berrah *et al.* 2011*a,b,c*) except for the C—O distances in the carboxylic group: C—O distances are 1.2524 (19) and 1.2553 (19) Å in (B) and 1.2418 (19) and 1.2631 (19) Å in (C), due to the transfer of the carboxylic group proton to the hetero-ring nitrogen atom. Perchlorate anions present quite regular tetrahedral geometry (Cl—O distances range from 1.4279 (13) to 1.4528 (12) Å and angles from 108.67 (8) to 110.11 (9)°) and are comparable to that reported in the literature (Bendjeddou *et al.* 2003; Berrah *et al.* 2012; Toumi Akriche *et al.* 2010).


All components of the structure are involved in an interesting hydrogen bond system in which all potential donors and acceptors are implicated: the H₂O molecule, the two 2-Aminopyrazin-1-i um 3-carboxylate zwitterions (B and C) and the 2-Amino-3-carboxypyrazin-1-i um cation (A) act as both hydrogen bond donors and acceptors (table 1). An extensive H-bonding system between the three different organic entities, allows development of wave-like extended chains which intersect to form double-sheets parallel to (1̄10) (Fig.2 and Fig.3). Perchlorate anions and water molecules connect these double-sheets to generate a three-dimensional network (Fig.2 and Fig.3).

S2. Experimental


The title compound was obtained by reacting 3-amino-pyrazine 2-carboxylic acid with some excess of perchloric acid in aqueous solution. Slow evaporation allows growth of well formed colourless prismatic crystals.

S3. Refinement


All H atoms were located in difference Fourier maps. The water molecule H atoms were refined with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$ while all the remaining H atoms were introduced in calculated positions and treated as riding on their parent atoms (C,N or O) with C—H = 0.93 Å, N—H = 0.86 Å and O—H = 0.82 Å with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C or N})$ and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$.

**Figure 1**


The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines.

**Figure 2**

A partial view of one extended double-sheet showing wave-like chains and H-bonds between the different components. Hydrogen bonds are shown as dashed lines.

**Figure 3**

A view of part of the packing of (I) showing how double-sheets are linked via perchlorate anions and water molecules. Hydrogen bonds are shown as dashed lines.

2-Amino-3-carboxypyrazin-1-i um perchlorate bis(2-aminopyrazin-1-i um-3-carboxylate) monohydrate*Crystal data*
 $M_r = 535.83$
Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 8.1332 (14)$ Å $b = 11.816 (2)$ Å $c = 11.850 (2)$ Å $\alpha = 95.696 (9)^\circ$ $\beta = 108.148 (8)^\circ$ $\gamma = 102.416 (8)^\circ$ $V = 1039.8 (3)$ Å³ $Z = 2$ $F(000) = 552$ $D_x = 1.712 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7298 reflections

 $\theta = 2.7\text{--}27.4^\circ$ $\mu = 0.27 \text{ mm}^{-1}$ $T = 150$ K

Prism, colourless

 $0.46 \times 0.27 \times 0.17$ mm*Data collection*Bruker APEXII
diffractometer

Graphite monochromator

CCD rotation images, thin slices scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2002) $T_{\min} = 0.855$, $T_{\max} = 0.955$

15575 measured reflections

4705 independent reflections

4165 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.057$ $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.7^\circ$ $h = -10 \rightarrow 10$ $k = -15 \rightarrow 15$ $l = -14 \rightarrow 15$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.038$ $wR(F^2) = 0.1$ $S = 1.04$

4705 reflections

332 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0379P)^2 + 0.495P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.39 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.48 \text{ e } \text{\AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.69001 (5)	0.36663 (3)	-0.00929 (3)	0.01747 (11)
O3	0.68759 (17)	0.43096 (11)	0.09976 (11)	0.0282 (3)
C2A	0.07122 (19)	-0.13735 (13)	0.05029 (13)	0.0142 (3)

O2	0.62112 (16)	0.24144 (10)	-0.01437 (11)	0.0248 (3)
O1A	0.32377 (15)	0.00920 (10)	0.18281 (10)	0.0214 (3)
H1A	0.4049	0.0661	0.1869	0.032*
O1W	0.55592 (16)	0.19617 (10)	0.19895 (12)	0.0222 (3)
H1W	0.663 (3)	0.2094 (19)	0.254 (2)	0.033*
H2W	0.566 (3)	0.205 (2)	0.132 (2)	0.033*
O2C	0.40678 (15)	0.39784 (9)	0.29630 (10)	0.0185 (2)
N2C	0.88489 (17)	0.53957 (11)	0.58581 (12)	0.0170 (3)
H2C	0.9651	0.6013	0.588	0.02*
O1	0.57705 (18)	0.40133 (11)	-0.11220 (11)	0.0298 (3)
N3C	0.63598 (17)	0.34284 (12)	0.58168 (12)	0.0176 (3)
O1B	0.11825 (16)	-0.26089 (10)	0.61036 (11)	0.0235 (3)
C5B	0.3196 (2)	0.08607 (14)	0.75159 (14)	0.0183 (3)
H5B	0.3426	0.1123	0.833	0.022*
C5C	0.7831 (2)	0.38892 (15)	0.68069 (15)	0.0197 (3)
H5C	0.7998	0.3523	0.7478	0.024*
O4	0.86893 (17)	0.38880 (12)	-0.01070 (14)	0.0359 (3)
N1A	-0.01642 (18)	-0.16806 (12)	-0.17083 (12)	0.0212 (3)
H11A	-0.0909	-0.2062	-0.2398	0.025*
H12A	0.0748	-0.1132	-0.1671	0.025*
N2B	0.34558 (17)	0.12346 (11)	0.56691 (12)	0.0176 (3)
H2B	0.3852	0.1715	0.5257	0.021*
C1C	0.4396 (2)	0.34074 (13)	0.38055 (14)	0.0156 (3)
C4B	0.3817 (2)	0.16113 (14)	0.68530 (14)	0.0186 (3)
H4B	0.4485	0.2378	0.7214	0.022*
C2C	0.61173 (19)	0.39387 (13)	0.48635 (14)	0.0147 (3)
C3C	0.7406 (2)	0.49782 (13)	0.48389 (14)	0.0150 (3)
O2B	-0.00600 (16)	-0.22021 (10)	0.42749 (10)	0.0239 (3)
O1C	0.34056 (15)	0.24560 (10)	0.38619 (10)	0.0219 (3)
C3A	-0.04216 (19)	-0.19367 (13)	-0.07064 (14)	0.0150 (3)
N2A	-0.18714 (16)	-0.28048 (11)	-0.08043 (11)	0.0152 (3)
H2A	-0.2602	-0.3146	-0.1513	0.018*
N1C	0.72791 (18)	0.55293 (12)	0.39109 (12)	0.0199 (3)
H11C	0.8106	0.6143	0.3961	0.024*
H12C	0.6369	0.5274	0.3257	0.024*
C5A	-0.1068 (2)	-0.26064 (14)	0.12868 (14)	0.0178 (3)
H5A	-0.1291	-0.2857	0.1957	0.021*
N3B	0.22488 (17)	-0.02635 (12)	0.70059 (12)	0.0173 (3)
C1B	0.0919 (2)	-0.19231 (14)	0.53518 (14)	0.0172 (3)
O2A	0.25539 (16)	0.00695 (10)	-0.01483 (10)	0.0239 (3)
C2B	0.1914 (2)	-0.06368 (13)	0.58521 (14)	0.0156 (3)
N3A	0.03789 (17)	-0.17121 (11)	0.14528 (12)	0.0164 (3)
C1A	0.2274 (2)	-0.03376 (13)	0.06933 (14)	0.0154 (3)
C4C	0.9077 (2)	0.48821 (15)	0.68396 (15)	0.0200 (3)
H4C	1.0073	0.52	0.7532	0.024*
N1B	0.2181 (2)	-0.01866 (13)	0.39418 (13)	0.0263 (3)
H11B	0.2591	0.0311	0.3549	0.032*
H12B	0.1569	-0.0888	0.3577	0.032*

C3B	0.2491 (2)	0.01272 (14)	0.51060 (14)	0.0170 (3)
C4A	-0.2213 (2)	-0.31536 (13)	0.01601 (14)	0.0168 (3)
H4A	-0.3216	-0.376	0.0063	0.02*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.01646 (19)	0.01438 (19)	0.0198 (2)	0.00210 (14)	0.00545 (15)	0.00173 (14)
O3	0.0371 (7)	0.0237 (6)	0.0195 (6)	0.0060 (5)	0.0069 (5)	-0.0027 (5)
C2A	0.0137 (7)	0.0142 (7)	0.0136 (7)	0.0036 (6)	0.0033 (6)	0.0016 (6)
O2	0.0290 (6)	0.0143 (6)	0.0267 (7)	-0.0020 (5)	0.0092 (5)	0.0016 (5)
O1A	0.0193 (6)	0.0231 (6)	0.0145 (6)	-0.0055 (4)	0.0039 (4)	0.0011 (5)
O1W	0.0196 (6)	0.0226 (6)	0.0202 (6)	-0.0005 (5)	0.0047 (5)	0.0046 (5)
O2C	0.0207 (6)	0.0158 (5)	0.0141 (5)	0.0003 (4)	0.0019 (4)	0.0031 (4)
N2C	0.0142 (6)	0.0147 (6)	0.0187 (7)	-0.0012 (5)	0.0050 (5)	0.0012 (5)
O1	0.0366 (7)	0.0292 (7)	0.0208 (6)	0.0125 (6)	0.0033 (5)	0.0057 (5)
N3C	0.0171 (6)	0.0169 (7)	0.0171 (7)	0.0018 (5)	0.0052 (5)	0.0038 (5)
O1B	0.0256 (6)	0.0160 (6)	0.0206 (6)	-0.0040 (5)	0.0023 (5)	0.0050 (5)
C5B	0.0214 (8)	0.0167 (8)	0.0134 (7)	0.0013 (6)	0.0044 (6)	0.0004 (6)
C5C	0.0184 (8)	0.0233 (8)	0.0156 (8)	0.0037 (6)	0.0036 (6)	0.0062 (6)
O4	0.0208 (6)	0.0277 (7)	0.0630 (10)	0.0046 (5)	0.0204 (7)	0.0096 (7)
N1A	0.0205 (7)	0.0232 (7)	0.0126 (7)	-0.0041 (5)	0.0026 (5)	0.0016 (5)
N2B	0.0201 (7)	0.0144 (6)	0.0180 (7)	-0.0002 (5)	0.0084 (5)	0.0058 (5)
C1C	0.0167 (7)	0.0146 (7)	0.0137 (7)	0.0004 (6)	0.0057 (6)	0.0006 (6)
C4B	0.0206 (8)	0.0136 (7)	0.0171 (8)	-0.0005 (6)	0.0045 (6)	-0.0011 (6)
C2C	0.0151 (7)	0.0126 (7)	0.0154 (7)	0.0011 (6)	0.0058 (6)	0.0015 (6)
C3C	0.0156 (7)	0.0131 (7)	0.0155 (7)	0.0017 (6)	0.0063 (6)	0.0003 (6)
O2B	0.0263 (6)	0.0210 (6)	0.0156 (6)	-0.0051 (5)	0.0031 (5)	0.0006 (5)
O1C	0.0221 (6)	0.0186 (6)	0.0165 (6)	-0.0066 (4)	0.0029 (5)	0.0036 (4)
C3A	0.0148 (7)	0.0153 (7)	0.0147 (7)	0.0046 (6)	0.0043 (6)	0.0022 (6)
N2A	0.0139 (6)	0.0154 (6)	0.0127 (6)	0.0018 (5)	0.0016 (5)	0.0007 (5)
N1C	0.0192 (7)	0.0173 (7)	0.0176 (7)	-0.0042 (5)	0.0040 (5)	0.0051 (5)
C5A	0.0190 (7)	0.0193 (8)	0.0152 (8)	0.0019 (6)	0.0080 (6)	0.0035 (6)
N3B	0.0182 (6)	0.0166 (7)	0.0153 (7)	0.0011 (5)	0.0053 (5)	0.0034 (5)
C1B	0.0157 (7)	0.0164 (8)	0.0172 (8)	-0.0014 (6)	0.0070 (6)	0.0009 (6)
O2A	0.0270 (6)	0.0226 (6)	0.0164 (6)	-0.0036 (5)	0.0056 (5)	0.0055 (5)
C2B	0.0148 (7)	0.0155 (7)	0.0142 (7)	0.0002 (6)	0.0043 (6)	0.0020 (6)
N3A	0.0170 (6)	0.0171 (7)	0.0142 (6)	0.0032 (5)	0.0053 (5)	0.0022 (5)
C1A	0.0152 (7)	0.0150 (7)	0.0143 (7)	0.0035 (6)	0.0036 (6)	0.0003 (6)
C4C	0.0173 (7)	0.0226 (8)	0.0161 (8)	0.0035 (6)	0.0025 (6)	0.0002 (6)
N1B	0.0368 (8)	0.0218 (7)	0.0159 (7)	-0.0043 (6)	0.0116 (6)	0.0017 (6)
C3B	0.0166 (7)	0.0163 (8)	0.0157 (8)	0.0008 (6)	0.0048 (6)	0.0027 (6)
C4A	0.0150 (7)	0.0149 (7)	0.0202 (8)	0.0016 (6)	0.0072 (6)	0.0029 (6)

Geometric parameters (\AA , $^\circ$)

C11—O4	1.4279 (13)	N2B—C4B	1.347 (2)
C11—O1	1.4362 (13)	N2B—C3B	1.354 (2)

C11—O3	1.4406 (13)	N2B—H2B	0.86
C11—O2	1.4528 (12)	C1C—O1C	1.2553 (19)
C2A—N3A	1.317 (2)	C1C—C2C	1.516 (2)
C2A—C3A	1.439 (2)	C4B—H4B	0.93
C2A—C1A	1.504 (2)	C2C—C3C	1.442 (2)
O1A—C1A	1.3103 (18)	C3C—N1C	1.321 (2)
O1A—H1A	0.82	O2B—C1B	1.2418 (19)
O1W—H1W	0.88 (2)	C3A—N2A	1.352 (2)
O1W—H2W	0.84 (2)	N2A—C4A	1.340 (2)
O2C—C1C	1.2524 (19)	N2A—H2A	0.86
N2C—C4C	1.345 (2)	N1C—H11C	0.86
N2C—C3C	1.353 (2)	N1C—H12C	0.86
N2C—H2C	0.86	C5A—N3A	1.349 (2)
N3C—C2C	1.314 (2)	C5A—C4A	1.365 (2)
N3C—C5C	1.350 (2)	C5A—H5A	0.93
O1B—C1B	1.2631 (19)	N3B—C2B	1.317 (2)
C5B—N3B	1.354 (2)	C1B—C2B	1.520 (2)
C5B—C4B	1.361 (2)	O2A—C1A	1.2112 (19)
C5B—H5B	0.93	C2B—C3B	1.432 (2)
C5C—C4C	1.364 (2)	C4C—H4C	0.93
C5C—H5C	0.93	N1B—C3B	1.321 (2)
N1A—C3A	1.321 (2)	N1B—H11B	0.86
N1A—H11A	0.86	N1B—H12B	0.86
N1A—H12A	0.86	C4A—H4A	0.93
O4—Cl1—O1	110.11 (9)	N1C—C3C—C2C	125.21 (14)
O4—Cl1—O3	110.07 (8)	N2C—C3C—C2C	115.84 (14)
O1—Cl1—O3	109.55 (8)	N1A—C3A—N2A	118.17 (14)
O4—Cl1—O2	109.26 (8)	N1A—C3A—C2A	125.89 (14)
O1—Cl1—O2	108.67 (8)	N2A—C3A—C2A	115.95 (13)
O3—Cl1—O2	109.15 (8)	C4A—N2A—C3A	122.52 (13)
N3A—C2A—C3A	121.68 (14)	C4A—N2A—H2A	118.7
N3A—C2A—C1A	118.80 (13)	C3A—N2A—H2A	118.7
C3A—C2A—C1A	119.47 (13)	C3C—N1C—H11C	120
C1A—O1A—H1A	109.5	C3C—N1C—H12C	120
H1W—O1W—H2W	110 (2)	H11C—N1C—H12C	120
C4C—N2C—C3C	122.50 (14)	N3A—C5A—C4A	121.69 (14)
C4C—N2C—H2C	118.8	N3A—C5A—H5A	119.2
C3C—N2C—H2C	118.8	C4A—C5A—H5A	119.2
C2C—N3C—C5C	119.78 (14)	C2B—N3B—C5B	119.75 (14)
N3B—C5B—C4B	120.86 (15)	O2B—C1B—O1B	126.42 (15)
N3B—C5B—H5B	119.6	O2B—C1B—C2B	118.62 (14)
C4B—C5B—H5B	119.6	O1B—C1B—C2B	114.96 (13)
N3C—C5C—C4C	121.16 (15)	N3B—C2B—C3B	121.61 (14)
N3C—C5C—H5C	119.4	N3B—C2B—C1B	117.00 (14)
C4C—C5C—H5C	119.4	C3B—C2B—C1B	121.38 (14)
C3A—N1A—H11A	120	C2A—N3A—C5A	119.10 (13)
C3A—N1A—H12A	120	O2A—C1A—O1A	124.13 (14)

H11A—N1A—H12A	120	O2A—C1A—C2A	121.45 (14)
C4B—N2B—C3B	122.34 (14)	O1A—C1A—C2A	114.38 (13)
C4B—N2B—H2B	118.8	N2C—C4C—C5C	119.23 (14)
C3B—N2B—H2B	118.8	N2C—C4C—H4C	120.4
O2C—C1C—O1C	125.77 (14)	C5C—C4C—H4C	120.4
O2C—C1C—C2C	116.62 (13)	C3B—N1B—H11B	120
O1C—C1C—C2C	117.59 (13)	C3B—N1B—H12B	120
N2B—C4B—C5B	119.43 (14)	H11B—N1B—H12B	120
N2B—C4B—H4B	120.3	N1B—C3B—N2B	119.36 (14)
C5B—C4B—H4B	120.3	N1B—C3B—C2B	124.66 (15)
N3C—C2C—C3C	121.48 (14)	N2B—C3B—C2B	115.96 (14)
N3C—C2C—C1C	116.98 (13)	N2A—C4A—C5A	119.01 (14)
C3C—C2C—C1C	121.52 (13)	N2A—C4A—H4A	120.5
N1C—C3C—N2C	118.95 (14)	C5A—C4A—H4A	120.5
C2C—N3C—C5C—C4C	0.2 (2)	C5B—N3B—C2B—C1B	177.24 (13)
C3B—N2B—C4B—C5B	-0.1 (2)	O2B—C1B—C2B—N3B	150.06 (15)
N3B—C5B—C4B—N2B	1.2 (2)	O1B—C1B—C2B—N3B	-29.6 (2)
C5C—N3C—C2C—C3C	0.9 (2)	O2B—C1B—C2B—C3B	-31.1 (2)
C5C—N3C—C2C—C1C	-177.26 (14)	O1B—C1B—C2B—C3B	149.24 (15)
O2C—C1C—C2C—N3C	171.42 (14)	C3A—C2A—N3A—C5A	-0.3 (2)
O1C—C1C—C2C—N3C	-6.8 (2)	C1A—C2A—N3A—C5A	177.01 (13)
O2C—C1C—C2C—C3C	-6.8 (2)	C4A—C5A—N3A—C2A	-1.3 (2)
O1C—C1C—C2C—C3C	175.03 (14)	N3A—C2A—C1A—O2A	-173.93 (14)
C4C—N2C—C3C—N1C	-179.80 (14)	C3A—C2A—C1A—O2A	3.5 (2)
C4C—N2C—C3C—C2C	-0.2 (2)	N3A—C2A—C1A—O1A	3.9 (2)
N3C—C2C—C3C—N1C	178.63 (15)	C3A—C2A—C1A—O1A	-178.67 (13)
C1C—C2C—C3C—N1C	-3.3 (2)	C3C—N2C—C4C—C5C	1.3 (2)
N3C—C2C—C3C—N2C	-0.9 (2)	N3C—C5C—C4C—N2C	-1.3 (2)
C1C—C2C—C3C—N2C	177.19 (13)	C4B—N2B—C3B—N1B	179.88 (15)
N3A—C2A—C3A—N1A	-177.91 (14)	C4B—N2B—C3B—C2B	-1.7 (2)
C1A—C2A—C3A—N1A	4.8 (2)	N3B—C2B—C3B—N1B	-179.13 (16)
N3A—C2A—C3A—N2A	2.0 (2)	C1B—C2B—C3B—N1B	2.1 (2)
C1A—C2A—C3A—N2A	-175.33 (12)	N3B—C2B—C3B—N2B	2.5 (2)
N1A—C3A—N2A—C4A	177.75 (14)	C1B—C2B—C3B—N2B	-176.22 (13)
C2A—C3A—N2A—C4A	-2.1 (2)	C3A—N2A—C4A—C5A	0.7 (2)
C4B—C5B—N3B—C2B	-0.3 (2)	N3A—C5A—C4A—N2A	1.1 (2)
C5B—N3B—C2B—C3B	-1.6 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1A—H1A \cdots O1W	0.82	1.71	2.5258 (18)	172
O1W—H1W \cdots O1B ⁱ	0.88 (2)	1.92 (2)	2.7873 (19)	169 (2)
O1W—H2W \cdots O2	0.84 (2)	1.99 (2)	2.8176 (19)	172 (2)
O1W—H1W \cdots N3B ⁱ	0.88 (2)	2.56 (2)	3.052 (2)	115.8 (17)
N2B—H2B \cdots O1C	0.86	1.92	2.6935 (18)	149
N1C—H11C \cdots O2B ⁱⁱ	0.86	2.11	2.958 (2)	170

N2C—H2C···O1B ⁱⁱ	0.86	1.76	2.6156 (19)	171
N2A—H2A···O2C ⁱⁱⁱ	0.86	1.80	2.6536 (17)	175
N1A—H11A···O1C ⁱⁱⁱ	0.86	2.14	2.9340 (19)	153
N1B—H11B···O1A	0.86	2.26	2.916 (2)	133
N1B—H11B···O1C	0.86	2.44	3.087 (2)	133
N1B—H12B···O2B	0.86	2.21	2.814 (2)	127
N1A—H12A···O2A	0.86	2.09	2.7038 (19)	128
N1C—H12C···O2C	0.86	2.06	2.6734 (19)	128
N2B—H2B···N3C	0.86	2.41	3.058 (2)	132
N1B—H12B···N3A	0.86	2.41	3.035 (2)	130
N1A—H12A···N3B ^{iv}	0.86	2.44	3.152 (2)	140
C4B—H4B···O1 ^v	0.93	2.41	3.267 (2)	153
C4C—H4C···O3 ^{vi}	0.93	2.45	3.350 (2)	162
C5A—H5A···O2B	0.93	2.58	3.336 (2)	138
C5B—H5B···O2 ^v	0.93	2.48	3.145 (2)	129
C5B—H5B···O2A ^v	0.93	2.48	3.164 (2)	130

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x+1, y+1, z$; (iii) $-x, -y, -z$; (iv) $x, y, z-1$; (v) $x, y, z+1$; (vi) $-x+2, -y+1, -z+1$.