

Dimethylammonium dichloridotriphenylstannate(IV)

Yaya Sow,^a Libasse Diop,^a Gabriele Kociok-Kohn^b and Kieran C. Molloy^{b*}

^aLaboratoire de Chimie Minerale et Analytique (LACHIMIA), Département de Chimie, Faculté des Sciences et Techniques, Université Cheikh Anta Diop Dakar, Senegal, and ^bDepartment of Chemistry, University of Bath, Bath BA2 7AY, England
Correspondence e-mail: yayasow81@yahoo.fr


Received 21 May 2012; accepted 25 June 2012

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.021; wR factor = 0.045; data-to-parameter ratio = 20.1.

The title salt, $[(\text{CH}_3)_2\text{NH}_2][\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]$, was obtained as a by-product of the reaction between bis(dimethylammonium) oxalate and triphenyltin chloride. In the stannate anion, the trigonal-bipyramidal coordination environment of the Sn^{IV} atom is defined by the phenyl groups in equatorial and the Cl atoms in axial positions. The cations are connected to adjacent anions through $\text{N}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen-bonding interactions, leading to a chain motif parallel to [100].

Related literature

For background to organotin(IV) chemistry, see: Chee *et al.* (2003); Evans & Karpel (1985); Gielen *et al.* (1995); Ng & Kumar Das (1997); Zhang *et al.* (2006). For compounds containing the $[\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]^-$ ion, see: Harrison *et al.* (1978); Ng (1995, 1999).



Experimental

Crystal data

$(\text{C}_2\text{H}_8\text{N})[\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]$
 $M_r = 466.98$
Monoclinic, Cc
 $a = 7.9865$ (1) \AA
 $b = 17.5031$ (3) \AA
 $c = 14.9484$ (3) \AA
 $\beta = 105.406$ (1) $^\circ$

$$V = 2014.53$$
 (6) \AA^3

$$Z = 4$$

Mo $K\alpha$ radiation

$$\mu = 1.54 \text{ mm}^{-1}$$

$$T = 150 \text{ K}$$

$$0.30 \times 0.20 \times 0.20 \text{ mm}$$

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan (*SORTAV*; Blessing, 1995)
 $T_{\min} = 0.656$, $T_{\max} = 0.749$

16595 measured reflections
4569 independent reflections
4469 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.045$
 $S = 1.07$
4569 reflections
227 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.89 \text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
2256 Friedel pairs
Flack parameter: -0.030 (12)

Table 1
Selected bond lengths (\AA).

Sn—C7	2.152 (2)	Sn—Cl2	2.6098 (6)
Sn—Cl3	2.152 (2)	Sn—Cl1	2.6153 (6)
Sn—C1	2.160 (2)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N—H1A \cdots Cl1	0.89 (3)	2.33 (3)	3.203 (2)	167 (3)
N—H1B \cdots Cl2 ⁱ	0.82 (3)	2.34 (3)	3.143 (2)	164 (3)
C2—H2 \cdots Cl2	0.95	2.67	3.309 (3)	125
C6—H6 \cdots Cl1	0.95	2.76	3.376 (2)	123
C8—H8 \cdots Cl1	0.95	2.70	3.344 (2)	126
C12—H12 \cdots Cl2	0.95	2.69	3.340 (2)	126

Symmetry code: (i) $x - 1, y, z$.

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2636).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Chee, C. F., Lo, K. M. & Ng, S. W. (2003). *Acta Cryst. E* **59**, m36–m37.
- Evans, C. J. & Karpel, S. (1985). *Organotin Compounds in Modern Technology*. *J. Organomet. Chem. Library*, Vol. 16, Amsterdam: Elsevier.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Gielen, M., Bouhdid, A., Kayser, S., Biesemans, M., De Vos, D., Mahieu, B. & Willem, R. (1995). *Appl. Organomet. Chem.* **9**, 251–257.
- Harrison, P. G., Molloy, K. C., Phillips, R. C., Smith, P. J. & Crowe, A. J. (1978). *J. Organomet. Chem.* **160**, 421–434.
- Ng, S. W. (1995). *Acta Cryst. C* **51**, 1124–1125.
- Ng, S. W. (1999). *Acta Cryst. C* **55**, IUC9900098.
- Ng, S. W. & Kumar Das, V. G. (1997). *Acta Cryst. C* **53**, 1034–1036.

metal-organic compounds

- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zhang, W.-L., Ma, J.-F. & Jiang, H. (2006). *Acta Cryst. E* **62**, m460–m461.

supporting information

Acta Cryst. (2012). E68, m1015–m1016 [https://doi.org/10.1107/S1600536812028905]

Dimethylammonium dichloridotriphenylstannate(IV)

Yaya Sow, Libasse Diop, Gabriele Kociok-Kohn and Kieran C. Molloy

S1. Comment

Three $[\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]^-$ stannate(IV) anion-containing compounds with 2,2'-iminodipyridinium (Ng, 1999), triphenyl-(benzoylmethyl)arsonium (Harrison *et al.*, 1978) and tetramethylammonium (Ng, 1995), have previously been reported. In our research of new organotin(IV) compounds, driven by the various applications found within this family (Chee *et al.*, 2003; Evans & Karpel 1985; Gielen *et al.*, 1995; Ng *et al.*, 1997; Zhang *et al.*, 2006), we have initiated here the study of the interactions between bis(dimethylammonium)oxalate and triphenyltin chloride which has yielded the title ionic product, $[(\text{CH}_3)_2\text{NH}_2][\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]$, (I).

The $[\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]^-$ anion has a trigonal-bipyramidal shape with the Sn(IV) atom in a *trans*- Cl_2C_3 environment (Fig. 1). The equatorial plane is defined by the three phenyl groups [Sn—C 2.152 (2), 2.152 (2) and 2.160 (2) Å] while the Sn—Cl distances are 2.6098 (6) and 2.6153 (6) Å. The latter distances are very close to those reported by Ng (1995, 1999), [2.598 (1) Å] but somehow longer and shorter than those reported by Harrison *et al.* (1978) [2.573 (7), 2.689 (6) Å] for the same kind of anion. The sum of the equatorial angles (360°) indicates a planar SnPh_3 residue, although the Cl—Sn—Cl angle deviates from linearity [$174.94 (2)^\circ$].


The $[\text{SnPh}_3\text{Cl}_2]^-$ anions are connected by the ammonium cations through a pair of similar N—H···Cl hydrogen bonds leading to an infinite chain structure parallel to [100] (Fig. 2), which is probably the origin of the Sn—Cl bond lengthening in comparison with $[(\text{CH}_3)_4\text{N}][\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}_2]$. In the crystal packing C—H···Cl interactions are also observed (Table 1).

S2. Experimental


All chemicals were purchased from Aldrich (Germany) and used without any further purification. When $((\text{CH}_3)_2\text{NH}_2)_2\text{C}_2\text{O}_4\text{nH}_2\text{O}$ (obtained as a powder on submitting a 2/1 ratio mixture of $[(\text{CH}_3)_2\text{NH}_2][\text{OH}]$ and oxalic acid in water to evaporate at 333 K) is allowed to react while stirring with an excess of $\text{Sn}(\text{C}_6\text{H}_5)_3\text{Cl}$, both as ethanolic solutions, over 2 h, a precipitate is obtained. After filtering the precipitate, slow solvent evaporation from the filtrate afforded colourless crystals of the title complex suitable for X-ray work.

S3. Refinement

Hydrogen atoms bonded to the N atom have been located in difference Fourier maps and have been freely refined. All other hydrogen atoms have been placed onto calculated position and refined using a riding model, with C—H distances of 0.95 Å for sp^2 carbon atoms, or 0.98 Å for sp^3 carbon atoms, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for the sp^2 carbon atoms and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for the sp^3 carbon atoms.

**Figure 1**


Molecular structure of the complex showing the numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The packing of the structure showing N—H···Cl hydrogen bonding interactions as dashed lines.

Dimethylammonium dichloridotriphenylstannate(IV)

Crystal data



$M_r = 466.98$

Monoclinic, Cc

Hall symbol: C -2yc

$a = 7.9865$ (1) Å

$b = 17.5031$ (3) Å

$c = 14.9484$ (3) Å

$\beta = 105.406$ (1)°

$V = 2014.53$ (6) Å³

$Z = 4$

$F(000) = 936$

$D_x = 1.540 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 12072 reflections

$\theta = 2.9\text{--}27.5$ °

$\mu = 1.54 \text{ mm}^{-1}$

$T = 150$ K

Block, colourless

0.30 × 0.20 × 0.20 mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

298 2.0 degree images with φ and ω scans

Absorption correction: multi-scan
(SORTAV; Blessing, 1995)

$T_{\min} = 0.656$, $T_{\max} = 0.749$

16595 measured reflections

4569 independent reflections

4469 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 3.5$ °

$h = -10 \rightarrow 10$

$k = -22 \rightarrow 22$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.021$

$wR(F^2) = 0.045$

$S = 1.07$

4569 reflections

227 parameters

2 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0204P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.89 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack (1983), 2256 Friedel
pairs

Absolute structure parameter: -0.030 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}*/U_{\text{eq}}$
Sn	0.547277 (14)	0.045301 (7)	0.791813 (12)	0.01712 (5)
Cl1	0.25286 (7)	0.11982 (3)	0.75104 (4)	0.02502 (13)
Cl2	0.84601 (8)	-0.02426 (3)	0.84731 (4)	0.02268 (12)

C1	0.4134 (3)	-0.06314 (13)	0.77445 (16)	0.0196 (5)
C2	0.4971 (3)	-0.12794 (14)	0.75338 (18)	0.0251 (5)
H2	0.6146	-0.1243	0.7510	0.030*
C3	0.4122 (3)	-0.19752 (14)	0.73581 (18)	0.0293 (6)
H3	0.4699	-0.2405	0.7191	0.035*
C4	0.2430 (3)	-0.20458 (14)	0.74251 (19)	0.0278 (6)
H4	0.1850	-0.2524	0.7314	0.033*
C5	0.1592 (3)	-0.14099 (14)	0.76558 (18)	0.0269 (5)
H5	0.0438	-0.1454	0.7711	0.032*
C6	0.2438 (3)	-0.07124 (14)	0.78062 (17)	0.0229 (5)
H6	0.1845	-0.0280	0.7955	0.028*
C7	0.6122 (3)	0.08981 (12)	0.67105 (15)	0.0189 (4)
C8	0.4818 (3)	0.11348 (13)	0.59360 (16)	0.0229 (5)
H8	0.3639	0.1126	0.5957	0.028*
C9	0.5229 (3)	0.13846 (14)	0.51318 (17)	0.0269 (5)
H9	0.4331	0.1537	0.4607	0.032*
C10	0.6947 (3)	0.14098 (13)	0.51011 (17)	0.0262 (5)
H10	0.7225	0.1573	0.4552	0.031*
C11	0.8259 (3)	0.11967 (13)	0.58705 (18)	0.0259 (5)
H11	0.9440	0.1228	0.5856	0.031*
C12	0.7843 (3)	0.09383 (13)	0.66590 (16)	0.0222 (5)
H12	0.8750	0.0784	0.7179	0.027*
C13	0.6162 (3)	0.10926 (12)	0.91933 (16)	0.0209 (5)
C14	0.6982 (3)	0.07444 (14)	1.00306 (17)	0.0246 (5)
H14	0.7210	0.0211	1.0046	0.029*
C15	0.7473 (3)	0.11681 (15)	1.08452 (18)	0.0315 (6)
H15	0.8030	0.0922	1.1413	0.038*
C16	0.7159 (4)	0.19439 (15)	1.0836 (2)	0.0340 (6)
H16	0.7496	0.2231	1.1395	0.041*
C17	0.6349 (3)	0.23011 (15)	1.0009 (2)	0.0318 (6)
H17	0.6148	0.2836	0.9998	0.038*
C18	0.5825 (3)	0.18794 (13)	0.91879 (18)	0.0257 (5)
H18	0.5240	0.2126	0.8626	0.031*
N	0.1313 (3)	0.09715 (12)	0.93699 (16)	0.0265 (5)
H1A	0.172 (4)	0.0954 (16)	0.887 (2)	0.034 (8)*
H1B	0.043 (4)	0.0718 (19)	0.916 (2)	0.032 (8)*
C30	0.0840 (5)	0.17409 (17)	0.9603 (3)	0.0535 (9)
H30A	0.0163	0.1706	1.0061	0.080*
H30B	0.0144	0.1993	0.9042	0.080*
H30C	0.1897	0.2038	0.9863	0.080*
C20	0.2435 (5)	0.0556 (2)	1.0161 (3)	0.0560 (10)
H20A	0.3495	0.0852	1.0420	0.084*
H20B	0.2743	0.0058	0.9952	0.084*
H20C	0.1815	0.0482	1.0639	0.084*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn	0.01594 (7)	0.01814 (7)	0.01734 (7)	-0.00084 (7)	0.00453 (5)	-0.00073 (8)
C11	0.0184 (3)	0.0280 (3)	0.0286 (3)	0.0049 (2)	0.0062 (2)	0.0038 (3)
C12	0.0170 (3)	0.0253 (3)	0.0250 (3)	0.0014 (2)	0.0043 (2)	0.0007 (3)
C1	0.0209 (11)	0.0241 (11)	0.0128 (11)	-0.0049 (10)	0.0027 (9)	-0.0002 (9)
C2	0.0238 (13)	0.0236 (12)	0.0317 (14)	-0.0029 (10)	0.0142 (11)	-0.0051 (10)
C3	0.0320 (13)	0.0230 (12)	0.0346 (15)	0.0016 (10)	0.0117 (12)	-0.0050 (11)
C4	0.0284 (13)	0.0219 (12)	0.0314 (14)	-0.0096 (11)	0.0052 (11)	-0.0040 (11)
C5	0.0205 (11)	0.0284 (12)	0.0314 (14)	-0.0053 (10)	0.0059 (10)	-0.0008 (11)
C6	0.0231 (12)	0.0239 (12)	0.0228 (13)	0.0021 (10)	0.0077 (10)	0.0005 (11)
C7	0.0223 (11)	0.0157 (10)	0.0180 (11)	-0.0031 (9)	0.0042 (9)	-0.0025 (9)
C8	0.0220 (12)	0.0231 (12)	0.0227 (13)	-0.0005 (9)	0.0043 (10)	-0.0002 (10)
C9	0.0339 (14)	0.0250 (12)	0.0200 (12)	0.0022 (10)	0.0036 (11)	-0.0004 (10)
C10	0.0422 (15)	0.0213 (12)	0.0185 (12)	0.0026 (10)	0.0136 (11)	0.0019 (10)
C11	0.0252 (13)	0.0252 (12)	0.0300 (14)	-0.0044 (10)	0.0122 (11)	-0.0017 (10)
C12	0.0230 (12)	0.0222 (11)	0.0202 (13)	-0.0020 (9)	0.0039 (10)	0.0003 (10)
C13	0.0193 (11)	0.0233 (11)	0.0207 (12)	-0.0027 (9)	0.0063 (10)	-0.0023 (10)
C14	0.0282 (13)	0.0231 (12)	0.0228 (13)	-0.0026 (10)	0.0075 (11)	-0.0014 (10)
C15	0.0358 (15)	0.0377 (15)	0.0213 (13)	-0.0058 (11)	0.0080 (12)	-0.0039 (11)
C16	0.0386 (15)	0.0391 (15)	0.0264 (15)	-0.0107 (12)	0.0127 (12)	-0.0134 (12)
C17	0.0346 (14)	0.0244 (12)	0.0406 (17)	-0.0057 (11)	0.0173 (13)	-0.0134 (12)
C18	0.0269 (13)	0.0223 (12)	0.0296 (14)	0.0001 (10)	0.0105 (11)	0.0002 (10)
N	0.0232 (11)	0.0300 (11)	0.0258 (12)	-0.0030 (9)	0.0058 (10)	-0.0033 (10)
C30	0.059 (2)	0.0319 (16)	0.081 (3)	0.0027 (14)	0.039 (2)	-0.0061 (16)
C20	0.045 (2)	0.078 (3)	0.041 (2)	0.0102 (16)	0.0024 (17)	0.0204 (17)

Geometric parameters (\AA , $^\circ$)

Sn—C7	2.152 (2)	C11—C12	1.383 (3)
Sn—C13	2.152 (2)	C11—H11	0.9500
Sn—C1	2.160 (2)	C12—H12	0.9500
Sn—Cl2	2.6098 (6)	C13—C14	1.390 (3)
Sn—Cl1	2.6153 (6)	C13—C18	1.403 (3)
C1—C6	1.389 (3)	C14—C15	1.390 (4)
C1—C2	1.395 (3)	C14—H14	0.9500
C2—C3	1.385 (3)	C15—C16	1.380 (4)
C2—H2	0.9500	C15—H15	0.9500
C3—C4	1.387 (3)	C16—C17	1.384 (4)
C3—H3	0.9500	C16—H16	0.9500
C4—C5	1.389 (3)	C17—C18	1.397 (4)
C4—H4	0.9500	C17—H17	0.9500
C5—C6	1.385 (3)	C18—H18	0.9500
C5—H5	0.9500	N—C30	1.466 (4)
C6—H6	0.9500	N—C20	1.473 (4)
C7—C8	1.399 (3)	N—H1A	0.89 (3)
C7—C12	1.399 (3)	N—H1B	0.82 (3)

C8—C9	1.398 (3)	C30—H30A	0.9800
C8—H8	0.9500	C30—H30B	0.9800
C9—C10	1.386 (4)	C30—H30C	0.9800
C9—H9	0.9500	C20—H20A	0.9800
C10—C11	1.386 (4)	C20—H20B	0.9800
C10—H10	0.9500	C20—H20C	0.9800
C7—Sn—C13	119.52 (8)	C12—C11—H11	120.1
C7—Sn—C1	116.05 (8)	C10—C11—H11	120.1
C13—Sn—C1	124.43 (9)	C11—C12—C7	121.7 (2)
C7—Sn—Cl2	91.83 (6)	C11—C12—H12	119.2
C13—Sn—Cl2	87.94 (6)	C7—C12—H12	119.2
C1—Sn—Cl2	90.57 (7)	C14—C13—C18	118.7 (2)
C7—Sn—Cl1	91.49 (6)	C14—C13—Sn	121.19 (16)
C13—Sn—Cl1	87.10 (6)	C18—C13—Sn	120.14 (17)
C1—Sn—Cl1	91.40 (7)	C13—C14—C15	120.7 (2)
Cl2—Sn—Cl1	174.94 (2)	C13—C14—H14	119.6
C6—C1—C2	117.8 (2)	C15—C14—H14	119.6
C6—C1—Sn	122.82 (18)	C16—C15—C14	120.5 (3)
C2—C1—Sn	119.33 (16)	C16—C15—H15	119.7
C3—C2—C1	121.2 (2)	C14—C15—H15	119.7
C3—C2—H2	119.4	C15—C16—C17	119.6 (3)
C1—C2—H2	119.4	C15—C16—H16	120.2
C2—C3—C4	120.2 (2)	C17—C16—H16	120.2
C2—C3—H3	119.9	C16—C17—C18	120.4 (2)
C4—C3—H3	119.9	C16—C17—H17	119.8
C3—C4—C5	119.3 (2)	C18—C17—H17	119.8
C3—C4—H4	120.3	C17—C18—C13	120.1 (2)
C5—C4—H4	120.3	C17—C18—H18	120.0
C6—C5—C4	120.0 (2)	C13—C18—H18	120.0
C6—C5—H5	120.0	C30—N—C20	113.8 (3)
C4—C5—H5	120.0	C30—N—H1A	113.9 (18)
C5—C6—C1	121.4 (2)	C20—N—H1A	112.2 (19)
C5—C6—H6	119.3	C30—N—H1B	110 (2)
C1—C6—H6	119.3	C20—N—H1B	109 (2)
C8—C7—C12	117.8 (2)	H1A—N—H1B	97 (3)
C8—C7—Sn	120.62 (17)	N—C30—H30A	109.5
C12—C7—Sn	121.57 (17)	N—C30—H30B	109.5
C9—C8—C7	120.8 (2)	H30A—C30—H30B	109.5
C9—C8—H8	119.6	N—C30—H30C	109.5
C7—C8—H8	119.6	H30A—C30—H30C	109.5
C10—C9—C8	120.0 (2)	H30B—C30—H30C	109.5
C10—C9—H9	120.0	N—C20—H20A	109.5
C8—C9—H9	120.0	N—C20—H20B	109.5
C11—C10—C9	120.1 (2)	H20A—C20—H20B	109.5
C11—C10—H10	120.0	N—C20—H20C	109.5
C9—C10—H10	120.0	H20A—C20—H20C	109.5
C12—C11—C10	119.7 (2)	H20B—C20—H20C	109.5

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N—H1 <i>A</i> ···Cl1	0.89 (3)	2.33 (3)	3.203 (2)	167 (3)
N—H1 <i>B</i> ···Cl2 ⁱ	0.82 (3)	2.34 (3)	3.143 (2)	164 (3)
C2—H2···Cl2	0.95	2.67	3.309 (3)	125
C6—H6···Cl1	0.95	2.76	3.376 (2)	123
C8—H8···Cl1	0.95	2.70	3.344 (2)	126
C12—H12···Cl2	0.95	2.69	3.340 (2)	126

Symmetry code: (i) $x-1, y, z$.