

N-(3-Methylphenyl)-2-nitrobenzene-sulfonamide

U. Chaithanya,^a Sabine Foro^b and B. Thimme Gowda^{a*}

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdab@yahoo.com


Received 27 July 2012; accepted 30 July 2012

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.041; wR factor = 0.108; data-to-parameter ratio = 14.9.

In the title compound, $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_4\text{S}$, the dihedral angle between the benzene rings is $73.64(7)^\circ$. The amide H atom exhibits bifurcated hydrogen bonding: an intramolecular N—H···O hydrogen bond generates an $S(7)$ motif while in the crystal, N—H···O(S) hydrogen bonds link the molecules into zigzag $C(4)$ chains running along the b axis.

Related literature

For studies on the effects of substituents on the structures and other aspects of *N*-(aryl)-amides, see: Alkan *et al.* (2011); Bowes *et al.* (2003); Gowda *et al.* (2000); Saeed *et al.* (2010); Shahwar *et al.* (2012) of *N*-arylsulfonamides, see: Chaithanya *et al.* (2012); Gowda *et al.* (2002) and of *N*-chloroaryl-sulfonamides, see: Gowda & Shetty (2004); Shetty & Gowda (2004). For hydrogen-bonding patterns and motifs, see: Adsmond *et al.* (2001); Allen *et al.* (1998); Bernstein *et al.* (1995); Etter (1990).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_4\text{S}$

$M_r = 292.31$

Monoclinic, $P2_1/c$

$a = 9.0292(6)\text{ \AA}$

$b = 9.6498(7)\text{ \AA}$

$c = 15.878(1)\text{ \AA}$

$\beta = 103.763(8)^\circ$

$V = 1343.73(16)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.26\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.40 \times 0.28 \times 0.28\text{ mm}$

Data collection

Oxford Diffraction Xcalibur

diffractometer with a Sapphire

CCD detector

Absorption correction: multi-scan
(*CrysAlis RED*; Oxford)

Diffraction, 2009)

$T_{\min} = 0.905$, $T_{\max} = 0.932$

4756 measured reflections

2750 independent reflections

2316 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.011$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.108$

$S = 1.06$

2750 reflections

185 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.38\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}1-\text{H}1\text{N}\cdots\text{O}1^i$	0.84 (2)	2.30 (2)	3.055 (2)	151 (2)
$\text{N}1-\text{H}1\text{N}\cdots\text{O}3$	0.84 (2)	2.39 (2)	2.894 (2)	120 (2)

Symmetry code: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the University Grants Commission, Government of India, New Delhi, for a special grant under the UGC-BSR one-time grant to faculty.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5988).

References

- Adsmond, D. A. & Grant, D. J. W. (2001). *J. Pharm. Sci.* **90**, 2058–2077.
- Alkan, C., Tek, Y. & Kahraman, D. (2011). *Turk. J. Chem.* **35**, 769–777.
- Allen, F. H., Raithby, P. R., Shields, G. P. & Taylor, R. (1998). *Chem. Commun.* pp. 1043–1044.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bowes, K. F., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2003). *Acta Cryst. C* **59**, o1–o3.
- Chaithanya, U., Foro, S. & Gowda, B. T. (2012). *Acta Cryst. E* **68**, o2576.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Gowda, B. T., Jyothi, K. & D'Souza, J. D. (2002). *Z. Naturforsch. Teil A*, **57**, 967–973.
- Gowda, B. T., Paulus, H. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 791–800.
- Gowda, B. T. & Shetty, M. (2004). *J. Phys. Org. Chem.* **17**, 848–864.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Saeed, A., Arshad, M. & Simpson, J. (2010). *Acta Cryst. E* **66**, o2808–o2809.
- Shahwar, D., Tahir, M. N., Chohan, M. M., Ahmad, N. & Raza, M. A. (2012). *Acta Cryst. E* **68**, o1160.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shetty, M. & Gowda, B. T. (2004). *Z. Naturforsch. Teil B*, **59**, 63–72.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o2627 [doi:10.1107/S1600536812034009]

N-(3-Methylphenyl)-2-nitrobenzenesulfonamide

U. Chaithanya, Sabine Foro and B. Thimme Gowda

S1. Comment

As part of studying the substituent effects on the structures and other aspects of *N*-(aryl)-amides (Alkan *et al.*, 2011; Bowes *et al.*, 2003; Gowda *et al.*, 2000; Saeed *et al.*, 2010; Shahwar *et al.*, 2012); *N*-arylsulfonamides (Chaithanya *et al.*, 2012; Gowda *et al.*, 2002) and *N*-chloroarylsulfonamides (Gowda & Shetty, 2004; Shetty & Gowda, 2004), in the present work, the crystal structure of *N*-(3-methylphenyl)-2-nitrobenzenesulfonamide has been determined (Fig. 1).

The conformation of the N—C bond in the —SO₂—NH—C segment has *gauche* torsion with respect to the S=O bonds (Fig. 1), similar to that observed in *N*-(3-chlorophenyl)-2-nitrobenzenesulfonamide (I) (Chaithanya *et al.*, 2012). Further, the conformation of the N—H bond in the —SO₂—NH— segment is *syn* to both the *ortho*-nitro group in the sulfonyl benzene ring and *meta*-methyl group in the anilino ring. The molecule is twisted at the S—N bond with the torsional angle of 46.97 (16)°, compared to the value of 48.46 (18)° in (I).

The dihedral angle between the sulfonyl and the anilino rings is 73.64 (7)°, compared to the value of 73.65 (7)° in (I).

The amide H-atom showed bifurcated intramolecular H-bonding with the O-atom of the *ortho*-nitro group in the sulfonyl benzene ring, generating S(7) motifs and the intermolecular H-bonding with the sulfonyl oxygen atom of the other molecule, generating C(4) motifs (Adsmond *et al.*, 2001; Allen *et al.*, 1998; Bernstein *et al.*, 1995; Etter, 1990).

In the crystal, the intermolecular N—H···O (S) hydrogen bonds (Table 1) link the molecules into chains. Part of the crystal structure is shown in Fig. 2.


S2. Experimental

The title compound was prepared by treating 2-nitrobenzenesulfonylchloride with 3-methylaniline in the stoichiometric ratio and boiling the reaction mixture for 15 minutes. The reaction mixture was then cooled to room temperature and added to ice cold water (100 ml). The resultant solid *N*-(3-methylphenyl)-2-nitrobenzenesulfonamide was filtered under suction and washed thoroughly with cold water and dilute HCl to remove the excess sulfonylchloride and aniline, respectively. It was then recrystallized to constant melting point (114° C) from dilute ethanol. The purity of the compound was checked and characterized by its infrared spectra.


Prism like brown single crystals of the title compound used in X-ray diffraction studies were grown in ethanolic solution by slow evaporation of the solvent at room temperature.

S3. Refinement

H atoms bonded to C were positioned with idealized geometry using a riding model with C_{aromatic}—H = 0.93 Å, C_{methyl}—H = 0.96 Å. The coordinates of the amino H atom were refined with the N—H distance restrained to 0.86 (2) Å. All H atoms were refined with isotropic displacement parameters set at 1.2 *U*_{eq}(C-aromatic, N) and 1.5 *U*_{eq} (C-methyl) of the parent atom. The (10 3 1) reflection had a poor disagreement with its calculated value and was omitted from the refinement.

**Figure 1**

Molecular structure of the title compound, showing the atom labelling scheme and with displacement ellipsoids drawn at the 50% probability level.

**Figure 2**

Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

N-(3-Methylphenyl)-2-nitrobenzenesulfonamide

Crystal data

C₁₃H₁₂N₂O₄S
 $M_r = 292.31$
 Monoclinic, P2₁/c
 Hall symbol: -P 2ybc
 $a = 9.0292 (6) \text{ \AA}$
 $b = 9.6498 (7) \text{ \AA}$
 $c = 15.878 (1) \text{ \AA}$
 $\beta = 103.763 (8)^\circ$
 $V = 1343.73 (16) \text{ \AA}^3$
 $Z = 4$

$F(000) = 608$
 $D_x = 1.445 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 2679 reflections
 $\theta = 2.6\text{--}27.7^\circ$
 $\mu = 0.26 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
 Prism, brown
 $0.40 \times 0.28 \times 0.28 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur
 diffractometer with a Sapphire CCD detector
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Rotation method data acquisition using ω scans

Absorption correction: multi-scan
 (CrysAlis RED; Oxford Diffraction, 2009)
 $T_{\min} = 0.905$, $T_{\max} = 0.932$
 4756 measured reflections
 2750 independent reflections

2316 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$
 $\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.6^\circ$

$h = -11 \rightarrow 7$
 $k = -12 \rightarrow 8$
 $l = -15 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.108$
 $S = 1.06$
2750 reflections
185 parameters
1 restraint
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[c^2(F_o^2) + (0.0514P)^2 + 0.5996P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.30 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.38 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Absorption correction: CrysAlis RED (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3713 (2)	0.13280 (19)	0.58299 (11)	0.0375 (4)
C2	0.4187 (2)	0.18945 (19)	0.51278 (12)	0.0401 (4)
C3	0.3604 (2)	0.1431 (2)	0.42923 (13)	0.0524 (5)
H3	0.3953	0.1805	0.3835	0.063*
C4	0.2502 (3)	0.0412 (2)	0.41374 (15)	0.0581 (6)
H4	0.2099	0.0105	0.3574	0.070*
C5	0.1998 (3)	-0.0150 (2)	0.48125 (15)	0.0562 (5)
H5	0.1243	-0.0826	0.4706	0.067*
C6	0.2620 (2)	0.0294 (2)	0.56580 (14)	0.0481 (5)
H6	0.2295	-0.0111	0.6114	0.058*
C7	0.22779 (19)	0.35535 (19)	0.69038 (11)	0.0365 (4)
C8	0.1874 (2)	0.4794 (2)	0.64721 (12)	0.0425 (4)
H8	0.2624	0.5337	0.6323	0.051*
C9	0.0373 (2)	0.5240 (3)	0.62577 (14)	0.0570 (6)
C10	-0.0715 (2)	0.4398 (3)	0.64794 (18)	0.0735 (8)
H10	-0.1730	0.4678	0.6344	0.088*
C11	-0.0324 (3)	0.3147 (3)	0.68975 (19)	0.0726 (8)
H11	-0.1080	0.2593	0.7030	0.087*
C12	0.1188 (2)	0.2707 (2)	0.71234 (14)	0.0526 (5)

H12	0.1458	0.1872	0.7412	0.063*
C13	-0.0030 (4)	0.6597 (3)	0.5786 (2)	0.0886 (10)
H13A	0.0591	0.6723	0.5378	0.133*
H13B	0.0148	0.7345	0.6197	0.133*
H13C	-0.1086	0.6588	0.5483	0.133*
N1	0.38724 (17)	0.32157 (16)	0.71545 (10)	0.0381 (3)
H1N	0.444 (2)	0.3866 (19)	0.7069 (13)	0.046*
N2	0.53010 (19)	0.30367 (18)	0.52293 (11)	0.0471 (4)
O1	0.39747 (18)	0.06967 (15)	0.74298 (9)	0.0562 (4)
O2	0.61404 (15)	0.18892 (17)	0.70619 (9)	0.0558 (4)
O3	0.52327 (18)	0.39619 (17)	0.57411 (10)	0.0595 (4)
O4	0.6207 (2)	0.30177 (19)	0.47697 (13)	0.0743 (5)
S1	0.45362 (5)	0.17305 (5)	0.69406 (3)	0.04018 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0392 (9)	0.0359 (9)	0.0394 (9)	0.0107 (7)	0.0134 (7)	0.0037 (7)
C2	0.0391 (9)	0.0423 (10)	0.0417 (9)	0.0137 (8)	0.0148 (7)	0.0070 (8)
C3	0.0578 (12)	0.0615 (13)	0.0408 (10)	0.0197 (10)	0.0174 (9)	0.0054 (9)
C4	0.0625 (13)	0.0605 (13)	0.0479 (12)	0.0183 (11)	0.0064 (10)	-0.0108 (10)
C5	0.0533 (12)	0.0471 (12)	0.0662 (14)	0.0030 (10)	0.0103 (10)	-0.0119 (10)
C6	0.0506 (11)	0.0412 (10)	0.0568 (12)	0.0038 (9)	0.0210 (9)	0.0012 (9)
C7	0.0335 (9)	0.0429 (10)	0.0351 (9)	-0.0014 (7)	0.0122 (7)	-0.0093 (7)
C8	0.0410 (10)	0.0463 (10)	0.0412 (10)	0.0033 (8)	0.0118 (8)	-0.0075 (8)
C9	0.0468 (11)	0.0674 (14)	0.0543 (12)	0.0179 (10)	0.0068 (9)	-0.0180 (11)
C10	0.0357 (11)	0.092 (2)	0.0902 (19)	0.0104 (12)	0.0092 (11)	-0.0365 (16)
C11	0.0441 (12)	0.0830 (19)	0.100 (2)	-0.0230 (12)	0.0362 (13)	-0.0365 (16)
C12	0.0487 (11)	0.0513 (12)	0.0646 (13)	-0.0104 (9)	0.0271 (10)	-0.0111 (10)
C13	0.091 (2)	0.089 (2)	0.0801 (18)	0.0511 (17)	0.0098 (15)	-0.0004 (15)
N1	0.0341 (8)	0.0415 (8)	0.0400 (8)	-0.0020 (6)	0.0114 (6)	-0.0005 (7)
N2	0.0455 (9)	0.0512 (10)	0.0494 (9)	0.0109 (7)	0.0208 (7)	0.0171 (8)
O1	0.0734 (10)	0.0498 (8)	0.0495 (8)	0.0103 (7)	0.0229 (7)	0.0204 (7)
O2	0.0380 (7)	0.0738 (10)	0.0532 (8)	0.0161 (7)	0.0063 (6)	0.0103 (7)
O3	0.0679 (10)	0.0580 (9)	0.0585 (9)	-0.0082 (8)	0.0270 (8)	0.0015 (8)
O4	0.0710 (11)	0.0763 (12)	0.0945 (13)	0.0046 (9)	0.0569 (10)	0.0142 (10)
S1	0.0418 (3)	0.0434 (3)	0.0359 (2)	0.00993 (19)	0.01056 (18)	0.00990 (18)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.384 (3)	C9—C10	1.384 (4)
C1—C2	1.397 (2)	C9—C13	1.510 (4)
C1—S1	1.7861 (19)	C10—C11	1.382 (4)
C2—C3	1.380 (3)	C10—H10	0.9300
C2—N2	1.475 (3)	C11—C12	1.393 (3)
C3—C4	1.380 (3)	C11—H11	0.9300
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.371 (3)	C13—H13A	0.9600

C4—H4	0.9300	C13—H13B	0.9600
C5—C6	1.394 (3)	C13—H13C	0.9600
C5—H5	0.9300	N1—S1	1.6203 (16)
C6—H6	0.9300	N1—H1N	0.839 (15)
C7—C8	1.384 (3)	N2—O3	1.219 (2)
C7—C12	1.386 (3)	N2—O4	1.219 (2)
C7—N1	1.437 (2)	O1—S1	1.4293 (14)
C8—C9	1.385 (3)	O2—S1	1.4234 (14)
C8—H8	0.9300		
C6—C1—C2	117.66 (18)	C11—C10—C9	121.2 (2)
C6—C1—S1	117.47 (14)	C11—C10—H10	119.4
C2—C1—S1	124.62 (15)	C9—C10—H10	119.4
C3—C2—C1	121.41 (19)	C10—C11—C12	120.9 (2)
C3—C2—N2	116.18 (17)	C10—C11—H11	119.5
C1—C2—N2	122.40 (17)	C12—C11—H11	119.5
C4—C3—C2	119.8 (2)	C7—C12—C11	117.8 (2)
C4—C3—H3	120.1	C7—C12—H12	121.1
C2—C3—H3	120.1	C11—C12—H12	121.1
C5—C4—C3	120.1 (2)	C9—C13—H13A	109.5
C5—C4—H4	119.9	C9—C13—H13B	109.5
C3—C4—H4	119.9	H13A—C13—H13B	109.5
C4—C5—C6	120.0 (2)	C9—C13—H13C	109.5
C4—C5—H5	120.0	H13A—C13—H13C	109.5
C6—C5—H5	120.0	H13B—C13—H13C	109.5
C1—C6—C5	121.08 (19)	C7—N1—S1	122.60 (12)
C1—C6—H6	119.5	C7—N1—H1N	113.2 (14)
C5—C6—H6	119.5	S1—N1—H1N	111.0 (14)
C8—C7—C12	121.04 (18)	O3—N2—O4	123.90 (19)
C8—C7—N1	117.49 (16)	O3—N2—C2	118.62 (15)
C12—C7—N1	121.39 (18)	O4—N2—C2	117.44 (18)
C7—C8—C9	121.1 (2)	O2—S1—O1	118.86 (9)
C7—C8—H8	119.4	O2—S1—N1	106.96 (9)
C9—C8—H8	119.4	O1—S1—N1	107.87 (8)
C10—C9—C8	117.9 (2)	O2—S1—C1	109.16 (9)
C10—C9—C13	122.2 (2)	O1—S1—C1	105.46 (9)
C8—C9—C13	119.9 (2)	N1—S1—C1	108.15 (8)
C6—C1—C2—C3	-0.8 (3)	C8—C7—C12—C11	-0.2 (3)
S1—C1—C2—C3	173.28 (14)	N1—C7—C12—C11	176.49 (18)
C6—C1—C2—N2	177.64 (16)	C10—C11—C12—C7	-0.9 (3)
S1—C1—C2—N2	-8.3 (2)	C8—C7—N1—S1	-129.88 (15)
C1—C2—C3—C4	1.6 (3)	C12—C7—N1—S1	53.3 (2)
N2—C2—C3—C4	-176.88 (17)	C3—C2—N2—O3	139.47 (18)
C2—C3—C4—C5	-0.7 (3)	C1—C2—N2—O3	-39.0 (2)
C3—C4—C5—C6	-1.1 (3)	C3—C2—N2—O4	-38.1 (2)
C2—C1—C6—C5	-1.0 (3)	C1—C2—N2—O4	143.39 (19)
S1—C1—C6—C5	-175.49 (15)	C7—N1—S1—O2	164.42 (14)

C4—C5—C6—C1	1.9 (3)	C7—N1—S1—O1	−66.64 (16)
C12—C7—C8—C9	1.0 (3)	C7—N1—S1—C1	46.97 (16)
N1—C7—C8—C9	−175.76 (16)	C6—C1—S1—O2	136.48 (15)
C7—C8—C9—C10	−0.8 (3)	C2—C1—S1—O2	−37.61 (17)
C7—C8—C9—C13	−180.0 (2)	C6—C1—S1—O1	7.70 (16)
C8—C9—C10—C11	−0.3 (3)	C2—C1—S1—O1	−166.39 (15)
C13—C9—C10—C11	178.9 (2)	C6—C1—S1—N1	−107.49 (15)
C9—C10—C11—C12	1.2 (4)	C2—C1—S1—N1	78.42 (16)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O1 ⁱ	0.84 (2)	2.30 (2)	3.055 (2)	151 (2)
N1—H1N···O3	0.84 (2)	2.39 (2)	2.894 (2)	120 (2)

Symmetry code: (i) $-x+1, y+1/2, -z+3/2$.