

Di- μ -azido- $\kappa^4 N^1:N^1'$ -bis({1-[(*E*)-phenyl-(pyridin-2-yl- κN)methylidene]thiosemicarbazidato- $\kappa^2 N^1,S$ }copper(II))

Roji J. Kunnath,^a M. R. Prathapachandra Kurup^a and Seik Weng Ng^{b,c*}

^aDepartment of Applied Chemistry, Cochin University of Science and Technology, Kochi 682 022, India, ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^cChemistry Department, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia

Correspondence e-mail: seikweng@um.edu.my


Received 14 August 2012; accepted 14 August 2012

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.045; wR factor = 0.134; data-to-parameter ratio = 18.3.

In the title compound, $[\text{Cu}_2(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})_2(\text{N}_3)_2]$, the Cu^{II} cation is N,N',S -chelated by the deprotonated Schiff base ligand and is coordinated by the azide anion, while an N atom from an adjacent azide anion bridges the Cu^{II} cation at the apical position with a longer $\text{Cu}-\text{N}$ distance of 2.533 (3) Å, completing the distorted N_4S square-pyramidal coordination geometry. A pair of azide anions bridge the two Cu^{II} cations, forming a centrosymmetric binuclear molecule. In the crystal, the binuclear molecules are linked by an $\text{N}-\text{H}\cdots\text{N}$ hydrogen bond into a ribbon running along the a axis.

Related literature

For the structure of the parent Schiff base, see: Casas *et al.* (2003).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})_2(\text{N}_3)_2]$
 $M_r = 721.78$
 Monoclinic, $P2_1/c$
 $a = 11.2462$ (12) Å
 $b = 7.2344$ (10) Å
 $c = 18.519$ (2) Å
 $\beta = 96.653$ (5)°

$V = 1496.5$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.61$ mm⁻¹
 $T = 295$ K
 $0.35 \times 0.30 \times 0.25$ mm

Data collection

Bruker Kappa APEXII diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\text{min}} = 0.604$, $T_{\text{max}} = 0.690$

13614 measured reflections
 3747 independent reflections
 2973 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.134$
 $S = 1.04$
 3747 reflections
 205 parameters
 2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.54$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.66$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H1}\cdots\text{N2}^i$	0.87 (1)	2.22 (1)	3.075 (3)	168 (4)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

RJK thanks the University Grants Commission (India) for a Junior Research Fellowship. We thank the Sophisticated Analytical Instruments Facility, Cochin University of S & T, for the diffraction measurements. We also thank the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5611).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2010). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Casas, J. S., Castellano, E. E., Ellena, J., Tasende, M. S. G., Sánchez, A., Sordo, J. & Vidarte, M. J. (2003). *Inorg. Chem.* **42**, 2584–2595.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2012). E68, m1195 [doi:10.1107/S1600536812035751]

Di- μ -azido- $\kappa^4N^1:N^{1'}$ -bis({1-[(*E*)-phenyl(pyridin-2-yl- κN)methylidene]thiosemicarbazidato- κ^2N^1,S }copper(II))

Roji J. Kunnath, M. R. Prathapachandra Kurup and Seik Weng Ng

S1. Comment

2-Benzoylpyridine thiosemicarbazone (Casas *et al.*, 2003) is a Schiff base that is capable of N,N',S -chelation to metal ions. The Cu^{II} atom in $[\text{Cu}(\text{N}_3)(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})]_2$ (Scheme I) is N,N',S -chelated by the deprotonated Schiff base, and it exists in a square pyramidal environment (Fig. 1). Two molecules are disposed about a center-of-inversion and the distance between the copper atom and their apical nitrogen atom of the other azide is 2.533 (3) Å. Adjacent inversion-related pairs of molecules are linked by an $\text{N}-\text{H}\cdots\text{N}$ hydrogen bond to form a ribbon running along the a -axis (Table 1).

S2. Experimental

The Schiff base ligand by heating 2-benzoylpyridine (0.183 g, 1 mmol) and thiosemicarbazide (0.091 g, 1 mmol) for 3 h. Copper acetate hydrate (0.199 g, 1 mmol) and sodium azide (0.065 g, 1 mmol) was added and the solution heated for another 2 h. Dark green colored crystals were obtained from the cool solution.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions ($\text{C}-\text{H}$ 0.93 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U(\text{C})$.

The amino H-atoms were located in a difference Fourier and were refined with a distance restraint of $\text{N}-\text{H}$ 0.88 ± 0.01 Å; their temperature factors tied by a factor of 1.2 times.

Omitted owing interference from the beam stop was (1 0 0).




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $[\text{Cu}(\text{N}_3)(\text{C}_{13}\text{H}_{11}\text{N}_4\text{S})]_2$ at the 570% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Di- μ -azido- $\kappa^4N^1:N^1'$ -bis({1-[(E)-phenyl(pyridin- 2-yl- κN)methylidene]thiosemicarbazidato- κ^2N^1,S }copper(II))

Crystal data

[Cu₂(C₁₃H₁₁N₄S)₂(N₃)₂]
M_r = 721.78
 Monoclinic, *P*2₁/*c*
 Hall symbol: -P 2ybc
a = 11.2462 (12) Å
b = 7.2344 (10) Å
c = 18.519 (2) Å
 β = 96.653 (5)°
V = 1496.5 (3) Å³
Z = 2

F(000) = 732
D_x = 1.602 Mg m⁻³
 Mo *K* α radiation, λ = 0.71073 Å
 Cell parameters from 4663 reflections
 θ = 3.0–28.3°
 μ = 1.61 mm⁻¹
T = 295 K
 Prism, dark green
 0.35 × 0.30 × 0.25 mm

Data collection

Bruker Kappa APEXII
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
T_{min} = 0.604, *T_{max}* = 0.690

13614 measured reflections
 3747 independent reflections
 2973 reflections with *I* > 2 σ (*I*)
R_{int} = 0.075
 θ_{\max} = 28.4°, θ_{\min} = 2.2°
h = -14→15
k = -9→9
l = -24→24

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2 σ (*F*²)] = 0.045
wR(*F*²) = 0.134
S = 1.04
 3747 reflections
 205 parameters
 2 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0694P)^2 + 0.1161P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.54 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.66 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> [*] / <i>U_{eq}</i>
Cu1	0.40816 (2)	0.66511 (5)	0.521018 (15)	0.03737 (14)
S1	0.31362 (6)	0.55753 (11)	0.61376 (3)	0.0469 (2)
N1	0.24572 (17)	0.7151 (3)	0.47503 (10)	0.0345 (4)
N2	0.14750 (19)	0.6402 (3)	0.50005 (12)	0.0408 (5)
N3	0.0841 (2)	0.4802 (5)	0.59265 (14)	0.0617 (8)
H1	0.023 (2)	0.456 (6)	0.5610 (15)	0.074*
H2	0.101 (3)	0.422 (5)	0.6343 (12)	0.074*
N4	0.44641 (18)	0.8236 (3)	0.43679 (11)	0.0377 (5)
N5	0.5749 (2)	0.6199 (4)	0.55657 (12)	0.0468 (6)
N6	0.6113 (2)	0.5803 (4)	0.61823 (13)	0.0487 (6)
N7	0.6497 (3)	0.5408 (5)	0.67545 (14)	0.0721 (9)
C1	0.1174 (2)	0.8573 (3)	0.37564 (12)	0.0340 (5)

C2	0.0861 (3)	0.7823 (4)	0.30754 (14)	0.0454 (6)
H2A	0.1412	0.7109	0.2861	0.054*
C3	-0.0264 (3)	0.8127 (5)	0.27110 (15)	0.0506 (7)
H3	-0.0470	0.7627	0.2251	0.061*
C4	-0.1078 (2)	0.9171 (5)	0.30306 (16)	0.0506 (7)
H4	-0.1848	0.9339	0.2796	0.061*
C5	-0.0752 (3)	0.9963 (5)	0.36967 (17)	0.0556 (8)
H5	-0.1294	1.0707	0.3906	0.067*
C6	0.0364 (2)	0.9669 (4)	0.40563 (14)	0.0456 (6)
H6	0.0576	1.0216	0.4507	0.055*
C7	0.2345 (2)	0.8130 (3)	0.41657 (12)	0.0335 (5)
C8	0.1738 (2)	0.5609 (4)	0.56389 (13)	0.0414 (6)
C9	0.3482 (2)	0.8757 (4)	0.39278 (13)	0.0349 (5)
C10	0.3555 (2)	0.9761 (4)	0.33033 (14)	0.0438 (6)
H10	0.2866	1.0115	0.3010	0.053*
C11	0.4675 (3)	1.0232 (5)	0.31207 (16)	0.0510 (7)
H11	0.4750	1.0910	0.2702	0.061*
C12	0.5668 (3)	0.9688 (5)	0.35637 (17)	0.0546 (7)
H12	0.6428	0.9978	0.3446	0.065*
C13	0.5534 (2)	0.8708 (4)	0.41863 (16)	0.0460 (6)
H13	0.6214	0.8364	0.4490	0.055*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.03127 (19)	0.0431 (2)	0.0358 (2)	0.00031 (11)	-0.00433 (13)	0.00303 (12)
S1	0.0438 (4)	0.0589 (5)	0.0356 (3)	-0.0033 (3)	-0.0049 (3)	0.0100 (3)
N1	0.0312 (9)	0.0386 (12)	0.0328 (9)	-0.0013 (8)	-0.0002 (7)	0.0049 (8)
N2	0.0332 (10)	0.0478 (14)	0.0404 (11)	-0.0046 (9)	-0.0003 (8)	0.0132 (9)
N3	0.0541 (15)	0.079 (2)	0.0504 (14)	-0.0192 (14)	-0.0032 (11)	0.0287 (14)
N4	0.0325 (10)	0.0390 (13)	0.0410 (11)	-0.0018 (8)	0.0018 (8)	-0.0002 (8)
N5	0.0366 (11)	0.0595 (16)	0.0412 (12)	0.0036 (11)	-0.0092 (9)	0.0006 (11)
N6	0.0408 (11)	0.0518 (16)	0.0499 (13)	0.0046 (10)	-0.0099 (10)	-0.0094 (11)
N7	0.081 (2)	0.083 (2)	0.0462 (14)	0.0153 (16)	-0.0196 (13)	-0.0044 (14)
C1	0.0341 (11)	0.0368 (14)	0.0308 (11)	-0.0008 (9)	0.0026 (9)	0.0077 (9)
C2	0.0456 (14)	0.0489 (17)	0.0402 (13)	0.0035 (12)	-0.0013 (10)	-0.0058 (12)
C3	0.0508 (16)	0.061 (2)	0.0370 (13)	-0.0032 (13)	-0.0064 (11)	0.0039 (12)
C4	0.0367 (13)	0.060 (2)	0.0527 (15)	0.0040 (12)	-0.0052 (11)	0.0219 (14)
C5	0.0421 (14)	0.065 (2)	0.0608 (17)	0.0200 (14)	0.0120 (12)	0.0086 (15)
C6	0.0455 (14)	0.0547 (18)	0.0368 (12)	0.0050 (12)	0.0057 (10)	0.0012 (11)
C7	0.0319 (11)	0.0343 (14)	0.0337 (11)	0.0022 (9)	0.0012 (9)	0.0002 (9)
C8	0.0409 (13)	0.0449 (17)	0.0370 (12)	-0.0025 (11)	-0.0006 (10)	0.0075 (10)
C9	0.0348 (11)	0.0332 (13)	0.0369 (11)	-0.0001 (10)	0.0050 (9)	-0.0004 (9)
C10	0.0446 (14)	0.0453 (17)	0.0418 (13)	-0.0020 (11)	0.0062 (11)	0.0050 (11)
C11	0.0551 (17)	0.0497 (18)	0.0508 (15)	-0.0051 (13)	0.0171 (13)	0.0057 (13)
C12	0.0425 (14)	0.055 (2)	0.0695 (19)	-0.0090 (13)	0.0208 (13)	-0.0018 (15)
C13	0.0336 (12)	0.0482 (17)	0.0557 (16)	-0.0032 (11)	0.0034 (11)	-0.0029 (12)

Geometric parameters (Å, °)

Cu1—N5	1.942 (2)	C2—C3	1.381 (4)
Cu1—N1	1.9578 (19)	C2—H2A	0.9300
Cu1—N4	2.022 (2)	C3—C4	1.373 (4)
Cu1—S1	2.2603 (8)	C3—H3	0.9300
Cu1—N5 ⁱ	2.533 (3)	C4—C5	1.371 (5)
S1—C8	1.729 (3)	C4—H4	0.9300
N1—C7	1.287 (3)	C5—C6	1.368 (4)
N1—N2	1.359 (3)	C5—H5	0.9300
N2—C8	1.316 (3)	C6—H6	0.9300
N3—C8	1.329 (4)	C7—C9	1.472 (3)
N3—H1	0.872 (10)	C9—C10	1.376 (3)
N3—H2	0.879 (10)	C10—C11	1.384 (4)
N4—C13	1.331 (3)	C10—H10	0.9300
N4—C9	1.348 (3)	C11—C12	1.364 (4)
N5—N6	1.202 (3)	C11—H11	0.9300
N6—N7	1.134 (3)	C12—C13	1.377 (4)
C1—C6	1.372 (4)	C12—H12	0.9300
C1—C2	1.380 (4)	C13—H13	0.9300
C1—C7	1.477 (3)		
N5—Cu1—N1	173.91 (9)	C2—C3—H3	120.1
N5—Cu1—N4	94.22 (9)	C5—C4—C3	119.7 (2)
N1—Cu1—N4	80.26 (8)	C5—C4—H4	120.1
N5—Cu1—S1	101.84 (7)	C3—C4—H4	120.1
N1—Cu1—S1	84.08 (6)	C6—C5—C4	120.5 (3)
N4—Cu1—S1	160.44 (7)	C6—C5—H5	119.7
N5—Cu1—N5 ⁱ	85.54 (9)	C4—C5—H5	119.7
N1—Cu1—N5 ⁱ	91.79 (8)	C5—C6—C1	120.5 (3)
N4—Cu1—N5 ⁱ	89.28 (8)	C5—C6—H6	119.8
S1—Cu1—N5 ⁱ	102.92 (6)	C1—C6—H6	119.8
C8—S1—Cu1	93.90 (9)	N1—C7—C9	114.7 (2)
C7—N1—N2	120.1 (2)	N1—C7—C1	123.1 (2)
C7—N1—Cu1	117.58 (17)	C9—C7—C1	122.3 (2)
N2—N1—Cu1	122.15 (15)	N2—C8—N3	116.7 (2)
C8—N2—N1	111.9 (2)	N2—C8—S1	125.6 (2)
C8—N3—H1	114 (2)	N3—C8—S1	117.71 (19)
C8—N3—H2	118 (3)	N4—C9—C10	122.1 (2)
H1—N3—H2	124 (4)	N4—C9—C7	114.3 (2)
C13—N4—C9	118.5 (2)	C10—C9—C7	123.6 (2)
C13—N4—Cu1	128.30 (19)	C9—C10—C11	118.7 (3)
C9—N4—Cu1	113.11 (16)	C9—C10—H10	120.7
N6—N5—Cu1	124.8 (2)	C11—C10—H10	120.7
N7—N6—N5	177.3 (3)	C12—C11—C10	119.1 (3)
C6—C1—C2	119.1 (2)	C12—C11—H11	120.4
C6—C1—C7	120.8 (2)	C10—C11—H11	120.4
C2—C1—C7	120.1 (2)	C11—C12—C13	119.4 (3)

C1—C2—C3	120.4 (3)	C11—C12—H12	120.3
C1—C2—H2A	119.8	C13—C12—H12	120.3
C3—C2—H2A	119.8	N4—C13—C12	122.2 (3)
C4—C3—C2	119.8 (3)	N4—C13—H13	118.9
C4—C3—H3	120.1	C12—C13—H13	118.9
N5—Cu1—S1—C8	-167.21 (13)	C2—C1—C6—C5	2.2 (4)
N1—Cu1—S1—C8	11.35 (12)	C7—C1—C6—C5	-175.2 (3)
N4—Cu1—S1—C8	48.2 (2)	N2—N1—C7—C9	-176.1 (2)
N5 ⁱ —Cu1—S1—C8	-79.13 (11)	Cu1—N1—C7—C9	-0.5 (3)
N4—Cu1—N1—C7	1.37 (19)	N2—N1—C7—C1	3.0 (4)
S1—Cu1—N1—C7	169.6 (2)	Cu1—N1—C7—C1	178.59 (18)
N5 ⁱ —Cu1—N1—C7	-87.6 (2)	C6—C1—C7—N1	65.7 (4)
N4—Cu1—N1—N2	176.9 (2)	C2—C1—C7—N1	-111.7 (3)
S1—Cu1—N1—N2	-14.88 (19)	C6—C1—C7—C9	-115.3 (3)
N5 ⁱ —Cu1—N1—N2	87.9 (2)	C2—C1—C7—C9	67.3 (3)
C7—N1—N2—C8	-173.5 (2)	N1—N2—C8—N3	-178.8 (3)
Cu1—N1—N2—C8	11.1 (3)	N1—N2—C8—S1	2.6 (4)
N5—Cu1—N4—C13	-0.4 (2)	Cu1—S1—C8—N2	-11.5 (3)
N1—Cu1—N4—C13	-177.8 (3)	Cu1—S1—C8—N3	170.0 (3)
S1—Cu1—N4—C13	144.9 (2)	C13—N4—C9—C10	-0.3 (4)
N5 ⁱ —Cu1—N4—C13	-85.9 (2)	Cu1—N4—C9—C10	-176.6 (2)
N5—Cu1—N4—C9	175.44 (18)	C13—N4—C9—C7	178.6 (2)
N1—Cu1—N4—C9	-1.97 (17)	Cu1—N4—C9—C7	2.2 (3)
S1—Cu1—N4—C9	-39.2 (3)	N1—C7—C9—N4	-1.2 (3)
N5 ⁱ —Cu1—N4—C9	89.97 (18)	C1—C7—C9—N4	179.7 (2)
N4—Cu1—N5—N6	155.8 (3)	N1—C7—C9—C10	177.6 (3)
S1—Cu1—N5—N6	-13.0 (3)	C1—C7—C9—C10	-1.5 (4)
N5 ⁱ —Cu1—N5—N6	-115.3 (3)	N4—C9—C10—C11	0.6 (4)
C6—C1—C2—C3	-2.0 (4)	C7—C9—C10—C11	-178.1 (3)
C7—C1—C2—C3	175.5 (3)	C9—C10—C11—C12	0.0 (5)
C1—C2—C3—C4	-0.5 (5)	C10—C11—C12—C13	-0.9 (5)
C2—C3—C4—C5	2.6 (5)	C9—N4—C13—C12	-0.7 (4)
C3—C4—C5—C6	-2.3 (5)	Cu1—N4—C13—C12	175.0 (2)
C4—C5—C6—C1	-0.1 (5)	C11—C12—C13—N4	1.3 (5)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H1 ⁱⁱ —N2 ⁱⁱ	0.87 (1)	2.22 (1)	3.075 (3)	168 (4)

Symmetry code: (ii) $-x, -y+1, -z+1$.