

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Bis[(1,1'-biphenyl-2,2'-diyl)di-*tert*-butylphosphonium] di- $\mu$ -chlorido-bis[di-chloridopalladate(II)]

Charmaine Arderne\* and Cedric W. Holzapfel

Department of Chemistry, University of Johannesburg, P O Box 524, Auckland Park, Johannesburg, 2006, South Africa

Correspondence e-mail: carderne@uj.ac.za


Received 3 August 2012; accepted 3 September 2012

 Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.015;  $wR$  factor = 0.037; data-to-parameter ratio = 22.3.

In the title compound,  $(\text{C}_{20}\text{H}_{26}\text{P})_2[\text{Pd}_2\text{Cl}_6]$ , the  $\text{Pd}^{\text{II}}$  atom within the hexachloridodipalladate(II) dianion has a square-planar geometry. It resides on a centre of inversion with the asymmetric unit containing half of the dianion and one phosphonium cation. Only weak  $\text{C}-\text{H}\cdots\pi$  interactions are present in the crystal structure.

## Related literature

For the structures of related  $\text{Pd}_{\text{II}}$  complexes and background to organopalladium-catalysed reactions, see: Ormondi *et al.* (2011); Williams *et al.* (2008); Migowski & DuPont (2007); d'OrLyé & Jutland (2005); Beletskaya & Cheprakov (2004). For a description of the Cambridge Structural Database, see: Allen (2002).



## Experimental

### Crystal data

$(\text{C}_{20}\text{H}_{26}\text{P})_2[\text{Pd}_2\text{Cl}_6]$   
 $M_r = 1020.26$   
 Triclinic,  $P\bar{1}$   
 $a = 8.3247$  (2) Å  
 $b = 11.2697$  (2) Å

$c = 11.7004$  (3) Å  
 $\alpha = 73.0982$  (6)°  
 $\beta = 85.0900$  (6)°  
 $\gamma = 82.2708$  (5)°  
 $V = 1039.49$  (4) Å<sup>3</sup>

$Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.36$  mm<sup>-1</sup>

$T = 100$  K  
 $0.29 \times 0.22 \times 0.20$  mm

### Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (*AXScale*; Bruker, 2010)  
 $T_{\text{min}} = 0.695$ ,  $T_{\text{max}} = 0.774$

39747 measured reflections  
 5184 independent reflections  
 5088 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.017$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$   
 $wR(F^2) = 0.037$   
 $S = 1.05$   
 5184 reflections

232 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.48$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

 $Cg7$  is the centroid of the  $\text{Pd1}, \text{Cl3}, \text{Pd1}', \text{Cl3}'$  ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C11}-\text{H11}\cdots Cg7^i$	0.95	2.68	3.5952 (13)	138

 Symmetry code: (i)  $x + 1, y, z - 1$ .

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *pubCIF* (Westrip, 2010) and *PLATON* (Spek, 2009).

The authors wish to acknowledge the University of Johannesburg for the use of their facilities and for funding for this project.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GG2094).

## References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.  
 Beletskaya, I. P. & Cheprakov, A. V. (2004). *J. Organomet. Chem.* **689**, 4055–4082.  
 Bruker (2010). *APEX2*, *AXScale* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.  
 Migowski, P. & DuPont, J. (2007). *Chem. Eur. J.* **13**, 32–39.  
 OrLyé, F. d' & Jutland, A. (2005). *Tetrahedron*, **61**, 9670–9678.  
 Ormondi, B., Shaw, M. L. & Holzapfel, C. W. (2011). *J. Organomet. Chem.* **696**, 3091–3096.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.  
 Williams, D. B. G., Shaw, M. L., Green, M. J. & Holzapfel, C. W. (2008). *Angew. Chem. Int. Ed.* **47**, 560–563.

## supporting information

*Acta Cryst.* (2012). E68, m1247 [https://doi.org/10.1107/S1600536812037786]

## Bis[(1,1'-biphenyl-2,2'-diyl)di-*tert*-butylphosphonium] di- $\mu$ -chlorido-bis-[dichloridopalladate(II)]

Charmaine Arderne and Cedric W. Holzapfel

### S1. Comment

As part of our continued studies (Williams *et al.*, 2008 and Ormondi *et al.*, 2011) of organopalladium catalysed reactions, we have found that certain palladocycles (Beletskaya & Cheprakov, 2004 and d'OrLyé & Jutland, 2005) are readily converted into highly catalytically active low-ligated Pd<sup>0</sup> complexes. We now report that treatment of one such palladocycle, namely acetato-(2'-di-*t*-butylphosphino-1,1'-diphenyl-2yl) palladium(II), with HCl at room temperature results in the formation of the title compound (I) in good yield. Formation of the complex appears to result from the acid-induced reductive elimination of Pd<sup>0</sup> from the palladocycle followed by oxidation of the palladium in the presence of air and chloride ions (Migowski & DuPont, 2007).

The structure of the title compound (I), [C<sub>20</sub>H<sub>26</sub>P. 0.5(Cl<sub>6</sub>Pd<sub>2</sub>)]<sub>2</sub> shows a square planar geometry for the Pd<sup>II</sup> atom within the hexachlorodipalladium(II) anion. The palladium atom sits on a centre of inversion and therefore the asymmetric unit contains half of the trichloropalladium(II) anion and one phosphonium cation. Figure 1 shows a diagram of the molecular structure of the asymmetric unit of (I). Weak interactions were observed in this structure where C—H⋯Cl and C—H⋯ $\pi$  are evident only.

### S2. Experimental

A solution of hydrogen chloride (142 mg; 4 mmol) in 18 ml of methanol was slowly added to a stirred solution of the palladocycle precursor, namely acetato-(2'-di-*t*-butylphosphino-1,1'-diphenyl-2yl) palladium(II) (493 mg; 1 mmol) in 17 ml of dichloromethane over a period of 10 minutes. The reaction mixture changed from colourless to dark purple and then to dark brown. After completion of the addition, stirring was discontinued and the reaction mixture left exposed to the air at room temperature. A red crystalline precipitate started to form after 45 minutes. After 24 h, the supernatant solution was removed, the solid material was with ether and dried *in vacuo*. The solid material (376 mg; 74%) was taken up in 15 ml of 2:1 dichloromethane:methanol and the resulting solution was exposed to the vapours of diethyl ether in a closed system for 24 h. Well formed, dark red prisms of the title compound (I) crystallized from the solution and a suitable single-crystal was selected for the single-crystal X-ray diffraction analysis.

### S3. Refinement

The H-atoms were geometrically positioned and refined in the riding-model approximation, with C—H = 0.97 Å, N—H = 0.89 Å, and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{N})$ . For (I), the highest peak in the final difference map is 0.60 Å from Cl4B and the deepest hole is 0.27 Å from Cl4B.




Figure 1

Molecular structure of the asymmetric unit of the title compound (I) with thermal displacement ellipsoids drawn at the 50% probability level.

**Bis[(1,1'-biphenyl-2,2'-diyl)di-*tert*-butylphosphonium] di- $\mu$ -chlorido-bis[dichloridopalladate(II)]**

*Crystal data*

(C<sub>20</sub>H<sub>26</sub>P)<sub>2</sub>[Pd<sub>2</sub>Cl<sub>6</sub>]

*M<sub>r</sub>* = 1020.26

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

*a* = 8.3247 (2) Å

*b* = 11.2697 (2) Å

*c* = 11.7004 (3) Å

$\alpha$  = 73.0982 (6)°

$\beta$  = 85.0900 (6)°

$\gamma$  = 82.2708 (5)°

*V* = 1039.49 (4) Å<sup>3</sup>

*Z* = 1

*F*(000) = 516

*D<sub>x</sub>* = 1.630 Mg m<sup>-3</sup>

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 9253 reflections

$\theta$  = 3.6–28.4°

$\mu$  = 1.36 mm<sup>-1</sup>

$T = 100$  K  $0.29 \times 0.22 \times 0.20$  mm  
 Prism, dark orange

*Data collection*

Bruker APEXII CCD diffractometer	39747 measured reflections
Radiation source: fine-focus sealed tube	5184 independent reflections
Graphite monochromator	5088 reflections with $I > 2\sigma(I)$
$\varphi$ and $\omega$ scans	$R_{\text{int}} = 0.017$
Absorption correction: multi-scan (AXScale; Bruker, 2010)	$\theta_{\text{max}} = 28.5^\circ$ , $\theta_{\text{min}} = 2.9^\circ$
$T_{\text{min}} = 0.695$ , $T_{\text{max}} = 0.774$	$h = -11 \rightarrow 11$
	$k = -15 \rightarrow 15$
	$l = -15 \rightarrow 15$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.015$	H-atom parameters constrained
$wR(F^2) = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.0139P)^2 + 0.6282P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
5184 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
232 parameters	$\Delta\rho_{\text{max}} = 0.41 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.48 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C16	0.06541 (14)	0.17413 (10)	0.35252 (11)	0.0171 (2)
H16A	0.1167	0.1453	0.4297	0.026*
H16B	0.1288	0.1356	0.2953	0.026*
H16C	-0.0451	0.1504	0.3628	0.026*
C2	0.51131 (13)	0.18374 (10)	0.38452 (10)	0.0150 (2)
C6	0.36510 (14)	0.28059 (11)	0.52998 (11)	0.0182 (2)
H6	0.2787	0.3363	0.5501	0.022*
C1	0.38564 (13)	0.26782 (10)	0.41465 (10)	0.0147 (2)
C3	0.61953 (14)	0.11512 (11)	0.47087 (11)	0.0186 (2)
H3	0.7057	0.0588	0.4515	0.022*
C5	0.47340 (15)	0.21017 (12)	0.61544 (11)	0.0213 (2)
H5	0.4601	0.2173	0.6947	0.026*
C4	0.60052 (15)	0.12966 (12)	0.58561 (11)	0.0215 (2)
H4	0.6753	0.0840	0.6441	0.026*

C17	0.32342 (14)	0.50983 (10)	0.22795 (11)	0.0161 (2)
C13	0.05899 (13)	0.31677 (10)	0.30507 (10)	0.0137 (2)
C7	0.51087 (13)	0.17674 (10)	0.26061 (10)	0.0149 (2)
C19	0.25132 (15)	0.58473 (11)	0.31437 (12)	0.0215 (2)
H19A	0.2870	0.5412	0.3951	0.032*
H19B	0.1326	0.5930	0.3149	0.032*
H19C	0.2883	0.6678	0.2884	0.032*
C18	0.26103 (15)	0.57096 (11)	0.10206 (11)	0.0211 (2)
H18A	0.1421	0.5790	0.1061	0.032*
H18B	0.3034	0.5190	0.0496	0.032*
H18C	0.2977	0.6539	0.0701	0.032*
C15	-0.03314 (14)	0.37709 (11)	0.39742 (11)	0.0181 (2)
H15A	-0.0455	0.4681	0.3642	0.027*
H15B	0.0279	0.3537	0.4702	0.027*
H15C	-0.1406	0.3479	0.4167	0.027*
C12	0.61554 (14)	0.09816 (11)	0.20828 (11)	0.0190 (2)
H12	0.7014	0.0445	0.2517	0.023*
C14	-0.02701 (14)	0.36283 (11)	0.18602 (11)	0.0180 (2)
H14A	0.0365	0.3286	0.1260	0.027*
H14B	-0.0369	0.4542	0.1584	0.027*
H14C	-0.1353	0.3350	0.1979	0.027*
C20	0.50976 (14)	0.50237 (12)	0.22269 (13)	0.0228 (2)
H20A	0.5562	0.4503	0.1713	0.034*
H20B	0.5491	0.4656	0.3034	0.034*
H20C	0.5426	0.5865	0.1900	0.034*
C9	0.36534 (14)	0.25847 (10)	0.07624 (10)	0.0157 (2)
H9	0.2817	0.3136	0.0314	0.019*
C8	0.38668 (13)	0.25742 (10)	0.19316 (10)	0.0137 (2)
C10	0.46880 (14)	0.17729 (11)	0.02614 (11)	0.0183 (2)
H10	0.4545	0.1754	-0.0530	0.022*
C11	0.59299 (15)	0.09897 (11)	0.09144 (12)	0.0205 (2)
H11	0.6639	0.0449	0.0557	0.025*
Cl3	0.06462 (4)	0.07944 (3)	1.07304 (2)	0.01962 (6)
Cl2	0.15259 (3)	0.29365 (2)	0.81900 (2)	0.01663 (5)
Cl1	0.01589 (3)	0.12552 (3)	0.67310 (2)	0.01763 (6)
P1	0.27397 (3)	0.34725 (3)	0.28353 (2)	0.01179 (5)
Pd1	0.041384 (9)	0.110164 (7)	0.868729 (7)	0.01193 (3)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C16	0.0169 (5)	0.0132 (5)	0.0208 (5)	-0.0040 (4)	0.0012 (4)	-0.0037 (4)
C2	0.0129 (5)	0.0132 (5)	0.0177 (5)	-0.0035 (4)	-0.0001 (4)	-0.0018 (4)
C6	0.0177 (5)	0.0209 (6)	0.0170 (5)	-0.0038 (4)	-0.0011 (4)	-0.0060 (4)
C1	0.0135 (5)	0.0149 (5)	0.0151 (5)	-0.0025 (4)	-0.0017 (4)	-0.0025 (4)
C3	0.0134 (5)	0.0170 (5)	0.0230 (6)	-0.0027 (4)	-0.0025 (4)	-0.0010 (4)
C5	0.0214 (6)	0.0260 (6)	0.0173 (5)	-0.0072 (5)	-0.0036 (4)	-0.0046 (5)
C4	0.0180 (5)	0.0228 (6)	0.0213 (6)	-0.0059 (4)	-0.0071 (4)	0.0009 (5)

C17	0.0149 (5)	0.0123 (5)	0.0208 (5)	-0.0032 (4)	0.0008 (4)	-0.0042 (4)
C13	0.0118 (5)	0.0128 (5)	0.0162 (5)	-0.0021 (4)	0.0006 (4)	-0.0037 (4)
C7	0.0127 (5)	0.0126 (5)	0.0181 (5)	-0.0020 (4)	0.0003 (4)	-0.0026 (4)
C19	0.0229 (6)	0.0161 (5)	0.0280 (6)	-0.0024 (4)	-0.0005 (5)	-0.0103 (5)
C18	0.0236 (6)	0.0157 (5)	0.0210 (6)	-0.0038 (4)	0.0000 (5)	-0.0001 (4)
C15	0.0161 (5)	0.0174 (5)	0.0204 (6)	-0.0012 (4)	0.0035 (4)	-0.0064 (4)
C12	0.0152 (5)	0.0159 (5)	0.0242 (6)	0.0015 (4)	0.0009 (4)	-0.0049 (5)
C14	0.0151 (5)	0.0192 (5)	0.0193 (5)	-0.0013 (4)	-0.0033 (4)	-0.0043 (4)
C20	0.0151 (5)	0.0189 (6)	0.0337 (7)	-0.0055 (4)	0.0013 (5)	-0.0052 (5)
C9	0.0152 (5)	0.0149 (5)	0.0169 (5)	-0.0023 (4)	0.0007 (4)	-0.0043 (4)
C8	0.0127 (5)	0.0117 (5)	0.0163 (5)	-0.0015 (4)	0.0014 (4)	-0.0039 (4)
C10	0.0193 (5)	0.0188 (5)	0.0184 (5)	-0.0039 (4)	0.0034 (4)	-0.0082 (4)
C11	0.0187 (5)	0.0173 (5)	0.0255 (6)	-0.0001 (4)	0.0045 (4)	-0.0088 (5)
Cl3	0.03037 (15)	0.01788 (13)	0.01368 (12)	-0.01237 (11)	0.00371 (10)	-0.00635 (10)
Cl2	0.01830 (12)	0.01205 (11)	0.01884 (13)	-0.00401 (9)	-0.00206 (10)	-0.00193 (10)
Cl1	0.02204 (13)	0.01686 (12)	0.01351 (12)	-0.00352 (10)	-0.00294 (9)	-0.00237 (10)
P1	0.01112 (12)	0.01095 (12)	0.01294 (12)	-0.00073 (9)	-0.00015 (9)	-0.00320 (10)
Pd1	0.01306 (4)	0.01056 (4)	0.01228 (4)	-0.00235 (3)	0.00107 (3)	-0.00344 (3)

*Geometric parameters (Å, °)*

C16—C13	1.5356 (15)	C19—H19C	0.9800
C16—H16A	0.9800	C18—H18A	0.9800
C16—H16B	0.9800	C18—H18B	0.9800
C16—H16C	0.9800	C18—H18C	0.9800
C2—C3	1.3945 (15)	C15—H15A	0.9800
C2—C1	1.4075 (15)	C15—H15B	0.9800
C2—C7	1.4751 (16)	C15—H15C	0.9800
C6—C1	1.3920 (16)	C12—C11	1.3930 (18)
C6—C5	1.3939 (16)	C12—H12	0.9500
C6—H6	0.9500	C14—H14A	0.9800
C1—P1	1.8003 (11)	C14—H14B	0.9800
C3—C4	1.3924 (18)	C14—H14C	0.9800
C3—H3	0.9500	C20—H20A	0.9800
C5—C4	1.3884 (18)	C20—H20B	0.9800
C5—H5	0.9500	C20—H20C	0.9800
C4—H4	0.9500	C9—C8	1.3911 (16)
C17—C19	1.5335 (16)	C9—C10	1.3913 (16)
C17—C18	1.5357 (17)	C9—H9	0.9500
C17—C20	1.5391 (16)	C8—P1	1.7945 (11)
C17—P1	1.8476 (11)	C10—C11	1.3890 (17)
C13—C15	1.5371 (15)	C10—H10	0.9500
C13—C14	1.5404 (15)	C11—H11	0.9500
C13—P1	1.8500 (11)	Cl3—Pd1 <sup>i</sup>	2.3166 (3)
C7—C12	1.3908 (16)	Cl3—Pd1	2.3349 (3)
C7—C8	1.4100 (15)	Cl2—Pd1	2.2791 (3)
C19—H19A	0.9800	Cl1—Pd1	2.2709 (3)
C19—H19B	0.9800	Pd1—Cl3 <sup>i</sup>	2.3166 (3)

C13—C16—H16A	109.5	H18A—C18—H18C	109.5
C13—C16—H16B	109.5	H18B—C18—H18C	109.5
H16A—C16—H16B	109.5	C13—C15—H15A	109.5
C13—C16—H16C	109.5	C13—C15—H15B	109.5
H16A—C16—H16C	109.5	H15A—C15—H15B	109.5
H16B—C16—H16C	109.5	C13—C15—H15C	109.5
C3—C2—C1	119.27 (11)	H15A—C15—H15C	109.5
C3—C2—C7	126.57 (11)	H15B—C15—H15C	109.5
C1—C2—C7	114.15 (10)	C7—C12—C11	119.26 (11)
C1—C6—C5	118.86 (11)	C7—C12—H12	120.4
C1—C6—H6	120.6	C11—C12—H12	120.4
C5—C6—H6	120.6	C13—C14—H14A	109.5
C6—C1—C2	121.02 (10)	C13—C14—H14B	109.5
C6—C1—P1	130.24 (9)	H14A—C14—H14B	109.5
C2—C1—P1	108.73 (8)	C13—C14—H14C	109.5
C4—C3—C2	119.62 (11)	H14A—C14—H14C	109.5
C4—C3—H3	120.2	H14B—C14—H14C	109.5
C2—C3—H3	120.2	C17—C20—H20A	109.5
C4—C5—C6	120.51 (12)	C17—C20—H20B	109.5
C4—C5—H5	119.7	H20A—C20—H20B	109.5
C6—C5—H5	119.7	C17—C20—H20C	109.5
C5—C4—C3	120.69 (11)	H20A—C20—H20C	109.5
C5—C4—H4	119.7	H20B—C20—H20C	109.5
C3—C4—H4	119.7	C8—C9—C10	118.76 (11)
C19—C17—C18	110.87 (10)	C8—C9—H9	120.6
C19—C17—C20	109.39 (10)	C10—C9—H9	120.6
C18—C17—C20	109.59 (10)	C9—C8—C7	121.31 (10)
C19—C17—P1	110.43 (8)	C9—C8—P1	129.83 (9)
C18—C17—P1	110.18 (8)	C7—C8—P1	108.86 (8)
C20—C17—P1	106.27 (8)	C11—C10—C9	120.21 (11)
C16—C13—C15	109.56 (9)	C11—C10—H10	119.9
C16—C13—C14	109.59 (9)	C9—C10—H10	119.9
C15—C13—C14	109.98 (9)	C10—C11—C12	121.24 (11)
C16—C13—P1	104.76 (7)	C10—C11—H11	119.4
C15—C13—P1	111.52 (8)	C12—C11—H11	119.4
C14—C13—P1	111.29 (8)	Pd1 <sup>i</sup> —C13—Pd1	94.884 (10)
C12—C7—C8	119.19 (11)	C8—P1—C1	93.92 (5)
C12—C7—C2	126.66 (10)	C8—P1—C17	108.77 (5)
C8—C7—C2	114.12 (10)	C1—P1—C17	109.07 (5)
C17—C19—H19A	109.5	C8—P1—C13	110.90 (5)
C17—C19—H19B	109.5	C1—P1—C13	111.69 (5)
H19A—C19—H19B	109.5	C17—P1—C13	119.48 (5)
C17—C19—H19C	109.5	C11—Pd1—C12	91.333 (10)
H19A—C19—H19C	109.5	C11—Pd1—C13 <sup>i</sup>	90.924 (10)
H19B—C19—H19C	109.5	C12—Pd1—C13 <sup>i</sup>	177.377 (10)
C17—C18—H18A	109.5	C11—Pd1—C13	175.919 (10)
C17—C18—H18B	109.5	C12—Pd1—C13	92.603 (10)

H18A—C18—H18B	109.5	C13 <sup>i</sup> —Pd1—C13	85.117 (10)
C17—C18—H18C	109.5		
C5—C6—C1—C2	1.22 (17)	C7—C8—P1—C17	107.86 (8)
C5—C6—C1—P1	-177.59 (9)	C9—C8—P1—C13	60.63 (12)
C3—C2—C1—C6	-1.94 (17)	C7—C8—P1—C13	-118.81 (8)
C7—C2—C1—C6	176.93 (10)	C6—C1—P1—C8	-176.59 (11)
C3—C2—C1—P1	177.11 (9)	C2—C1—P1—C8	4.49 (9)
C7—C2—C1—P1	-4.03 (12)	C6—C1—P1—C17	72.01 (12)
C1—C2—C3—C4	0.77 (17)	C2—C1—P1—C17	-106.92 (8)
C7—C2—C3—C4	-177.94 (11)	C6—C1—P1—C13	-62.25 (12)
C1—C6—C5—C4	0.65 (18)	C2—C1—P1—C13	118.82 (8)
C6—C5—C4—C3	-1.81 (19)	C19—C17—P1—C8	-170.94 (8)
C2—C3—C4—C5	1.08 (18)	C18—C17—P1—C8	66.24 (9)
C3—C2—C7—C12	1.82 (19)	C20—C17—P1—C8	-52.40 (9)
C1—C2—C7—C12	-176.95 (11)	C19—C17—P1—C1	-69.77 (9)
C3—C2—C7—C8	-179.97 (11)	C18—C17—P1—C1	167.42 (8)
C1—C2—C7—C8	1.26 (14)	C20—C17—P1—C1	48.77 (9)
C8—C7—C12—C11	-1.44 (17)	C19—C17—P1—C13	60.37 (10)
C2—C7—C12—C11	176.69 (11)	C18—C17—P1—C13	-62.44 (10)
C10—C9—C8—C7	0.39 (16)	C20—C17—P1—C13	178.91 (8)
C10—C9—C8—P1	-178.98 (9)	C16—C13—P1—C8	52.91 (9)
C12—C7—C8—C9	1.04 (16)	C15—C13—P1—C8	171.33 (8)
C2—C7—C8—C9	-177.32 (10)	C14—C13—P1—C8	-65.44 (9)
C12—C7—C8—P1	-179.47 (9)	C16—C13—P1—C1	-50.42 (9)
C2—C7—C8—P1	2.18 (12)	C15—C13—P1—C1	68.00 (9)
C8—C9—C10—C11	-1.41 (17)	C14—C13—P1—C1	-168.77 (8)
C9—C10—C11—C12	1.01 (18)	C16—C13—P1—C17	-179.38 (7)
C7—C12—C11—C10	0.44 (18)	C15—C13—P1—C17	-60.96 (10)
C9—C8—P1—C1	175.63 (11)	C14—C13—P1—C17	62.27 (9)
C7—C8—P1—C1	-3.81 (8)	Pd1 <sup>i</sup> —C13—Pd1—C12	-178.700 (11)
C9—C8—P1—C17	-72.71 (12)	Pd1 <sup>i</sup> —C13—Pd1—C13 <sup>i</sup>	0.0

Symmetry code: (i)  $-x, -y, -z+2$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

Cg7 is the centroid of the Pd1,Cl3,Pd1',Cl3' ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C11—H11 $\cdots$ Cg7 <sup>ii</sup>	0.95	2.68	3.5952 (13)	138

Symmetry code: (ii)  $x+1, y, z-1$ .