

# *rac*-4-Chloro-2-[{2-[(3-chloro-6-hydroxy-2,4-dimethylbenzyl)(methyl)amino]propyl}(methyl)amino)methyl]-3,5-dimethylphenol

Augusto Rivera,<sup>a\*</sup> Dency José Pacheco,<sup>a</sup> Jaime Ríos-Motta,<sup>a</sup> Mauricio Maldonado<sup>a</sup> and Michael Bolte<sup>b</sup>

<sup>a</sup>Departamento de Química, Facultad de Ciencias, Universidad Nacional de Colombia, Sede Bogotá, Cra 30 No. 45-03, Bogotá, Código Postal 111321, Colombia, and <sup>b</sup>Institut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany  
Correspondence e-mail: ariverau@unal.edu.co


Received 7 September 2012; accepted 18 September 2012

Key indicators: single-crystal X-ray study;  $T = 173\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ; disorder in main residue;  $R$  factor = 0.064;  $wR$  factor = 0.159; data-to-parameter ratio = 14.3.

The title compound,  $C_{23}H_{32}Cl_2N_2O_2$ , a potential chiral ligand for coordination chemistry, was prepared by a two-step reaction. The molecule is located on a crystallographic centre of inversion. As a result, the methyl group bonded to the methylene group is disordered over two equally occupied positions, sharing the same site as the H atom of the chiral C atom. As a further consequence of the crystallographic centrosymmetry, the 1,2-diaminopropane unit adopts an antiperiplanar conformation and the two benzene rings are coplanar. The central chain is in an all-*trans* arrangement. An intramolecular O—H $\cdots$ N hydrogen bond makes an *S*(6) ring motif. A C—H $\cdots$  $\pi$  interaction links the molecules into one-dimensional chains along the [001] direction.

## Related literature

For the synthesis of the title compound, see: Rivera *et al.* (2010); Burke (1949). For the uses of tetrahydrosalens in coordination chemistry, see: Atwood (1997). For related structures, see: Rivera *et al.* (2011); Xu *et al.* (2009). For reference bond-length data, see: Allen *et al.* (1987). For graph-set analysis of hydrogen bonds, see: Bernstein *et al.* (1995).



## Experimental

### Crystal data

$C_{23}H_{32}Cl_2N_2O_2$	$V = 1129.19(19)\text{ \AA}^3$
$M_r = 439.41$	$Z = 2$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 9.5011(8)\text{ \AA}$	$\mu = 0.31\text{ mm}^{-1}$
$b = 11.9060(13)\text{ \AA}$	$T = 173\text{ K}$
$c = 9.9824(9)\text{ \AA}$	$0.25 \times 0.22 \times 0.08\text{ mm}$
$\beta = 90.348(7)^\circ$	

### Data collection

Stoe IPDS II two-circle diffractometer	11151 measured reflections
Absorption correction: multi-scan ( <i>X</i> -AREA; Stoe & Cie, 2001)	2055 independent reflections
$T_{\min} = 0.927$ , $T_{\max} = 0.976$	1758 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.074$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.159$	$\Delta\rho_{\text{max}} = 0.59\text{ e \AA}^{-3}$
$S = 1.14$	$\Delta\rho_{\text{min}} = -0.35\text{ e \AA}^{-3}$
2055 reflections	
144 parameters	

**Table 1**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$Cg1$  is the centroid of the C11–C16 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 $\cdots$ N1	0.82 (5)	1.87 (5)	2.614 (3)	150 (4)
C1—H1B $\cdots$ Cg1 <sup>i</sup>	0.99	2.83	3.709 (3)	148

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *X*-AREA (Stoe & Cie, 2001); cell refinement: *X*-AREA; data reduction: *X*-AREA; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Dirección de Investigaciones, Sede Bogotá (DIB) de la Universidad Nacional de Colombia, for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2069).

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Atwood, D. A. (1997). *Coord. Chem. Rev.* **165**, 267–296.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Burke, W. J. (1949). *J. Am. Chem. Soc.* **71**, 609–612.
- Rivera, A., Rojas, J. J., Ríos-Motta, J., Dušek, M. & Fejfarová, K. (2011). *Acta Cryst. E67*, o1391.
- Rivera, A., Rojas, J. J., Salazar-Barrios, J., Maldonado, M. & Ríos-Motta, J. (2010). *Molecules*, **15**, 4102–4110.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stoe & Cie (2001). *X*-AREA. Stoe & Cie, Darmstadt, Germany.
- Xu, Y.-M., Gao, S. & Ng, S. W. (2009). *Acta Cryst. E65*, o3151.

# supporting information

*Acta Cryst.* (2012). E68, o2997 [https://doi.org/10.1107/S1600536812039694]

## ***rac*-4-Chloro-2-[{2-[(3-chloro-6-hydroxy-2,4-dimethylbenzyl)(methyl)amino]-propyl}(methyl)amino)methyl]-3,5-dimethylphenol**

**Augusto Rivera, Dency José Pacheco, Jaime Ríos-Motta, Mauricio Maldonado and Michael Bolte**

### S1. Comment

Compound **I**,  $C_{23}H_{32}Cl_2N_2O_2$ , a new chiral *N,N*-dimethylated tetrahydrosalen ( $H_2[H4]$ salen) was obtained by reacting the bis-benzoxazine (**II**) with sodium borohydride using our procedure reported earlier (Rivera *et al.* 2010). The intermediate **II** was prepared by condensing 1,2-diaminopropane with formaldehyde and 4-chloro-3,5-dimethylphenol employing the general procedure of Burke (1949). The synthetic route for the title compound reported herein is illustrated in Fig. 1. The molecular structure and atom-numbering scheme for title compound,  $C_{23}H_{32}Cl_2N_2O_2$ , are shown in Fig. 2. The bond/newcifs lengths (Allen *et al.*, 1987) and angles are normal and similar to those observed for related structures (Rivera *et al.* 2010; Xu *et al.* 2009). In the title molecule, the 1,2-propanodiamine unit adopts an antiperiplanar conformation with an N1—C3—C3a—N1a torsion angle of -180.0 (2) $^{\circ}$ . As a consequence of this conformation, both benzene rings are parallel to each other. The central chain —CH<sub>2</sub>—N(CH<sub>3</sub>)—CH<sub>2</sub>—CH(CH<sub>3</sub>)—N(CH<sub>3</sub>)—CH<sub>2</sub>— is found in an all-*trans* arrangement. The two symmetry-related methyl substituents in the molecule (C2 and C2<sup>i</sup>, (i) = 1 - *x*, 1 - *y*, 2 - *z*) are orientated in an antiperiplanar arrangement (pseudo torsion angle CH<sub>3</sub>—N···N—CH<sub>3</sub> = 180.00 $^{\circ}$ ). The C2 and C4 methyl groups are almost (+)-synclinal [C2—N1—C3—C4 torsion angle = 48.0 (4) $^{\circ}$ ], a conformation stabilized by an intramolecular O—H···N hydrogen bond. The relationship of C4 methyl to C2<sup>i</sup> is defined by the pseudo torsion angle C4—C3···N1a—C2a, which is 114.64 (3) $^{\circ}$ .

The intramolecular O—H···N hydrogen bond (Table 1) makes an S(6) ring motif (Bernstein *et al.*, 1995), contrasting with the related structure (Xu *et al.*, 2009). In the crystal structure, intermolecular C1—H1B···Cg1(1 - *x*, 1 - *y*, 1 - *z*) interaction links the molecules into one-dimensional chains. C1···Cg1 is 3.709 (3) $\text{\AA}$ , H1···Cg1 is 2.83 $\text{\AA}$  and the angle at H1 is 148 $^{\circ}$ . Cg1 is the centroid of the C11—C16 ring.


### S2. Experimental

Sodium borohydride (3.0 mmol, 0.11 g) was added to a solution of 3,3'-(propane-1,2-diyl)bis(6-chloro-5,7-dimethyl-3,4-dihydro-2 h-benzo[*e*][1,3]oxazina) (435 mg, 1 mmol) (**II**) in ethanol (15 ml), and the mixture was stirred magnetically for 30 min at room temperature. After completion of the reaction, the mixture was poured into ice-cold water, neutralized with ammonium chloride (12 ml), and extracted with CHCl<sub>3</sub> (3 × 10 ml). The combined extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated. Recrystallization from ethanol afforded (**I**) in 91% yield. m.p. 424 K.

### S3. Refinement


All H atoms bonded to C were refined using a riding model with fixed individual displacement parameters [ $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ ] with C—H ranging from 0.95  $\text{\AA}$  to 0.99  $\text{\AA}$ . The hydroxyl H atom was isotropically refined. The methyl group C4 is disordered over two centrosymmetrically related positions each with a 50% occupancy.

The completeness of the data is 99.4% with eleven reflections missing for a full completeness. Since no reflection was omitted on purpose, this is most probably due to the data collection strategy using an area detector.



**Figure 1**

Synthetic route for the title compound.



**Figure 2**

A perspective view of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii. Only one of the two positions of the disordered methyl group is shown. Hydrogen bonds are drawn as dashed lines.

*rac*-4-Chloro-2-[{(2-[(3-chloro-6-hydroxy-2,4-dimethylbenzyl)(methyl)amino]propyl)(methyl)amino)methyl]-3,5-dimethylphenol

#### Crystal data



M<sub>r</sub> = 439.41

Monoclinic, P2<sub>1</sub>/n

Hall symbol: -P 2yn

a = 9.5011 (8) Å

b = 11.9060 (13) Å

c = 9.9824 (9) Å

β = 90.348 (7)°

V = 1129.19 (19) Å<sup>3</sup>

Z = 2

$$F(000) = 468$$

$$D_x = 1.292 \text{ Mg m}^{-3}$$

$$\text{Mo } K\alpha \text{ radiation, } \lambda = 0.71073 \text{ Å}$$

Cell parameters from 11821 reflections

$$\theta = 2.4\text{--}28.1^\circ$$

$$\mu = 0.31 \text{ mm}^{-1}$$

$$T = 173 \text{ K}$$

Plate, colourless

0.25 × 0.22 × 0.08 mm

*Data collection*

Stoe IPDS II two-circle  
 diffractometer  
 Radiation source: Genix 3D  $\text{I}\mu\text{S}$  microfocus X-ray source  
 Genix 3D multilayer optics monochromator  
 $\omega$  scans  
 Absorption correction: multi-scan (*X-AREA*; Stoe & Cie, 2001)  
 $T_{\min} = 0.927$ ,  $T_{\max} = 0.976$   
 11151 measured reflections  
 2055 independent reflections  
 1758 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.074$   
 $\theta_{\max} = 25.3^\circ$ ,  $\theta_{\min} = 3.4^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -14 \rightarrow 14$   
 $l = -11 \rightarrow 11$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.159$   
 $S = 1.14$   
 2055 reflections  
 144 parameters  
 0 restraints  
 Primary atom site location: structure-invariant direct methods  
 Secondary atom site location: difference Fourier map  
 Hydrogen site location: inferred from neighbouring sites  
 H atoms treated by a mixture of independent and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0629P)^2 + 0.952P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.59 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.**  $^1\text{H}$  NMR (400.1 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.07 (d,  $^3J = 6.6$  Hz, 3H), 2.18 (s, 3H), 2.25 (s, 3H), 2.30 (s, 6H), 2.31 (s, 6H), 2.35 (dd,  $^2J_{\text{gem}} = 12.6$ ,  $^3J = 6.9$  Hz, 1H), 2.62 (dd,  $^2J_{\text{gem}} = 12.6$ ,  $^3J = 6.7$  Hz, 1H), 3.03–3.11 (m, 1H), 3.69 (d,  $^2J_{\text{gem}} = 14.0$ , 1H), 3.74 (d,  $^2J_{\text{gem}} = 14.0$ , 1H), 3.81 (s, 2H), 6.62 (s, 2H).  $^{13}\text{C}$  NMR (100.6 MHz,  $\text{CDCl}_3$ ):  $\delta$  11.17, 16.74, 16.82, 21.22, 21.24, 35.19, 41.9, 53.68, 54.20, 58.57, 60.32, 116.57, 116.58, 118.48, 118.94, 125.26, 125.39, 133.96, 134.04, 136.62, 136.73, 156.76, 156.78. FT-IR (KBr) ( $\nu$ ,  $\text{cm}^{-1}$ ): 3406 (O—H, broad, m), 2960 ( $\text{CH}_3$  asym, st), 2922 ( $\text{CH}_2$  asym, st), 2855 ( $\text{CH}_3$  sym, st), 2802 ( $\text{CH}_2$  sym, st), 1613 (—C=C, st), 1320 (C—N, st), 668 (C—Cl, st).

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.74219 (10)	0.71021 (8)	0.25120 (9)	0.0590 (3)	
N1	0.5987 (3)	0.54354 (18)	0.8459 (2)	0.0383 (6)	
O1	0.7205 (2)	0.39166 (17)	0.6972 (3)	0.0485 (6)	
H1	0.687 (5)	0.421 (4)	0.764 (5)	0.071 (13)*	
C1	0.5450 (3)	0.5882 (3)	0.7189 (3)	0.0410 (7)	
H1A	0.5303	0.6702	0.7282	0.049*	
H1B	0.4525	0.5535	0.6992	0.049*	
C2	0.7218 (4)	0.6084 (3)	0.8905 (3)	0.0479 (8)	
H2A	0.7945	0.6057	0.8214	0.072*	

H2B	0.7588	0.5762	0.9739	0.072*	
H2C	0.6940	0.6866	0.9059	0.072*	
C3	0.4831 (3)	0.5398 (2)	0.9448 (3)	0.0464 (8)	
H3	0.3984	0.5110	0.8981	0.056*	
H3'	0.4654	0.6157	0.9814	0.056*	0.50
C4	0.4443 (7)	0.6608 (5)	1.0048 (6)	0.0403 (13)	0.50
H4A	0.5114	0.6802	1.0760	0.060*	0.50
H4B	0.3489	0.6588	1.0416	0.060*	0.50
H4C	0.4489	0.7172	0.9335	0.060*	0.50
C11	0.6426 (3)	0.5669 (2)	0.6023 (3)	0.0346 (6)	
C12	0.6469 (3)	0.6424 (2)	0.4950 (3)	0.0370 (6)	
C13	0.7364 (3)	0.6175 (2)	0.3882 (3)	0.0386 (7)	
C14	0.8205 (3)	0.5226 (2)	0.3842 (3)	0.0372 (7)	
C15	0.8119 (3)	0.4494 (2)	0.4901 (3)	0.0411 (7)	
H15	0.8676	0.3832	0.4896	0.049*	
C16	0.7241 (3)	0.4695 (2)	0.5979 (3)	0.0371 (6)	
C17	0.5559 (4)	0.7462 (3)	0.4936 (4)	0.0539 (9)	
H17A	0.6010	0.8052	0.5473	0.081*	
H17B	0.4636	0.7284	0.5314	0.081*	
H17C	0.5440	0.7725	0.4012	0.081*	
C18	0.9182 (3)	0.4993 (3)	0.2693 (3)	0.0506 (8)	
H18A	0.9715	0.4305	0.2874	0.076*	
H18B	0.9836	0.5624	0.2590	0.076*	
H18C	0.8632	0.4900	0.1867	0.076*	

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0693 (6)	0.0651 (6)	0.0429 (5)	0.0039 (4)	0.0184 (4)	0.0072 (4)
N1	0.0397 (13)	0.0333 (12)	0.0421 (14)	-0.0020 (9)	0.0198 (11)	0.0028 (10)
O1	0.0586 (14)	0.0358 (11)	0.0515 (14)	0.0070 (9)	0.0185 (11)	0.0011 (10)
C1	0.0323 (15)	0.0451 (16)	0.0457 (18)	0.0006 (12)	0.0128 (12)	0.0010 (13)
C2	0.0512 (19)	0.0565 (18)	0.0361 (17)	-0.0031 (15)	0.0128 (14)	-0.0041 (14)
C3	0.0454 (18)	0.0442 (16)	0.0499 (19)	0.0062 (13)	0.0217 (14)	0.0130 (14)
C4	0.049 (3)	0.040 (3)	0.032 (3)	0.007 (2)	0.015 (3)	0.008 (2)
C11	0.0263 (13)	0.0379 (14)	0.0396 (16)	-0.0048 (10)	0.0067 (11)	-0.0064 (11)
C12	0.0332 (14)	0.0396 (15)	0.0383 (16)	-0.0025 (11)	0.0055 (12)	-0.0041 (12)
C13	0.0372 (15)	0.0436 (15)	0.0350 (16)	-0.0044 (12)	0.0060 (12)	-0.0047 (12)
C14	0.0300 (14)	0.0469 (16)	0.0348 (15)	-0.0060 (12)	0.0059 (11)	-0.0117 (12)
C15	0.0362 (15)	0.0422 (15)	0.0448 (17)	0.0034 (12)	0.0040 (13)	-0.0134 (13)
C16	0.0359 (15)	0.0342 (14)	0.0413 (16)	-0.0045 (11)	0.0069 (12)	-0.0036 (12)
C17	0.061 (2)	0.0508 (18)	0.050 (2)	0.0165 (15)	0.0152 (16)	0.0071 (15)
C18	0.0440 (17)	0.067 (2)	0.0415 (18)	0.0017 (15)	0.0125 (14)	-0.0143 (15)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Cl1—C13	1.758 (3)	C4—H4B	0.9800
N1—C1	1.464 (4)	C4—H4C	0.9800

N1—C2	1.468 (4)	C11—C16	1.396 (4)
N1—C3	1.483 (4)	C11—C12	1.399 (4)
O1—C16	1.358 (4)	C12—C13	1.400 (4)
O1—H1	0.82 (5)	C12—C17	1.508 (4)
C1—C11	1.514 (4)	C13—C14	1.385 (4)
C1—H1A	0.9900	C14—C15	1.373 (4)
C1—H1B	0.9900	C14—C18	1.506 (4)
C2—H2A	0.9800	C15—C16	1.386 (4)
C2—H2B	0.9800	C15—H15	0.9500
C2—H2C	0.9800	C17—H17A	0.9800
C3—C3 <sup>i</sup>	1.486 (6)	C17—H17B	0.9800
C3—C4	1.604 (6)	C17—H17C	0.9800
C3—H3	0.9886	C18—H18A	0.9800
C3—H3'	0.9905	C18—H18B	0.9800
C4—H3'	0.6186	C18—H18C	0.9800
C4—H4A	0.9800		
C1—N1—C2	110.1 (2)	C16—C11—C12	119.5 (3)
C1—N1—C3	109.4 (2)	C16—C11—C1	120.4 (3)
C2—N1—C3	113.9 (2)	C12—C11—C1	120.1 (2)
C16—O1—H1	108 (3)	C11—C12—C13	117.9 (3)
N1—C1—C11	113.1 (2)	C11—C12—C17	121.0 (3)
N1—C1—H1A	108.9	C13—C12—C17	121.1 (3)
C11—C1—H1A	108.9	C14—C13—C12	123.3 (3)
N1—C1—H1B	108.9	C14—C13—Cl1	117.9 (2)
C11—C1—H1B	108.9	C12—C13—Cl1	118.8 (2)
H1A—C1—H1B	107.8	C15—C14—C13	117.3 (3)
N1—C2—H2A	109.5	C15—C14—C18	120.7 (3)
N1—C2—H2B	109.5	C13—C14—C18	122.1 (3)
H2A—C2—H2B	109.5	C14—C15—C16	121.9 (3)
N1—C2—H2C	109.5	C14—C15—H15	119.1
H2A—C2—H2C	109.5	C16—C15—H15	119.1
H2B—C2—H2C	109.5	O1—C16—C15	117.9 (3)
N1—C3—C3 <sup>i</sup>	110.8 (3)	O1—C16—C11	121.9 (3)
N1—C3—C4	113.2 (3)	C15—C16—C11	120.3 (3)
C3 <sup>i</sup> —C3—C4	110.2 (4)	C12—C17—H17A	109.5
N1—C3—H3	107.5	C12—C17—H17B	109.5
C3 <sup>i</sup> —C3—H3	107.5	H17A—C17—H17B	109.5
C4—C3—H3	107.4	C12—C17—H17C	109.5
N1—C3—H3'	110.3	H17A—C17—H17C	109.5
C3 <sup>i</sup> —C3—H3'	110.1	H17B—C17—H17C	109.5
H3—C3—H3'	110.5	C14—C18—H18A	109.5
C3—C4—H4A	109.5	C14—C18—H18B	109.5
C3—C4—H4B	109.5	H18A—C18—H18B	109.5
H4A—C4—H4B	109.5	C14—C18—H18C	109.5
C3—C4—H4C	109.5	H18A—C18—H18C	109.5
H4A—C4—H4C	109.5	H18B—C18—H18C	109.5
H4B—C4—H4C	109.5		

C2—N1—C1—C11	69.4 (3)	C11—C12—C13—Cl1	179.3 (2)
C3—N1—C1—C11	−164.7 (2)	C17—C12—C13—Cl1	0.3 (4)
C1—N1—C3—C3 <sup>i</sup>	159.9 (3)	C12—C13—C14—C15	1.2 (4)
C2—N1—C3—C3 <sup>i</sup>	−76.3 (4)	Cl1—C13—C14—C15	−178.2 (2)
C1—N1—C3—C4	−75.7 (4)	C12—C13—C14—C18	−178.6 (3)
C2—N1—C3—C4	48.0 (4)	Cl1—C13—C14—C18	2.0 (4)
N1—C1—C11—C16	33.1 (4)	C13—C14—C15—C16	−0.7 (4)
N1—C1—C11—C12	−149.8 (3)	C18—C14—C15—C16	179.1 (3)
N1—C3—C3 <sup>i</sup> —N1 <sup>i</sup>	−180.0 (2)	C14—C15—C16—O1	178.9 (3)
C16—C11—C12—C13	−1.4 (4)	C14—C15—C16—C11	−0.9 (4)
C1—C11—C12—C13	−178.5 (3)	C12—C11—C16—O1	−177.8 (3)
C16—C11—C12—C17	177.5 (3)	C1—C11—C16—O1	−0.7 (4)
C1—C11—C12—C17	0.4 (4)	C12—C11—C16—C15	2.0 (4)
C11—C12—C13—C14	−0.2 (4)	C1—C11—C16—C15	179.1 (3)
C17—C12—C13—C14	−179.1 (3)		

Symmetry code: (i)  $-x+1, -y+1, -z+2$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^{\circ}$ )

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1—H1 $\cdots$ N1	0.82 (5)	1.87 (5)	2.614 (3)	150 (4)
C1—H1B $\cdots$ Cg1 <sup>ii</sup>	0.99	2.83	3.709 (3)	148

Symmetry code: (ii)  $-x+1, -y+1, -z+1$ .