

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-(2-Fluorobenzoyl)-1-[2-(4-hydroxyphenyl)-2-oxoethyl]piperazin-1-ium trifluoroacetate

Fuyong Bian, Yi Jin, Shaoming Chi, Guojun Shi and Sichuan Xu*

College of Chemical Science and Technology, Yunnan University, Kunming City, People's Republic of China

Correspondence e-mail: sichuan@ynu.edu.cn


Received 5 September 2012; accepted 13 October 2012

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.073; wR factor = 0.240; data-to-parameter ratio = 13.2.

In the crystal structure of the title compound, $\text{C}_{19}\text{H}_{20}\text{FN}_2\text{O}_3^+ \cdot \text{C}_2\text{F}_3\text{O}_2^-$, $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link two cations and two anions into a 22-atom ring. These rings are further linked into a three dimensional network by weak $\text{C}-\text{H} \cdots \text{O}$ contacts.

Related literature

For the preparation for the title compound, see: Hoff *et al.* (2005); Wallén *et al.* (2003); Stachulski *et al.* (2006). For similar structures, see: Luedtkea & Mach (2003); Rok *et al.* (2007); Friedel & Crafts (1932*a,b*). For the applications of similar compounds, see: Wise (1996).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{20}\text{FN}_2\text{O}_3^+ \cdot \text{C}_2\text{F}_3\text{O}_2^-$
 $M_r = 456.39$
 Monoclinic, $P2_1/n$
 $a = 10.055$ (3) Å
 $b = 9.601$ (2) Å
 $c = 21.946$ (5) Å

 $\beta = 91.960$ (4)°
 $V = 2117.4$ (9) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

 $\mu = 0.13$ mm⁻¹
 $T = 298$ K
 $0.23 \times 0.20 \times 0.18$ mm

Data collection

 Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.956$, $T_{\max} = 0.971$
 12540 measured reflections
 3824 independent reflections
 1534 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.070$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.240$
 $S = 1.03$
 3824 reflections
 289 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N1}-\text{H1N} \cdots \text{O5}^{\text{I}}$	0.91	1.87	2.710 (5)	152
$\text{O1}-\text{H1} \cdots \text{O5}^{\text{II}}$	0.82	1.95	2.697 (5)	151
$\text{C8}-\text{H8B} \cdots \text{O3}^{\text{I}}$	0.97	2.22	2.996 (6)	136

 Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Support from the National Natural Science Foundation of China (21163024) is acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: QM2083).

References

- Bruker (2001). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Friedel, C. & Crafts, J. M. (1932*a*). *Compt. Rend.* **84**, 1392–1395.
- Friedel, C. & Crafts, J. M. (1932*b*). *Compt. Rend.* **84**, 1450–1450.
- Hoff, B., Strandberg, E., Ulrich, A. S., Tieleman, D. P. & Posten, C. (2005). *Biophys. J.* **88**, 1818–1827.
- Luedtkea, R. R. & Mach, R. H. (2003). *Curr. Pharm. Des.* **9**, 643–671.
- Rok, F., Stanislav, G. & Danijel, K. (2007). *Tetrahedron*, **63**, 10698–10708.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stachulski, A. V., Berry, N. G., Low, A. C. L., Moores, S. L., Row, E., Warhurst, D. C., Adagu, I. S. & Rossignol, J.-F. (2006). *J. Med. Chem.* **49**, 1450–1454.
- Wallén, E. A. A., Christiaans, J. A. M., Jarho, E. M., Forsberg, M. M., Venäläinen, J. I., Mannisto, P. T. & Gynther, J. (2003). *J. Med. Chem.* **46**, 4543–4551.
- Wise, R. A. (1996). *Curr Opin Neurobiol.* **6**, 243–251.

supporting information

Acta Cryst. (2012). E68, o3243 [doi:10.1107/S1600536812042857]

4-(2-Fluorobenzoyl)-1-[2-(4-hydroxyphenyl)-2-oxoethyl]piperazin-1-ium tri-fluoroacetate


Fuyong Bian, Yi Jin, Shaoming Chi, Guojun Shi and Sichuan Xu

S1. Experimental

The title compound was obtained through the following four steps: 2-fluorobenzoic acid (700 mg, 5 mmol) was dissolved into 3 ml of SOCl_2 , the mixture stirred at 1033 K for 5 h and then slowly cooled to room temperature before removing excess SOCl_2 . After preparing an ethanol solution (10 ml) of 2-fluorobenzoyl chloride (317 mg, 2 mmol) and 1-boc-piperazine (372 mg, 4 mmol), potassium carbonate (552 mg, 4 mmol) was added to the solution. The mixture was refluxed for 12 h and then cooled to room temperature; extraction and concentration were performed to obtain an oil. After mixing the oil (a dichloromethane solution, 10 ml) with TFA (1.5 ml) and stirring for 12 h to complete the reaction, the mixture was poured into water. The aqueous layer was extracted by ethyl acetate, and then the organic layer was separated, dried using sodium sulfate to obtain 4-(2-fluorobenzoyl)piperazin-1-ium trifluoroacetate. Potassium carbonate (276 mg, 2 mmol) was added to the prepared ethanol solution (10 ml) of (2-fluorophenyl)(piperazin-1-yl)methanone (208 mg, 1 mmol) and 2-chloro-1-(4-hydroxyphenyl)ethanone (341 mg, 2 mmol), which was refluxed for 5 h and mixed by adding water. Again the aqueous layer was extracted by ethyl acetate for the separation of the organic layer, washed using anhydrous ethyl acetate, and dried using sodium sulfate, filtered, and concentrated in vacuum. The purification of residue by using silica gel column chromatography and eluting with EtOAc–petroleum ether (1:1) produced the pale-yellow solid (yield: 164 mg, 48%) of 4-(2-fluorobenzoyl)-1-[2-(4-hydroxyphenyl)-2-oxoethyl]piperazin-1-ium trifluoroacetate.

S2. Refinement

H atoms were placed in calculated positions [$\text{C—H} = 0.93$ (aromatic) and $\text{O—H} = 0.82$ or 0.96 \AA (methyl group)] and refined using a riding model approximation with $U_{\text{iso}}(\text{H})$ constrained to 1.2 (aromatic) or 1.5 (methyl, O—H) times U_{eq} of the respective parent atom.

**Figure 1**

Ellipsoid plot

4-(2-Fluorobenzoyl)-1-[2-(4-hydroxyphenyl)-2-oxoethyl]piperazin-1-ium trifluoroacetate*Crystal data* $C_{19}H_{20}FN_2O_3^+ \cdot C_2F_3O_2^-$ $M_r = 456.39$ Monoclinic, $P2_1/n$ $a = 10.055 (3) \text{ \AA}$ $b = 9.601 (2) \text{ \AA}$ $c = 21.946 (5) \text{ \AA}$ $\beta = 91.960 (4)^\circ$ $V = 2117.4 (9) \text{ \AA}^3$ $Z = 4$ $F(000) = 944$ $D_x = 1.432 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 892 reflections

 $\theta = 2.3\text{--}17.0^\circ$ $\mu = 0.13 \text{ mm}^{-1}$ $T = 298 \text{ K}$

Parallelepiped, colourless

 $0.23 \times 0.20 \times 0.18 \text{ mm}$ *Data collection*Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ϕ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2001)

 $T_{\min} = 0.956$, $T_{\max} = 0.971$

12540 measured reflections

3824 independent reflections

1534 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.070$ $\theta_{\max} = 25.4^\circ$, $\theta_{\min} = 1.9^\circ$ $h = -12 \rightarrow 12$ $k = 0 \rightarrow 11$ $l = 0 \rightarrow 16$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.073$ $wR(F^2) = 0.240$ $S = 1.03$

3824 reflections

289 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.6641 (4)	0.7921 (4)	0.35902 (18)	0.1420 (13)
F2	-0.0352 (4)	0.0629 (4)	0.3531 (2)	0.1733 (18)
F3	-0.0010 (4)	0.2786 (4)	0.35550 (19)	0.1465 (15)
F4	-0.0132 (4)	0.1720 (4)	0.27235 (19)	0.1558 (16)
N1	0.9794 (3)	0.7317 (3)	0.17362 (14)	0.0623 (9)
H1N	1.0634	0.7032	0.1657	0.081*
N2	0.9521 (4)	0.6187 (4)	0.29247 (17)	0.0772 (10)
O1	0.7573 (3)	0.9834 (3)	-0.15804 (14)	0.0945 (11)
H1	0.7304	1.0632	-0.1533	0.123*
O2	1.0042 (3)	0.6279 (3)	0.05966 (14)	0.0850 (9)
O3	0.7802 (4)	0.5059 (4)	0.33367 (18)	0.1197 (13)
O4	0.2142 (4)	0.0147 (4)	0.3220 (2)	0.1226 (14)
O5	0.2540 (3)	0.2358 (4)	0.34091 (16)	0.0951 (11)
C1	0.7985 (4)	0.9296 (5)	-0.1033 (2)	0.0742 (12)
C2	0.8442 (5)	0.7938 (5)	-0.1037 (2)	0.0813 (13)
H2	0.8444	0.7442	-0.1401	0.106*
C3	0.8891 (4)	0.7321 (4)	-0.0505 (2)	0.0753 (12)
H3	0.9200	0.6409	-0.0512	0.098*
C4	0.8891 (4)	0.8047 (4)	0.00491 (19)	0.0642 (11)
C5	0.8382 (4)	0.9385 (4)	0.0040 (2)	0.0720 (12)
H5	0.8336	0.9874	0.0405	0.094*
C6	0.7943 (4)	1.0014 (5)	-0.0493 (2)	0.0766 (12)
H6	0.7618	1.0921	-0.0488	0.100*
C7	0.9441 (4)	0.7384 (4)	0.06037 (19)	0.0653 (11)
C8	0.9287 (5)	0.8139 (4)	0.12025 (18)	0.0711 (12)
H8A	0.9763	0.9016	0.1190	0.092*
H8B	0.8353	0.8346	0.1253	0.092*
C9	0.8978 (5)	0.6055 (4)	0.1846 (2)	0.0846 (14)
H9A	0.8992	0.5449	0.1493	0.110*
H9B	0.8063	0.6325	0.1908	0.110*
C10	0.9516 (5)	0.5284 (4)	0.2399 (2)	0.0872 (14)
H10A	0.8969	0.4473	0.2472	0.113*
H10B	1.0415	0.4968	0.2329	0.113*
C11	1.0351 (5)	0.7411 (4)	0.28498 (18)	0.0728 (12)
H11A	1.1268	0.7127	0.2803	0.095*

H11B	1.0314	0.7994	0.3210	0.095*
C12	0.9874 (4)	0.8226 (4)	0.22957 (18)	0.0684 (11)
H12A	0.9003	0.8614	0.2368	0.089*
H12B	1.0480	0.8993	0.2228	0.089*
C13	0.8623 (5)	0.6010 (6)	0.3364 (2)	0.0864 (14)
C14	0.8670 (5)	0.6952 (5)	0.3901 (2)	0.0782 (13)
C15	0.7657 (6)	0.7849 (6)	0.4023 (3)	0.0935 (15)
C16	0.7596 (7)	0.8661 (6)	0.4533 (3)	0.1098 (19)
H16	0.6881	0.9251	0.4598	0.143*
C17	0.8654 (8)	0.8552 (7)	0.4944 (3)	0.117 (2)
H17	0.8663	0.9095	0.5295	0.152*
C18	0.9690 (7)	0.7671 (8)	0.4851 (3)	0.133 (2)
H18	1.0400	0.7609	0.5133	0.173*
C19	0.9659 (6)	0.6875 (7)	0.4329 (3)	0.124 (2)
H19	1.0354	0.6255	0.4269	0.161*
C20	0.1836 (5)	0.1356 (6)	0.3310 (2)	0.0810 (14)
C21	0.0321 (5)	0.1635 (6)	0.3290 (3)	0.0932 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.138 (3)	0.151 (3)	0.135 (3)	0.049 (2)	-0.022 (2)	0.000 (2)
F2	0.103 (3)	0.157 (3)	0.262 (5)	-0.037 (2)	0.036 (3)	0.053 (3)
F3	0.120 (3)	0.135 (3)	0.185 (4)	0.039 (2)	0.010 (3)	-0.046 (3)
F4	0.137 (3)	0.195 (4)	0.131 (3)	0.028 (3)	-0.059 (3)	-0.024 (3)
N1	0.068 (2)	0.0591 (19)	0.060 (2)	-0.0027 (16)	0.0064 (16)	0.0012 (15)
N2	0.089 (3)	0.071 (2)	0.073 (2)	-0.011 (2)	0.016 (2)	0.0063 (19)
O1	0.112 (3)	0.094 (2)	0.076 (2)	-0.0032 (18)	-0.0173 (19)	0.0037 (17)
O2	0.099 (2)	0.075 (2)	0.081 (2)	0.0236 (17)	0.0098 (17)	-0.0025 (16)
O3	0.113 (3)	0.121 (3)	0.127 (3)	-0.045 (2)	0.034 (2)	0.004 (2)
O4	0.106 (3)	0.093 (3)	0.169 (4)	0.002 (2)	0.004 (3)	-0.007 (3)
O5	0.088 (2)	0.093 (2)	0.104 (3)	-0.0177 (19)	0.0044 (19)	-0.0028 (19)
C1	0.072 (3)	0.071 (3)	0.079 (3)	-0.004 (2)	-0.009 (2)	0.008 (2)
C2	0.092 (3)	0.086 (3)	0.066 (3)	0.002 (3)	-0.008 (3)	-0.006 (2)
C3	0.078 (3)	0.067 (3)	0.082 (3)	-0.001 (2)	0.013 (2)	-0.009 (2)
C4	0.067 (3)	0.059 (2)	0.067 (3)	-0.003 (2)	0.003 (2)	-0.011 (2)
C5	0.079 (3)	0.064 (3)	0.072 (3)	0.000 (2)	0.001 (2)	-0.009 (2)
C6	0.076 (3)	0.073 (3)	0.080 (3)	-0.001 (2)	-0.002 (2)	0.000 (2)
C7	0.065 (3)	0.060 (3)	0.072 (3)	-0.005 (2)	0.011 (2)	-0.005 (2)
C8	0.087 (3)	0.063 (2)	0.063 (3)	0.012 (2)	-0.001 (2)	-0.003 (2)
C9	0.097 (3)	0.068 (3)	0.088 (3)	-0.030 (3)	0.010 (3)	-0.002 (2)
C10	0.104 (4)	0.062 (3)	0.096 (4)	-0.016 (2)	0.017 (3)	0.010 (3)
C11	0.081 (3)	0.077 (3)	0.061 (3)	-0.016 (2)	0.001 (2)	0.010 (2)
C12	0.077 (3)	0.064 (2)	0.064 (3)	-0.004 (2)	0.004 (2)	-0.004 (2)
C13	0.083 (3)	0.085 (3)	0.092 (4)	-0.005 (3)	0.012 (3)	0.019 (3)
C14	0.068 (3)	0.095 (3)	0.073 (3)	-0.007 (3)	0.013 (2)	0.026 (3)
C15	0.090 (4)	0.099 (4)	0.091 (4)	0.010 (3)	0.003 (3)	0.029 (3)
C16	0.148 (6)	0.087 (4)	0.097 (4)	0.010 (4)	0.034 (4)	0.014 (3)

C17	0.140 (6)	0.142 (6)	0.070 (4)	-0.021 (5)	0.015 (4)	0.003 (3)
C18	0.095 (5)	0.203 (7)	0.102 (5)	0.026 (5)	0.004 (4)	0.006 (5)
C19	0.095 (4)	0.185 (6)	0.092 (4)	0.025 (4)	0.014 (4)	-0.026 (4)
C20	0.105 (4)	0.070 (3)	0.067 (3)	-0.012 (3)	0.000 (3)	0.003 (2)
C21	0.078 (3)	0.098 (4)	0.104 (4)	0.001 (3)	-0.001 (3)	0.000 (3)

Geometric parameters (Å, °)

F1—C15	1.372 (6)	C6—H6	0.9300
F2—C21	1.301 (6)	C7—C8	1.513 (5)
F3—C21	1.298 (6)	C8—H8A	0.9700
F4—C21	1.312 (6)	C8—H8B	0.9700
N1—C8	1.488 (5)	C9—C10	1.505 (6)
N1—C9	1.488 (5)	C9—H9A	0.9700
N1—C12	1.506 (5)	C9—H9B	0.9700
N1—H1N	0.9100	C10—H10A	0.9700
N2—C13	1.353 (6)	C10—H10B	0.9700
N2—C10	1.443 (5)	C11—C12	1.510 (5)
N2—C11	1.454 (5)	C11—H11A	0.9700
O1—C1	1.360 (5)	C11—H11B	0.9700
O1—H1	0.8200	C12—H12A	0.9700
O2—C7	1.221 (4)	C12—H12B	0.9700
O3—C13	1.231 (5)	C13—C14	1.486 (7)
O4—C20	1.219 (5)	C14—C19	1.346 (7)
O5—C20	1.209 (5)	C14—C15	1.368 (7)
C1—C6	1.373 (6)	C15—C16	1.367 (7)
C1—C2	1.382 (6)	C16—C17	1.375 (8)
C2—C3	1.372 (6)	C16—H16	0.9300
C2—H2	0.9300	C17—C18	1.363 (8)
C3—C4	1.402 (6)	C17—H17	0.9300
C3—H3	0.9300	C18—C19	1.377 (8)
C4—C5	1.383 (5)	C18—H18	0.9300
C4—C7	1.465 (6)	C19—H19	0.9300
C5—C6	1.376 (6)	C20—C21	1.546 (7)
C5—H5	0.9300		
C8—N1—C9	112.6 (3)	N2—C10—H10B	109.7
C8—N1—C12	110.0 (3)	C9—C10—H10B	109.7
C9—N1—C12	110.7 (3)	H10A—C10—H10B	108.2
C8—N1—H1N	107.8	N2—C11—C12	110.0 (3)
C9—N1—H1N	107.8	N2—C11—H11A	109.7
C12—N1—H1N	107.8	C12—C11—H11A	109.7
C13—N2—C10	120.7 (4)	N2—C11—H11B	109.7
C13—N2—C11	125.5 (4)	C12—C11—H11B	109.7
C10—N2—C11	112.4 (3)	H11A—C11—H11B	108.2
C1—O1—H1	109.5	N1—C12—C11	111.3 (3)
O1—C1—C6	123.7 (4)	N1—C12—H12A	109.4
O1—C1—C2	116.3 (4)	C11—C12—H12A	109.4

C6—C1—C2	120.0 (4)	N1—C12—H12B	109.4
C3—C2—C1	120.1 (4)	C11—C12—H12B	109.4
C3—C2—H2	120.0	H12A—C12—H12B	108.0
C1—C2—H2	120.0	O3—C13—N2	121.4 (5)
C2—C3—C4	121.0 (4)	O3—C13—C14	119.5 (5)
C2—C3—H3	119.5	N2—C13—C14	119.0 (5)
C4—C3—H3	119.5	C19—C14—C15	115.9 (5)
C5—C4—C3	117.4 (4)	C19—C14—C13	121.5 (5)
C5—C4—C7	123.0 (4)	C15—C14—C13	122.3 (5)
C3—C4—C7	119.6 (4)	C16—C15—C14	125.0 (6)
C6—C5—C4	121.9 (4)	C16—C15—F1	119.0 (6)
C6—C5—H5	119.1	C14—C15—F1	116.0 (5)
C4—C5—H5	119.1	C15—C16—C17	116.0 (6)
C1—C6—C5	119.7 (4)	C15—C16—H16	122.0
C1—C6—H6	120.2	C17—C16—H16	122.0
C5—C6—H6	120.2	C18—C17—C16	121.8 (6)
O2—C7—C4	122.7 (4)	C18—C17—H17	119.1
O2—C7—C8	119.5 (4)	C16—C17—H17	119.1
C4—C7—C8	117.8 (4)	C17—C18—C19	118.2 (6)
N1—C8—C7	112.8 (3)	C17—C18—H18	120.9
N1—C8—H8A	109.0	C19—C18—H18	120.9
C7—C8—H8A	109.0	C14—C19—C18	123.0 (6)
N1—C8—H8B	109.0	C14—C19—H19	118.5
C7—C8—H8B	109.0	C18—C19—H19	118.5
H8A—C8—H8B	107.8	O5—C20—O4	129.6 (6)
N1—C9—C10	110.3 (4)	O5—C20—C21	116.0 (5)
N1—C9—H9A	109.6	O4—C20—C21	114.5 (5)
C10—C9—H9A	109.6	F3—C21—F2	107.8 (5)
N1—C9—H9B	109.6	F3—C21—F4	106.6 (5)
C10—C9—H9B	109.6	F2—C21—F4	105.3 (5)
H9A—C9—H9B	108.1	F3—C21—C20	113.7 (5)
N2—C10—C9	109.9 (4)	F2—C21—C20	112.7 (5)
N2—C10—H10A	109.7	F4—C21—C20	110.3 (5)
C9—C10—H10A	109.7		
O1—C1—C2—C3	179.2 (4)	N2—C11—C12—N1	-54.0 (5)
C6—C1—C2—C3	-2.0 (7)	C10—N2—C13—O3	1.8 (7)
C1—C2—C3—C4	0.3 (7)	C11—N2—C13—O3	167.4 (4)
C2—C3—C4—C5	2.1 (6)	C10—N2—C13—C14	179.3 (4)
C2—C3—C4—C7	-176.9 (4)	C11—N2—C13—C14	-15.1 (7)
C3—C4—C5—C6	-2.8 (6)	O3—C13—C14—C19	106.7 (6)
C7—C4—C5—C6	176.1 (4)	N2—C13—C14—C19	-70.8 (6)
O1—C1—C6—C5	-180.0 (4)	O3—C13—C14—C15	-66.8 (7)
C2—C1—C6—C5	1.4 (7)	N2—C13—C14—C15	115.7 (5)
C4—C5—C6—C1	1.1 (7)	C19—C14—C15—C16	0.8 (8)
C5—C4—C7—O2	-170.3 (4)	C13—C14—C15—C16	174.7 (5)
C3—C4—C7—O2	8.6 (6)	C19—C14—C15—F1	-179.4 (5)
C5—C4—C7—C8	7.6 (6)	C13—C14—C15—F1	-5.6 (7)

C3—C4—C7—C8	-173.5 (4)	C14—C15—C16—C17	0.6 (8)
C9—N1—C8—C7	-67.6 (4)	F1—C15—C16—C17	-179.1 (5)
C12—N1—C8—C7	168.4 (3)	C15—C16—C17—C18	-1.0 (9)
O2—C7—C8—N1	-6.2 (6)	C16—C17—C18—C19	0.0 (10)
C4—C7—C8—N1	175.8 (3)	C15—C14—C19—C18	-1.9 (9)
C8—N1—C9—C10	-178.6 (4)	C13—C14—C19—C18	-175.8 (6)
C12—N1—C9—C10	-55.1 (5)	C17—C18—C19—C14	1.6 (11)
C13—N2—C10—C9	106.7 (5)	O5—C20—C21—F3	-19.3 (7)
C11—N2—C10—C9	-60.7 (5)	O4—C20—C21—F3	160.9 (5)
N1—C9—C10—N2	58.3 (5)	O5—C20—C21—F2	-142.4 (5)
C13—N2—C11—C12	-108.2 (5)	O4—C20—C21—F2	37.8 (7)
C10—N2—C11—C12	58.4 (5)	O5—C20—C21—F4	100.3 (5)
C8—N1—C12—C11	178.4 (3)	O4—C20—C21—F4	-79.5 (6)
C9—N1—C12—C11	53.4 (5)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N \cdots O5 ⁱ	0.91	1.87	2.710 (5)	152
O1—H1 \cdots O5 ⁱⁱ	0.82	1.95	2.697 (5)	151
C8—H8B \cdots O3 ⁱ	0.97	2.22	2.996 (6)	136

Symmetry codes: (i) $-x+3/2, y+1/2, -z+1/2$; (ii) $x+1/2, -y+3/2, z-1/2$.