

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-[4-Acetyl-1,3-phenylenebis(oxy)]-diacetic acid

Jian-Guo Wang,* Ping Yan, Chan-juan Zhong and Guo-zhen Yu

School of Chemistry & Environmental Engineering, Jiujiang University, Jiujiang 332005, People's Republic of China


Correspondence e-mail: jgwang117@163.com

Received 9 November 2012; accepted 21 November 2012

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.120; data-to-parameter ratio = 11.6.

In the title compound, $\text{C}_{12}\text{H}_{12}\text{O}_7$, the dihedral angles between the benzene ring and the mean planes of the 3-carboxymethoxy, 1-carboxymethoxy and acetyl substituents are 8.67 (7), 7.81 (6) and 10.3 (18) $^\circ$, respectively. In the crystal, molecules are linked by typical carboxylic acid $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a zigzag chain. $\text{C}-\text{H}\cdots\text{O}$ interactions also occur.

Related literature

For a related structure, see: Zhang *et al.* (2007).

Experimental

Crystal data

 $\text{C}_{12}\text{H}_{12}\text{O}_7$
 $M_r = 268.22$
 Triclinic, $P\bar{1}$
 $a = 5.1351$ (6) Å

 $b = 7.8346$ (9) Å
 $c = 15.6157$ (18) Å
 $\alpha = 86.217$ (2) $^\circ$
 $\beta = 81.321$ (1) $^\circ$
 $\gamma = 72.101$ (2) $^\circ$
 $V = 590.86$ (12) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

 $\mu = 0.13$ mm⁻¹
 $T = 298$ K
 $0.12 \times 0.10 \times 0.10$ mm

Data collection

 Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.985$, $T_{\max} = 0.987$

 3902 measured reflections
 2049 independent reflections
 1860 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.120$
 $S = 1.05$
 2049 reflections

 176 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³
Table 1
 Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.82	1.83	2.6530 (14)	180
$\text{O6}-\text{H6}\cdots\text{O5}^{\text{ii}}$	0.82	1.83	2.6352 (15)	167
$\text{C2}-\text{H2}\cdots\text{O7}^{\text{iii}}$	0.93	2.44	3.3566 (17)	168
$\text{C4}-\text{H4}\cdots\text{O6}^{\text{iv}}$	0.93	2.52	3.4392 (19)	169

 Symmetry codes: (i) $-x+3, -y+2, -z$; (ii) $-x-1, -y+2, -z+1$; (iii) $x-1, y+1, z$; (iv) $-x, -y+1, -z+1$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

This study was funded by Jiangxi Provincial Department of Education (GJJ08433) and the Doctoral Research Initiating Project of Jiujiang University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2075).

References

- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2003). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Zhang, C., Li, H., Liu, D. & Liu, M. (2007). *Acta Cryst.* **E63**, o4210.

supporting information

Acta Cryst. (2012). E68, o3470 [doi:10.1107/S1600536812047897]

2,2'-[4-Acetyl-1,3-phenylenebis(oxy)]diacetic acid

Jian-Guo Wang, Ping Yan, Chan-juan Zhong and Guo-zhen Yu

S1. Comment

The title compound is a potential inhibitor of mushroom tyrosinase.

The dihedral angles between the mean planes of the benzene ring and those of the 2-carboxymethoxyl,(O1-C8-C7-O3-O2), 4-carboxymethoxyl (O6-C10-C9-O4-O5), and acetyl, (C11-C12-O7), substituents are 8.67 (7)°, 7.81 (6)°, and 10.3 (18)° respectively.

In the crystal, the molecules are linked by typical carboxylic acid O—H···O hydrogen bonding forming a one-dimensional zig-zag chain Table 1, Figure 2.


This chain is linked to anti-parallel chains by C—H···O weak hydrogen bonds to form a two dimensional sheet stabilizing the supramolecular structure, Table 1, Figure 2.

S2. Experimental


2,4-dihydroxyacetophenone (10.64 g, 0.07 mol) and potassium hydroxide (10.58 g, 0.189 mol) were dissolved in dry acetone (100 ml) in a three-neck flask. Then ethyl bromoacetate (28.06 g, 0.168 mol) was dropwise added at room temperature and vigorously stirred for 3 h. The progress of the reaction was monitored by TCL (Si gel, developing solvent $V(\text{acetone})/V(\text{petroleum ether}) = 1:2$). After suction filtration and distillation to remove the solvent, a white solid was obtained, 19.76 g, yield 87.1%. This solid was dissolved in acetone (30 ml) and 20% aq. NaOH (50 ml) was added. The reaction mixture was stirred at 323 K for 1.5 h. After acidifying the mixture with dilute hydrochloric acid to pH=3, followed by suction filtration and washing the residue with water, the target product was prepared. After recrystallization from ethanol, crystalline colorless needles were obtained.

S3. Refinement

All the carbon-bonded hydrogen atoms were located at their ideal positions with the C—H=0.93 Å, C—H=0.96 Å, C—H=0.97 Å and $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$. All the hydrogen atoms bonded to the oxygen atoms were located from the difference maps and refined with the restraints of O—H=0.82 (1) Å and $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

View of the molecule of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The crystal packing for (I), with hydrogen bonds shown as dashed lines. Hydrogen atoms not involved in the hydrogen bonding have been omitted.

2,2'-[4-Acetyl-1,3-phenylenebis(oxy)]diacetic acid

Crystal data

C₁₂H₁₂O₇ $M_r = 268.22$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 5.1351$ (6) Å $b = 7.8346$ (9) Å $c = 15.6157$ (18) Å $\alpha = 86.217$ (2)° $\beta = 81.321$ (1)° $\gamma = 72.101$ (2)° $V = 590.86$ (12) Å³ $Z = 2$ $F(000) = 280$ $D_x = 1.508$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3359 reflections

 $\theta = 2.6$ – 31.8 ° $\mu = 0.13$ mm⁻¹ $T = 298$ K

Block, colourless

 $0.12 \times 0.10 \times 0.10$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2003)

 $T_{\min} = 0.985$, $T_{\max} = 0.987$

3902 measured reflections

2049 independent reflections

1860 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.016$ $\theta_{\max} = 25.0$ °, $\theta_{\min} = 1.3$ ° $h = -6 \rightarrow 6$ $k = -9 \rightarrow 9$ $l = -18 \rightarrow 18$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.036$ $wR(F^2) = 0.120$ $S = 1.05$

2049 reflections

176 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0795P)^2 + 0.1078P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.24$ e Å⁻³ $\Delta\rho_{\min} = -0.23$ e Å⁻³

Extinction correction: SHELXL97 (Sheldrick,

2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.036 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.9386 (3)	0.64900 (17)	0.22991 (8)	0.0282 (3)
C2	0.6877 (3)	0.75023 (17)	0.27294 (9)	0.0291 (3)

H2	0.6254	0.8735	0.2630	0.035*
C3	0.5297 (3)	0.66705 (18)	0.33088 (9)	0.0290 (3)
C4	0.6221 (3)	0.48246 (18)	0.34697 (9)	0.0332 (3)
H4	0.5166	0.4266	0.3858	0.040*
C5	0.8736 (3)	0.38464 (17)	0.30388 (9)	0.0329 (3)
H5	0.9361	0.2618	0.3152	0.040*
C6	1.0383 (3)	0.46109 (17)	0.24430 (9)	0.0299 (3)
C7	1.0249 (3)	0.90973 (18)	0.15901 (10)	0.0349 (3)
H7A	1.0211	0.9696	0.2118	0.042*
H7B	0.8424	0.9519	0.1412	0.042*
C8	1.2360 (3)	0.94824 (18)	0.08886 (9)	0.0329 (3)
C9	0.1228 (3)	0.70250 (18)	0.43152 (9)	0.0323 (3)
H9A	0.2214	0.6572	0.4806	0.039*
H9B	0.0869	0.6029	0.4068	0.039*
C10	-0.1449 (3)	0.84177 (18)	0.46089 (9)	0.0321 (3)
C11	1.3024 (3)	0.33765 (19)	0.20074 (10)	0.0363 (3)
C12	1.4694 (4)	0.3989 (2)	0.12491 (12)	0.0524 (5)
H12A	1.6153	0.2977	0.1007	0.079*
H12B	1.5473	0.4845	0.1436	0.079*
H12C	1.3527	0.4537	0.0818	0.079*
O1	1.1504 (2)	1.10815 (14)	0.05426 (8)	0.0472 (3)
H1	1.2707	1.1235	0.0164	0.071*
O2	1.4614 (2)	0.84065 (14)	0.06836 (7)	0.0463 (3)
O3	1.1036 (2)	0.72247 (13)	0.17277 (7)	0.0406 (3)
O4	0.28549 (19)	0.77925 (13)	0.36867 (7)	0.0369 (3)
O5	-0.2075 (2)	0.99593 (13)	0.43053 (7)	0.0421 (3)
O6	-0.2994 (2)	0.78060 (14)	0.51955 (8)	0.0487 (3)
H6	-0.4400	0.8616	0.5357	0.073*
O7	1.3789 (3)	0.18327 (14)	0.22649 (9)	0.0579 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0251 (6)	0.0269 (6)	0.0304 (6)	-0.0088 (5)	0.0037 (5)	0.0021 (5)
C2	0.0258 (7)	0.0232 (6)	0.0344 (7)	-0.0059 (5)	0.0025 (5)	0.0039 (5)
C3	0.0231 (6)	0.0288 (7)	0.0317 (7)	-0.0059 (5)	0.0019 (5)	0.0009 (5)
C4	0.0300 (7)	0.0300 (7)	0.0372 (7)	-0.0111 (5)	0.0041 (6)	0.0061 (5)
C5	0.0315 (7)	0.0233 (6)	0.0408 (8)	-0.0071 (5)	0.0005 (6)	0.0034 (5)
C6	0.0279 (7)	0.0247 (7)	0.0344 (7)	-0.0065 (6)	0.0003 (5)	-0.0003 (5)
C7	0.0299 (7)	0.0264 (7)	0.0420 (8)	-0.0064 (5)	0.0089 (6)	0.0039 (6)
C8	0.0306 (7)	0.0283 (7)	0.0375 (7)	-0.0101 (6)	0.0031 (6)	0.0028 (5)
C9	0.0266 (7)	0.0322 (7)	0.0342 (7)	-0.0087 (6)	0.0057 (5)	0.0036 (6)
C10	0.0292 (7)	0.0322 (7)	0.0328 (7)	-0.0105 (6)	0.0051 (5)	-0.0010 (5)
C11	0.0314 (7)	0.0283 (7)	0.0448 (8)	-0.0061 (6)	0.0033 (6)	-0.0045 (6)
C12	0.0432 (9)	0.0373 (8)	0.0602 (10)	-0.0011 (7)	0.0226 (8)	-0.0061 (7)
O1	0.0391 (6)	0.0365 (6)	0.0568 (7)	-0.0098 (5)	0.0118 (5)	0.0142 (5)
O2	0.0342 (6)	0.0394 (6)	0.0540 (7)	-0.0066 (5)	0.0142 (5)	0.0094 (5)
O3	0.0337 (6)	0.0259 (5)	0.0518 (6)	-0.0064 (4)	0.0185 (4)	0.0047 (4)

O4	0.0270 (5)	0.0298 (5)	0.0446 (6)	-0.0047 (4)	0.0126 (4)	0.0052 (4)
O5	0.0344 (6)	0.0333 (6)	0.0508 (6)	-0.0076 (4)	0.0102 (5)	0.0047 (4)
O6	0.0371 (6)	0.0384 (6)	0.0574 (7)	-0.0071 (5)	0.0228 (5)	0.0052 (5)
O7	0.0491 (7)	0.0285 (6)	0.0774 (9)	0.0037 (5)	0.0135 (6)	0.0050 (5)

Geometric parameters (Å, °)

C1—O3	1.3596 (16)	C8—O2	1.2154 (18)
C1—C2	1.3862 (19)	C8—O1	1.3033 (17)
C1—C6	1.4170 (18)	C9—O4	1.4166 (15)
C2—C3	1.3884 (18)	C9—C10	1.4979 (19)
C2—H2	0.9300	C9—H9A	0.9700
C3—O4	1.3642 (17)	C9—H9B	0.9700
C3—C4	1.3949 (19)	C10—O5	1.2323 (17)
C4—C5	1.381 (2)	C10—O6	1.2863 (17)
C4—H4	0.9300	C11—O7	1.2126 (18)
C5—C6	1.3927 (19)	C11—C12	1.496 (2)
C5—H5	0.9300	C12—H12A	0.9600
C6—C11	1.4966 (19)	C12—H12B	0.9600
C7—O3	1.4079 (16)	C12—H12C	0.9600
C7—C8	1.5050 (18)	O1—H1	0.8200
C7—H7A	0.9700	O6—H6	0.8200
C7—H7B	0.9700		
O3—C1—C2	122.68 (11)	O2—C8—C7	122.72 (12)
O3—C1—C6	116.25 (11)	O1—C8—C7	112.58 (12)
C2—C1—C6	121.06 (12)	O4—C9—C10	109.47 (11)
C1—C2—C3	119.83 (11)	O4—C9—H9A	109.8
C1—C2—H2	120.1	C10—C9—H9A	109.8
C3—C2—H2	120.1	O4—C9—H9B	109.8
O4—C3—C2	114.79 (11)	C10—C9—H9B	109.8
O4—C3—C4	124.56 (12)	H9A—C9—H9B	108.2
C2—C3—C4	120.65 (12)	O5—C10—O6	124.81 (13)
C5—C4—C3	118.49 (12)	O5—C10—C9	122.91 (12)
C5—C4—H4	120.8	O6—C10—C9	112.28 (11)
C3—C4—H4	120.8	O7—C11—C12	119.44 (13)
C4—C5—C6	123.14 (12)	O7—C11—C6	118.94 (13)
C4—C5—H5	118.4	C12—C11—C6	121.60 (12)
C6—C5—H5	118.4	C11—C12—H12A	109.5
C5—C6—C1	116.82 (12)	C11—C12—H12B	109.5
C5—C6—C11	117.25 (11)	H12A—C12—H12B	109.5
C1—C6—C11	125.93 (12)	C11—C12—H12C	109.5
O3—C7—C8	106.80 (11)	H12A—C12—H12C	109.5
O3—C7—H7A	110.4	H12B—C12—H12C	109.5
C8—C7—H7A	110.4	C8—O1—H1	109.5
O3—C7—H7B	110.4	C1—O3—C7	119.30 (10)
C8—C7—H7B	110.4	C3—O4—C9	117.02 (10)
H7A—C7—H7B	108.6	C10—O6—H6	109.5

O2—C8—O1	124.70 (13)		
O3—C1—C2—C3	-179.66 (12)	O3—C7—C8—O1	163.06 (12)
C6—C1—C2—C3	-0.3 (2)	O4—C9—C10—O5	2.36 (19)
C1—C2—C3—O4	-179.26 (11)	O4—C9—C10—O6	-178.61 (11)
C1—C2—C3—C4	0.6 (2)	C5—C6—C11—O7	-9.3 (2)
O4—C3—C4—C5	179.78 (12)	C1—C6—C11—O7	171.32 (14)
C2—C3—C4—C5	-0.1 (2)	C5—C6—C11—C12	168.98 (14)
C3—C4—C5—C6	-0.8 (2)	C1—C6—C11—C12	-10.4 (2)
C4—C5—C6—C1	1.1 (2)	C2—C1—O3—C7	2.9 (2)
C4—C5—C6—C11	-178.34 (13)	C6—C1—O3—C7	-176.45 (12)
O3—C1—C6—C5	178.87 (12)	C8—C7—O3—C1	-176.25 (11)
C2—C1—C6—C5	-0.52 (19)	C2—C3—O4—C9	-176.96 (11)
O3—C1—C6—C11	-1.8 (2)	C4—C3—O4—C9	3.1 (2)
C2—C1—C6—C11	178.85 (12)	C10—C9—O4—C3	-175.88 (11)
O3—C7—C8—O2	-17.6 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots O2 ⁱ	0.82	1.83	2.6530 (14)	180
O6—H6 \cdots O5 ⁱⁱ	0.82	1.83	2.6352 (15)	167
C2—H2 \cdots O7 ⁱⁱⁱ	0.93	2.44	3.3566 (17)	168
C4—H4 \cdots O6 ^{iv}	0.93	2.52	3.4392 (19)	169

Symmetry codes: (i) $-x+3, -y+2, -z$; (ii) $-x-1, -y+2, -z+1$; (iii) $x-1, y+1, z$; (iv) $-x, -y+1, -z+1$.