

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

6-[4-Chloro-2-(trifluoromethyl)phenyl]-3-fluoro-2-methylpyridine

S. Sreenivasa,^{a*} K. E. Manojkumar,^a P. A. Suchetan,^b N. R. Mohan,^a B. S. Palakshamurthy,^c T. Srinivasan^d and D. Velmurgan^d

^aDepartment of Studies and Research in Chemistry, Tumkur University, Tumkur, Karnataka 572 103, India, ^bDepartment of Studies and Research in Chemistry, U.C.S, Tumkur University, Tumkur, Karnataka 572 103, India, ^cDepartment of Studies and Research in Physics, U.C.S., Tumkur University, Tumkur, Karnataka 572 103, India, and ^dCentre of Advanced Study in Crystallography and Biophysics, University of Madras Guindy Campus, Chennai 600 025, India
Correspondence e-mail: drsreenivasa@yahoo.co.in


Received 14 October 2012; accepted 8 November 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.042; wR factor = 0.129; data-to-parameter ratio = 17.5.

In the title compound, $\text{C}_{13}\text{H}_8\text{ClF}_4\text{N}$, the dihedral angle between the benzene and pyridine rings is $59.8(3)^\circ$. In the crystal, molecules are stacked in columns along the b axis through weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the biological activity of pyridine derivatives, see: Patrick & Kinsmar (1996); Hishmat *et al.* (1990); Doshi *et al.* (1999); Bhatt *et al.* (2001).



Experimental

Crystal data

$\text{C}_{13}\text{H}_8\text{ClF}_4\text{N}$
 $M_r = 289.65$
Monoclinic, $P2_1/n$
 $a = 13.1813(7)$ Å
 $b = 4.5837(3)$ Å

$c = 20.4448(11)$ Å
 $\beta = 92.441(3)^\circ$
 $V = 1234.14(12)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.34$ mm⁻¹
 $T = 293$ K

 $0.2 \times 0.18 \times 0.16$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
11077 measured reflections

3033 independent reflections
2116 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.129$
 $S = 1.01$
3033 reflections

173 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the N1/C8–C12 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C13}-\text{H13B}\cdots\text{Cg1}^{\text{i}}$	0.96	2.83	3.606 (2)	138

Symmetry code: (i) $x, y + 1, z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 2012); software used to prepare material for publication: SHELXL97.

Authors thank Dr. S. C. Sharma, Vice Chancellor, Tumkur University, Tumkur for his constant encouragement and G.B. Sadananda, Department of Studies and Research in Physics, U.C.S. Tumkur University, Tumkur, for his help and valuable suggestions. BSP thanks Dr. H. C. Devaraje Gowda, Department of Physics, Yuvarajas College (constituent), University of Mysore for his guidance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5209).

References

- Bhatt, A. H., Parekh, M. H., Parikh, K. A. & Parikh, A. R. (2001). *J. Indian Chem. Soc.* **40**, 57–21.
Bruker (2004). APEX2 and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
Doshi, R., Kagthara, P. & Parekh, H. (1999). *Indian J. Chem.* **38**, 348–352.
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
Hishmat, O. H., Abdel Galil, F. M. & Farrag, D. S. (1990). *Pharmazie*, **45**, 793–795.
Patrick, G. L. & Kinsmar, O. S. (1996). *Eur. J. Med. Chem.* **31**, 615–624.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2012). E68, o3370 [doi:10.1107/S1600536812046211]

6-[4-Chloro-2-(trifluoromethyl)phenyl]-3-fluoro-2-methylpyridine

S. Sreenivasa, K. E. Manojkumar, P. A. Suchetan, N. R. Mohan, B. S. Palakshamurthy, T. Srinivasan and D. Velmurgan

S1. Comment

Pyridine derivatives of different heterocyclic nucleus have shown potent pharmacological properties like antifungal (Patrick & Kinsmar, 1996), antitubercular (Hishmat *et al.*, 1990) and antibacterial (Doshi *et al.*, 1999; Bhatt *et al.*, 2001). Keeping in view of the biological importance of this class of compound, we synthesized the title compound to study its X-ray crystal structure.


In the title compound, the dihedral angle between the least square planes through the benzene ring and the pyridine ring is 59.8 (3)°. In the absence of hydrogen bonds, the structure is stabilized by a weak intermolecular C—H... π interaction (Table 1).

S2. Experimental


2-Methyl-3-fluoropyridine-6-boronic acid (8.4 mmol) was taken in a mixture of dioxane (20 ml) and water (5 ml) at RT under nitrogen atmosphere. The reaction mixture was degassed with argon for 10 min, and K₂CO₃ (23.1 mmol) and dichlorobis(triphenylphosphene)-palladium(II) (0.231 mmol) were added and again degassed for 10 min. 2-Bromo-5-chlorobenzotrifluoride (7.7 mmol) was added and then reaction mixture was heated to 100 °C for 1 h (reaction was monitored by TLC), and the reaction mass was cooled to RT, diluted with ethyl acetate, filtered over celite, and washed with ethyl acetate. The filtrate was washed with water and brine, and the ethyl acetate layer was dried with anhydrous Na₂SO₄ and concentrated. The crude product was purified by triturating with petroleum ether. Single crystals of the title compound used for X-ray diffraction studies were obtained from slow evaporation of the solution of the compound in petroleum ether-ethyl acetate mixture (1:1).

S3. Refinement

The H atoms were positioned with idealized geometry (C—H = 0.93–0.96 Å) using a riding model with $U_{iso}(H) = 1.2$ or $1.5U_{eq}(C)$.

**Figure 1**

Molecular structure of the title compound, showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A packing diagram of the title compound, showing C—H... π interactions (dotted lines).

6-[4-Chloro-2-(trifluoromethyl)phenyl]-3-fluoro-2-methylpyridine

Crystal data

C₁₃H₈ClF₄N $M_r = 289.65$ Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

 $a = 13.1813 (7) \text{ \AA}$ $b = 4.5837 (3) \text{ \AA}$ $c = 20.4448 (11) \text{ \AA}$ $\beta = 92.441 (3)^\circ$ $V = 1234.14 (12) \text{ \AA}^3$ $Z = 4$ $F(000) = 584$

Prism

 $D_x = 1.559 \text{ Mg m}^{-3}$

Melting point: 408 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2116 reflections

 $\theta = 1.8\text{--}26.8^\circ$ $\mu = 0.34 \text{ mm}^{-1}$ $T = 293 \text{ K}$

Prism, colourless

 $0.2 \times 0.18 \times 0.16 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 1.20 pixels mm^{-1}

multi-scan

11077 measured reflections

3033 independent reflections

2116 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.031$ $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 1.8^\circ$ $h = -17 \rightarrow 17$ $k = -5 \rightarrow 6$ $l = -27 \rightarrow 27$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.042$ $wR(F^2) = 0.129$ $S = 1.01$

3033 reflections

173 parameters

0 restraints

0 constraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0549P)^2 + 0.4806P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.001$ $\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.83131 (5)	0.3848 (2)	0.03175 (3)	0.0883 (3)
F1	1.12038 (13)	1.0835 (4)	0.09837 (7)	0.0978 (6)
F2	1.18924 (9)	0.8102 (3)	0.17220 (10)	0.0964 (5)
F3	1.11540 (11)	1.2001 (3)	0.19839 (7)	0.0781 (4)

F4	1.09214 (11)	1.1818 (4)	0.46328 (6)	0.0853 (5)
N1	0.93981 (11)	1.0496 (3)	0.31763 (7)	0.0451 (3)
C1	0.87112 (13)	0.6135 (5)	0.21835 (9)	0.0528 (5)
H1	0.8368	0.5883	0.2568	0.063*
C2	0.83018 (14)	0.4976 (5)	0.16091 (9)	0.0589 (5)
H2	0.7693	0.395	0.1605	0.071*
C3	0.88083 (15)	0.5361 (5)	0.10418 (9)	0.0565 (5)
C4	0.96922 (14)	0.6939 (5)	0.10389 (9)	0.0552 (5)
H4	1.0015	0.7239	0.0649	0.066*
C5	1.01029 (13)	0.8085 (4)	0.16185 (9)	0.0472 (4)
C6	0.96206 (12)	0.7668 (4)	0.22062 (9)	0.0445 (4)
C7	1.10812 (16)	0.9742 (5)	0.15823 (11)	0.0609 (5)
C8	1.00181 (13)	0.8725 (4)	0.28603 (9)	0.0453 (4)
C9	1.09464 (14)	0.7800 (5)	0.31331 (11)	0.0587 (5)
H9	1.1349	0.6502	0.291	0.07*
C10	1.12574 (16)	0.8841 (5)	0.37399 (11)	0.0638 (6)
H10	1.1876	0.8282	0.3937	0.077*
C11	1.06333 (16)	1.0706 (5)	0.40409 (10)	0.0571 (5)
C12	0.96959 (14)	1.1523 (4)	0.37664 (9)	0.0485 (4)
C13	0.90004 (18)	1.3525 (5)	0.41073 (10)	0.0642 (6)
H13A	0.8421	1.3959	0.3823	0.096*
H13B	0.9352	1.5301	0.422	0.096*
H13C	0.8779	1.2605	0.4498	0.096*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0773 (4)	0.1319 (7)	0.0551 (3)	-0.0171 (4)	-0.0033 (3)	-0.0247 (4)
F1	0.1080 (12)	0.1124 (13)	0.0752 (9)	-0.0472 (10)	0.0308 (8)	0.0014 (9)
F2	0.0459 (7)	0.0725 (10)	0.1719 (16)	-0.0025 (6)	0.0166 (8)	-0.0082 (10)
F3	0.0911 (9)	0.0542 (8)	0.0897 (10)	-0.0197 (7)	0.0125 (7)	-0.0126 (7)
F4	0.0926 (10)	0.0965 (11)	0.0645 (8)	0.0001 (8)	-0.0251 (7)	-0.0226 (8)
N1	0.0467 (8)	0.0453 (8)	0.0433 (7)	0.0016 (6)	0.0034 (6)	0.0002 (6)
C1	0.0444 (9)	0.0704 (13)	0.0438 (9)	-0.0028 (9)	0.0057 (7)	0.0011 (9)
C2	0.0436 (9)	0.0797 (15)	0.0535 (11)	-0.0116 (10)	0.0012 (8)	-0.0005 (10)
C3	0.0505 (10)	0.0717 (14)	0.0471 (10)	0.0010 (10)	0.0000 (8)	-0.0079 (10)
C4	0.0539 (10)	0.0649 (13)	0.0477 (10)	0.0011 (9)	0.0122 (8)	-0.0030 (9)
C5	0.0432 (9)	0.0463 (10)	0.0526 (10)	0.0013 (8)	0.0078 (7)	-0.0038 (8)
C6	0.0410 (8)	0.0463 (10)	0.0464 (9)	0.0061 (7)	0.0032 (7)	-0.0021 (8)
C7	0.0592 (12)	0.0535 (12)	0.0710 (13)	-0.0077 (10)	0.0164 (10)	-0.0069 (11)
C8	0.0431 (9)	0.0460 (10)	0.0470 (9)	0.0014 (7)	0.0024 (7)	-0.0006 (8)
C9	0.0483 (10)	0.0615 (13)	0.0656 (12)	0.0120 (9)	-0.0042 (9)	-0.0081 (10)
C10	0.0507 (11)	0.0709 (15)	0.0684 (13)	0.0043 (10)	-0.0145 (9)	-0.0026 (11)
C11	0.0626 (12)	0.0567 (12)	0.0508 (10)	-0.0059 (10)	-0.0099 (9)	-0.0062 (9)
C12	0.0557 (10)	0.0437 (10)	0.0460 (9)	-0.0040 (8)	0.0030 (8)	0.0001 (8)
C13	0.0785 (14)	0.0619 (14)	0.0526 (11)	0.0043 (11)	0.0078 (10)	-0.0098 (10)

Geometric parameters (Å, °)

C1—C3	1.737 (2)	C5—C7	1.501 (3)
F1—C7	1.339 (3)	C6—C1	1.388 (3)
F2—C7	1.328 (3)	C6—C5	1.396 (2)
F3—C7	1.322 (2)	C8—C9	1.389 (2)
F4—C11	1.353 (2)	C8—C6	1.496 (2)
N1—C8	1.338 (2)	C9—C10	1.375 (3)
N1—C12	1.338 (2)	C9—H9	0.93
C1—C2	1.378 (3)	C10—H10	0.93
C1—H1	0.93	C11—C10	1.352 (3)
C2—H2	0.93	C11—C12	1.387 (3)
C3—C2	1.374 (3)	C12—C13	1.491 (3)
C4—C3	1.372 (3)	C13—H13A	0.96
C4—H4	0.93	C13—H13B	0.96
C5—C4	1.385 (3)	C13—H13C	0.96
C8—N1—C12	119.16 (15)	F2—C7—C5	112.85 (18)
C2—C1—C6	121.99 (17)	F1—C7—C5	111.83 (18)
C2—C1—H1	119	N1—C8—C9	122.48 (17)
C6—C1—H1	119	N1—C8—C6	115.49 (15)
C3—C2—C1	118.89 (18)	C9—C8—C6	121.94 (17)
C3—C2—H2	120.6	C10—C9—C8	118.67 (19)
C1—C2—H2	120.6	C10—C9—H9	120.7
C4—C3—C2	120.98 (18)	C8—C9—H9	120.7
C4—C3—C1	119.69 (15)	C11—C10—C9	117.78 (18)
C2—C3—C1	119.33 (16)	C11—C10—H10	121.1
C3—C4—C5	119.83 (17)	C9—C10—H10	121.1
C3—C4—H4	120.1	F4—C11—C10	119.50 (18)
C5—C4—H4	120.1	F4—C11—C12	118.13 (19)
C4—C5—C6	120.57 (17)	C10—C11—C12	122.36 (18)
C4—C5—C7	117.07 (17)	N1—C12—C11	119.46 (18)
C6—C5—C7	122.37 (17)	N1—C12—C13	118.39 (17)
C1—C6—C5	117.69 (17)	C11—C12—C13	122.15 (18)
C1—C6—C8	117.57 (16)	C12—C13—H13A	109.5
C5—C6—C8	124.74 (16)	C12—C13—H13B	109.5
F3—C7—F2	105.85 (18)	H13A—C13—H13B	109.5
F3—C7—F1	105.45 (18)	C12—C13—H13C	109.5
F2—C7—F1	106.32 (18)	H13A—C13—H13C	109.5
F3—C7—C5	113.91 (17)	H13B—C13—H13C	109.5

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the N1/C8—C12 ring.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C13—H13B \cdots Cg1 ⁱ	0.96	2.83	3.606 (2)	138

Symmetry code: (i) *x*, *y*+1, *z*.