

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Pyrimidine-2,4-diamine acetone monosolvate

 Sergiu Draguta,^{a*} Bhupinder Sandhu,^b Victor N. Khrustalev,^c Marina S. Fonari^d and Tatiana V. Timofeeva^b

^aD. Ghitu Institute of Electronic Engineering and Nanotechnologies, 3/3 Academy Street, MD-2028, Chisinau, Republic of Moldova, ^bDepartment of Biology & Chemistry, New Mexico Highlands University, 803 University Avenue, Las Vegas, NM 87701, USA, ^cX-Ray Structural Centre, A.N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 28 Vavilov Street, B-334, Moscow 119991, Russian Federation, and ^dInstitute of Applied Physics Academy of Science of Moldova, 5 Academy Street, MD-2028, Chisinau, Republic of Moldova.

Correspondence e-mail: sergiudraguta@gmail.com


Received 1 January 2013; accepted 11 January 2013

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.048; wR factor = 0.139; data-to-parameter ratio = 17.1.

In the title compound, $C_4H_6N_4 \cdot C_3H_6O$, the pyrimidine-2,4-diamine molecule is nearly planar (r.m.s. deviation = 0.005 Å), with the endocyclic angles covering the range 114.36 (10)–126.31 (10)°. In the crystal, $N-H \cdots N$ and $N-H \cdots O$ hydrogen bonds link the molecules into ribbons along [101], and weak $C-H \cdots \pi$ interactions consolidate further the crystal packing.

Related literature

For the biological activity of pyrimidine derivatives, see: Hall *et al.* (1993); Gengeliczki *et al.* (2011). For the crystal structures of related compounds, see: Bertolasi *et al.* (2002); Draguta *et al.* (2012). For bond lengths in organic compounds, see: Allen *et al.* (1987). For hydrogen-bonding graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $C_4H_6N_4 \cdot C_3H_6O$
 $M_r = 168.21$

 Monoclinic, $P2_1/c$
 $a = 8.1594$ (15) Å
 $b = 12.728$ (2) Å
 $c = 8.7663$ (16) Å
 $\beta = 99.395$ (3)°
 $V = 898.2$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

 Bruker APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2003)
 $T_{min} = 0.974$, $T_{max} = 0.982$

 9693 measured reflections
 2170 independent reflections
 1752 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.051$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.139$
 $S = 1.07$
 2170 reflections
 127 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.28$ e Å⁻³
 $\Delta\rho_{min} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

 C_g is the centroid of the pyrimidine ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N2-H2A \cdots N1^i$	0.875 (18)	2.191 (18)	3.0608 (18)	177.3 (15)
$N2-H2B \cdots O1$	0.871 (16)	2.247 (19)	3.0990 (17)	164.7 (16)
$N4-H4A \cdots O1^{ii}$	0.879 (17)	2.170 (18)	2.9141 (16)	142.2 (15)
$N4-H4B \cdots N3^{ii}$	0.900 (18)	2.120 (19)	3.0171 (17)	174.9 (15)
$C9-H9C \cdots C_g^{iii}$	0.96	2.63	3.5484 (17)	159

 Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The authors are grateful for NSF support via DMR grant 0934212 (PREM) and CHE 0832622.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5381).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bertolasi, V., Pretto, L., Gilli, P., Ferretti, V. & Gilli, G. (2002). *New J. Chem.* **26**, 1559–1566.
- Bruker (2001). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
- Draguta, S., Khrustalev, V. N., Fonari, M. S., Antipin, M. Y. & Timofeeva, T. V. (2012). *Acta Cryst.* **E68**, o3353.
- Gengeliczki, Z., Callahan, M. P., Kabeláč, M., Rijs, A. M. & de Vries, M. S. (2011). *J. Phys. Chem. A*, **115**, 11423–11427.
- Hall, I. A., Campbell, K. L., Chambers, M. D. & Davis, C. N. (1993). *J. Am. Vet. Med. Assoc.* **202**, 1959–1962.
- Sheldrick, G. M. (2003). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2013). E69, o251 [doi:10.1107/S1600536813001025]

Pyrimidine-2,4-diamine acetone monosolvate

Sergiu Draguta, Bhupinder Sandhu, Victor N. Khrustalev, Marina S. Fonari and Tatiana V. Timofeeva

S1. Comment

Pyrimidine derivatives are biologically important compounds, because they occur in nature as components of nucleic acids. Pyrimidine-2,4-diamine reacts with 3,4,5-trimethoxybenzyl to form trimethoprim which acts as the nucleic acid inhibitor (Hall *et al.*, 1993) as well as with 3,7-dimethylxanthine to form clusters which represent potential alternate nucleobase pairs, geometrically equivalent to guanine-cytosine (Gengeliczki *et al.* 2011). In the area of drug design and pharmacore mapping, there are several compounds comprising the 2,4-diaminopyrimidinium cations and β - or ζ -diketoenolate anions bound into supramolecular synthons by intermolecular hydrogen bonds (Bertolasi *et al.*, 2002). The presented here crystal structure of the title compound, C₄H₆N₄.C₃H₆O, (**I**) (Figure 1) was determined to study its hydrogen bonding system and to use it in the future for the design of co-crystals with particular properties.

The asymmetric unit of **I** consists of a pyrimidine-2,4-diamine molecule and an acetone solvate molecule. The pyrimidine-2,4-diamine molecule is planar (r.m.s. = 0.005 Å), with the endocyclic angles covering range of 114.36 (10)–126.31 (10)°. The endocyclic angles at the C2, C4 and C6 carbon atoms adjacent to the N1 and N3 heteroatoms are larger than 120°, and those at the other atoms of the ring are smaller than 120°. The analogous distribution of the endocyclic angles was recently observed by us within the related pyridine-2,5-diamine (Draguta *et al.*, 2012). The bond lengths have the usual values (Allen *et al.*, 1987).

In the crystal, each pyrimidine molecule is connected to two others ones by the intermolecular N2—H2A[⋯]N1ⁱ and N4—H4B[⋯]N3ⁱⁱ hydrogen bonds (centrosymmetric R_2^2 (8) ring motifs (Bernstein *et al.*, 1995); Table 1), forming the infinite ribbons toward [101] (Figure 2). The acetone molecules are bonded to the ribbons as pendant molecules *via* two intermolecular N2—H2B[⋯]O1 and N4—H4A[⋯]O1ⁱⁱ hydrogen bonds (Table 1, Figure 2), and are almost coplanar to the ribbon planes (the dihedral angle between the 2,4-pyrimidine and acetone molecules is 7.33 (2)°). The ribbons are packed into layers parallel to (1 0 $\bar{1}$), with the interlayer distance of 3.8827 (16) Å). The layers are linked to each other by the intermolecular C9—H9C[⋯] π (pyrimidine ring) interactions (Table 1).

S2. Experimental

The compound **I** was obtained commercially (Aldrich) as a fine-crystalline powder. Crystals suitable for the X-ray diffraction study were grown by slow evaporation from acetone solution. The crystals of **I** are very sensitive to air and moisture. Therefore, to keep the quality of the crystal during the experiment, the crystal of **I** was mounted in vaseline oil.

S3. Refinement

The hydrogen atoms of the amino groups were localized in the difference Fourier maps and refined isotropically. The other hydrogen atoms were placed in the calculated positions with C—H = 0.93 Å (CH-groups) and 0.96 Å (CH₃-groups) and refined in the riding model with fixed isotropic displacement parameters [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$] for the CH₃-groups and

$1.2U_{eq}(C)$ for the CH-groups].




Figure 1

Molecular structure of **I**. Displacement ellipsoids are shown at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A portion of the crystal packing showing the H-bonded ribbons toward [101]. Dashed lines indicate the intermolecular N—H...N and N—H...O hydrogen bonds.

Pyrimidine-2,4-diamine acetone monosolvate

Crystal data

$C_4H_6N_4 \cdot C_3H_6O$

$M_r = 168.21$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 8.1594$ (15) Å

$b = 12.728$ (2) Å

$c = 8.7663$ (16) Å

$\beta = 99.395$ (3)°

$V = 898.2$ (3) Å³

$Z = 4$

$F(000) = 360$

$D_x = 1.244$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8851 reflections

$\theta = 2.3$ – 32.1 °

$\mu = 0.09$ mm⁻¹

$T = 296$ K

Prism, colourless

$0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker APEXII CCD diffractometer	9693 measured reflections
Radiation source: fine-focus sealed tube	2170 independent reflections
Graphite monochromator	1752 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.051$
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	$\theta_{\text{max}} = 28.0^\circ$, $\theta_{\text{min}} = 2.5^\circ$
$T_{\text{min}} = 0.974$, $T_{\text{max}} = 0.982$	$h = -10 \rightarrow 10$
	$k = -16 \rightarrow 16$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.048$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.139$	$w = 1/[\sigma^2(F_o^2) + (0.0884P)^2]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
2170 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
127 parameters	$\Delta\rho_{\text{max}} = 0.28 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.28 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.12301 (13)	0.39554 (8)	0.11848 (12)	0.0224 (3)
C2	0.19664 (15)	0.47490 (9)	0.20658 (14)	0.0191 (3)
N2	0.14277 (14)	0.57309 (9)	0.16744 (13)	0.0255 (3)
H2A	0.0691 (19)	0.5838 (13)	0.0852 (18)	0.025 (4)*
H2B	0.197 (2)	0.6249 (13)	0.2183 (19)	0.033 (4)*
N3	0.31691 (12)	0.46530 (8)	0.33213 (12)	0.0187 (3)
C4	0.36727 (15)	0.36728 (9)	0.37483 (14)	0.0190 (3)
N4	0.48650 (14)	0.35701 (8)	0.49903 (13)	0.0240 (3)
H4A	0.5237 (19)	0.2944 (14)	0.5294 (18)	0.033 (4)*
H4B	0.539 (2)	0.4120 (14)	0.5498 (19)	0.031 (4)*
C5	0.29627 (16)	0.27903 (10)	0.29098 (15)	0.0245 (3)
H5	0.3283	0.2109	0.3202	0.029*
C6	0.17840 (16)	0.29891 (10)	0.16502 (15)	0.0248 (3)
H6	0.1329	0.2417	0.1070	0.030*
O1	0.28515 (12)	0.78374 (7)	0.30884 (11)	0.0285 (3)
C7	0.30086 (19)	0.97014 (11)	0.31631 (19)	0.0333 (4)

H7A	0.3699	0.9583	0.4144	0.050*
H7B	0.2067	1.0120	0.3308	0.050*
H7C	0.3635	1.0063	0.2489	0.050*
C8	0.24238 (15)	0.86704 (9)	0.24592 (14)	0.0223 (3)
C9	0.12660 (17)	0.86932 (11)	0.09351 (16)	0.0297 (3)
H9A	0.1316	0.8032	0.0418	0.045*
H9B	0.1592	0.9246	0.0303	0.045*
H9C	0.0151	0.8816	0.1112	0.045*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0249 (6)	0.0168 (5)	0.0226 (5)	-0.0012 (4)	-0.0046 (4)	-0.0021 (4)
C2	0.0203 (6)	0.0162 (6)	0.0199 (6)	-0.0007 (4)	0.0003 (5)	-0.0009 (4)
N2	0.0304 (6)	0.0153 (6)	0.0260 (6)	-0.0001 (4)	-0.0093 (5)	0.0003 (4)
N3	0.0209 (5)	0.0125 (5)	0.0212 (5)	-0.0003 (4)	-0.0012 (4)	0.0001 (4)
C4	0.0190 (6)	0.0155 (6)	0.0217 (6)	-0.0001 (4)	0.0006 (4)	0.0003 (4)
N4	0.0255 (6)	0.0134 (5)	0.0288 (6)	0.0008 (4)	-0.0080 (4)	0.0012 (4)
C5	0.0270 (7)	0.0138 (6)	0.0303 (7)	0.0007 (5)	-0.0025 (5)	-0.0011 (5)
C6	0.0279 (7)	0.0162 (6)	0.0277 (6)	-0.0012 (5)	-0.0027 (5)	-0.0046 (5)
O1	0.0325 (5)	0.0184 (5)	0.0314 (5)	0.0024 (4)	-0.0042 (4)	0.0037 (4)
C7	0.0376 (8)	0.0210 (7)	0.0424 (8)	-0.0053 (6)	0.0099 (6)	-0.0066 (6)
C8	0.0217 (6)	0.0182 (6)	0.0268 (6)	0.0007 (5)	0.0029 (5)	0.0014 (5)
C9	0.0271 (7)	0.0311 (7)	0.0289 (7)	0.0032 (5)	-0.0017 (5)	0.0059 (5)

Geometric parameters (Å, °)

N1—C6	1.3503 (16)	C5—H5	0.9300
N1—C2	1.3513 (16)	C6—H6	0.9300
C2—N2	1.3505 (16)	O1—C8	1.2196 (15)
C2—N3	1.3552 (16)	C7—C8	1.4948 (18)
N2—H2A	0.871 (16)	C7—H7A	0.9600
N2—H2B	0.875 (18)	C7—H7B	0.9600
N3—C4	1.3474 (15)	C7—H7C	0.9600
C4—N4	1.3428 (16)	C8—C9	1.5054 (18)
C4—C5	1.4137 (17)	C9—H9A	0.9600
N4—H4A	0.879 (17)	C9—H9B	0.9600
N4—H4B	0.900 (18)	C9—H9C	0.9600
C5—C6	1.3644 (18)		
C6—N1—C2	114.36 (11)	N1—C6—H6	117.6
N2—C2—N1	116.78 (11)	C5—C6—H6	117.6
N2—C2—N3	116.88 (11)	C8—C7—H7A	109.5
N1—C2—N3	126.32 (11)	C8—C7—H7B	109.5
C2—N2—H2A	120.5 (11)	H7A—C7—H7B	109.5
C2—N2—H2B	116.9 (11)	C8—C7—H7C	109.5
H2A—N2—H2B	121.8 (15)	H7A—C7—H7C	109.5
C4—N3—C2	117.17 (10)	H7B—C7—H7C	109.5

N4—C4—N3	117.59 (11)	O1—C8—C7	121.87 (12)
N4—C4—C5	121.69 (11)	O1—C8—C9	120.66 (11)
N3—C4—C5	120.71 (11)	C7—C8—C9	117.47 (12)
C4—N4—H4A	120.2 (11)	C8—C9—H9A	109.5
C4—N4—H4B	123.3 (10)	C8—C9—H9B	109.5
H4A—N4—H4B	116.2 (15)	H9A—C9—H9B	109.5
C6—C5—C4	116.64 (12)	C8—C9—H9C	109.5
C6—C5—H5	121.7	H9A—C9—H9C	109.5
C4—C5—H5	121.7	H9B—C9—H9C	109.5
N1—C6—C5	124.78 (11)		
C6—N1—C2—N2	-178.34 (11)	C2—N3—C4—C5	-0.04 (18)
C6—N1—C2—N3	0.25 (19)	N4—C4—C5—C6	-178.77 (12)
N2—C2—N3—C4	177.85 (11)	N3—C4—C5—C6	1.21 (19)
N1—C2—N3—C4	-0.75 (19)	C2—N1—C6—C5	1.10 (19)
C2—N3—C4—N4	179.95 (11)	C4—C5—C6—N1	-1.8 (2)

Hydrogen-bond geometry (Å, °)

C_g is the centroid of the pyrimidine ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2 <i>A</i> ...N1 ⁱ	0.875 (18)	2.191 (18)	3.0608 (18)	177.3 (15)
N2—H2 <i>B</i> ...O1	0.871 (16)	2.247 (19)	3.0990 (17)	164.7 (16)
N4—H4 <i>A</i> ...O1 ⁱⁱ	0.879 (17)	2.170 (18)	2.9141 (16)	142.2 (15)
N4—H4 <i>B</i> ...N3 ⁱⁱ	0.900 (18)	2.120 (19)	3.0171 (17)	174.9 (15)
C9—H9 <i>C</i> ...C _g ⁱⁱⁱ	0.96	2.63	3.5484 (17)	159

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+1, -y+1, -z+1$; (iii) $-x, y+1/2, -z+1/2$.