

$b = 11.0309 (4) \text{ \AA}$	$Z = 4$
$c = 13.8161 (6) \text{ \AA}$	$\text{Mo } K\alpha \text{ radiation}$
$\alpha = 102.557 (2)^\circ$	$\mu = 0.30 \text{ mm}^{-1}$
$\beta = 100.646 (2)^\circ$	$T = 293 \text{ K}$
$\gamma = 103.704 (2)^\circ$	$0.35 \times 0.30 \times 0.25 \text{ mm}$
$V = 1296.29 (9) \text{ \AA}^3$	

Methyl 2-[(2-chloroquinolin-3-yl)- (hydroxy)methyl]acrylate

T. Anuradha,^a J. Srinivasan,^b P. R. Seshadri^{a*} and M. Bakthadoss^b

^aPost Graduate and Research Department of Physics, Agurchand Manmull Jain College, Chennai 600 114, India, and ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India
Correspondence e-mail: seshadri_pr@yahoo.com


Received 22 April 2013; accepted 21 May 2013

Key indicators: single-crystal X-ray study; $T = 293 \text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$; R factor = 0.032; wR factor = 0.093; data-to-parameter ratio = 13.0.

There are two independent molecules (*A* and *B*) in the asymmetric unit of the title compound, $C_{14}H_{12}\text{ClNO}_3$. The mean planes of the methyl ester unit ($\text{C}_{\text{methyl}}-\text{O}-\text{C}=\text{O}$; r.m.s. deviation = 0.051 \AA for molecule *A* and 0.016 \AA for molecule *B*) and the chloroquinoline ring system (r.m.s. deviation = 0.023 \AA for molecule *A* and 0.014 \AA for molecule *B*) form dihedral angles of $63.5 (1)^\circ$ in molecule *A* and $78.1 (1)^\circ$ in molecule *B*. The main difference between the two independent molecules is reflected in the $(\text{H})\text{O}-\text{C}-\text{C}=\text{C}(\text{H}_2)$ torsion angle which is $-109.7 (2)^\circ$ in molecule *A* and $10.6 (2)^\circ$ in molecule *B*. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is observed in molecule *A*. In the crystal, molecules *A* and *B* are linked into pairs *via* bifurcated $\text{O}-\text{H}\cdots(\text{N},\text{Cl})$ hydrogen bonds and a weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond links pairs of molecules into chains along [100].

Related literature

For the biological activity of quinoline compounds, see: Biavatti *et al.* (2002); Towers *et al.* (1981); Shen *et al.* (1999). For their luminescent properties, see: Montes *et al.* (2006). For applications of acrylate compounds, see: Bhatia *et al.* (2007); Sharma (2011). For conformational aspects of methyl esters, see: Dunitz & Schweizer (1982). For resonance effects in acrylates, see: Merlino (1971); Varghese *et al.* (1986).



Experimental

Crystal data

$C_{14}H_{12}\text{ClNO}_3$
 $M_r = 277.70$

Triclinic, $P\bar{1}$
 $a = 9.2614 (4) \text{ \AA}$

Data collection

Bruker SMART APEXII diffractometer
14402 measured reflections

4466 independent reflections
3767 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.093$
 $S = 1.03$
4466 reflections

344 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.29 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots\text{A}$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
O1A—H1A···O2A	0.82	2.24	2.8372 (19)	130
O1B—H1B···Cl1A ⁱ	0.82	2.79	3.5040 (12)	147
O1B—H1B···N1A ⁱ	0.82	2.16	2.8609 (17)	144
CSA—H5A···O1B ⁱⁱ	0.93	2.56	3.451 (2)	162

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*, *PLATON* and *publCIF* (Westrip, 2010).

The authors thank Dr Babu Varghese, SAIF, IIT-Madras, India, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5607).

References

- Bhatia, S. P., Wellington, G. A., Cochiaro, J., Lalko, J., Letizia, C. S. & Api, A. M. (2007). *J. Food Chem. Toxicol.* **45**, S113–S119.
- Biavatti, M. W., Vieira, P. C., da Silva, M. F. G. F., Fernandes, J. B., Victor, S. R., Pagnocca, F. C., Albuquerque, S., Caracelli, I. & Zukerman-Schpector, J. (2002). *J. Braz. Chem. Soc.* **13**, 66–70.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Ins., Madison, Wisconsin, USA.
- Dunitz, J. D. & Schweizer, B. W. (1982). *Helv. Chim. Acta* **65**, 1547–1554.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Merlino, S. (1971). *Acta Cryst.* **B27**, 2491–2492.
- Montes, V. A., Pohl, R., Shinar, J. & Anzenbacher, P. Jr (2006). *Chem. Eur. J.* **12**, 4523–4535.
- Sharma, P. (2011). *J. Chem. Pharm. Res.* **3**, 403–423.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shen, A. Y., Wu, S. N. & Chiu, C. T. (1999). *J. Pharm. Pharmacol.* **51**, 543–548.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Towers, G. H. N., Graham, E. A., Spenser, I. D. & Abramowski, Z. (1981). *Planta Med.* **41**, 136–142.
- Varghese, B., Srinivasan, S., Padmanabhan, P. V. & Ramadas, S. R. (1986). *Acta Cryst.* **C42**, 1544–1546.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2013). E69, o990 [doi:10.1107/S1600536813014050]

Methyl 2-[(2-chloroquinolin-3-yl)(hydroxy)methyl]acrylate

T. Anuradha, J. Srinivasan, P. R. Seshadri and M. Bakthadoss

S1. Comment

The quinoline ring is found in compounds with antifungal (Biavatti *et al.*, 2002), antibacterial (Towers *et al.*, 1981) and anticancer (Shen *et al.*, 1999) properties. Quinoline derivatives have been used for their luminescent properties as organic light-emitting diode (OLED) materials (Montes *et al.*, 2006). Methyl acrylate is an ingredient used in many fragrances and decorative cosmetics (Bhatia *et al.*, 2007; Sharma, 2011). In view of the potential importance of the title compound its crystal structure is presented herein.

The asymmetric unit of the title compound contains the two independent molecules, A and B (Fig. 1). The dihedral angle between the mean plane of methyl ester unit ($C_{13}/C_{14}/O_2/O_3$, r.m.s deviation = -0.051 Å for A and -0.016 Å for B) and the chloroquinolin ring system ($C_1—C_9/N_1/C_{11}$, r.m.s deviation = 0.023 Å for A and -0.014 Å for B) is 63.5 (1)° in molecule A and 78.6 (1)° in molecule B. The main difference between the two independent molecules is reflected in the $O_1—C_{10}—C_{11}—C_{12}$ torsion angle which is -109.7 (2)° in molecule A and 10.6 (2)° in molecule B.

The methyl ester moiety adopts an extended conformation as reflected by the torsion angles for $C_{11}—C_{13}—C_{14}—O_3$ = 177.7 (2)° in A and 178.4 (1)° in B. The extended conformation is supported by the fact that the bond angles involving the carbonyl O atoms are invariably expanded (Dunitz & Schweizer, 1982). The significant difference in the bond lengths of the $C_{13}—O_3$ = 1.322 (3) Å (A) 1.331 (3) Å (B) versus $C_{14}—O_3$ = 1.447 (3) Å (A) and 1.447 (2) Å (B) can be attributed to a partial contribution from the $O^-—C=O^+—C$ resonance structure of the $O_2—C_{13}—O_3—C_{14}$ group (Merlino, 1971). This feature, commonly observed in the carboxylic ester group of these substituents in various compounds has been shown to give average values of 1.340 Å and 1.447 Å respectively for these bonds (Varghese *et al.*, 1986).


In the crystal molecule A and B are linked into pairs via bifurcated $O—H\cdots(N,Cl)$ hydrogen bonds (Fig. 2) and a weak $C—H\cdots O$ hydrogen bond links pairs of molecules into chains along [100].

S2. Experimental


A mixture of 2-chloroquinoline-3-carbaldehyde (0.1 g, 0.52 mmol), methyl acrylate (0.071 ml, 0.78 mmol), and DABCO (0.017 g, 0.15 mmol), was kept at room temperature for 7 days. After completion of the reaction (indicated by TLC), the reaction mixture was extracted with ethylacetate (3 times 15 ml). The combined organic layer subsequently washed with dil.HCl and dried over anhydrous Na_2SO_4 . The solvent was evaporated under reduced pressure. The crude product was obtained and purified by column chromatography eluting with 8% ethylacetate in hexane afforded the alcohol methyl 2-((2-chloroquinolin-3-yl)(hydroxy)methyl)acrylate as a colourless solid. X-ray quality crystals were obtained by slow evaporation of a solution of the title compound in ethylacetate.

S3. Refinement

Hydrogen atoms were positioned geometrically and allowed to ride on their parent atoms, with $C—H$ = 0.93–0.98 Å, $O—H$ = 0.82° and $U_{iso}(H)$ = 1.5 $U_{eq}(C)$ for methyl and hydroxyl H atoms and 1.2 $U_{eq}(C)$ for other H atoms.

**Figure 1**

The asymmetric unit of the title compound showing 30% probability displacement ellipsoids. H atoms are shown as spheres of arbitrary radius.

**Figure 2**

Part of the crystal structure with hydrogen bonds shown as dashed lines.

Methyl 2-[(2-chloroquinolin-3-yl)(hydroxy)methyl]acrylate

Crystal data

$C_{14}H_{12}ClNO_3$

$M_r = 277.70$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.2614 (4) \text{ \AA}$

$b = 11.0309 (4) \text{ \AA}$

$c = 13.8161 (6) \text{ \AA}$

$\alpha = 102.557 (2)^\circ$

$\beta = 100.646 (2)^\circ$

$\gamma = 103.704 (2)^\circ$

$V = 1296.29 (9) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 576$
 $D_x = 1.423 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 4551 reflections

$\theta = 2.0\text{--}25.0^\circ$
 $\mu = 0.30 \text{ mm}^{-1}$
 $T = 293 \text{ K}$
Block, colourless
 $0.35 \times 0.30 \times 0.25 \text{ mm}$

Data collection

Bruker SMART APEXII
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scans
14402 measured reflections
4466 independent reflections

3767 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.0^\circ$
 $h = -11 \rightarrow 10$
 $k = -12 \rightarrow 13$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.093$
 $S = 1.03$
4466 reflections
344 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0462P)^2 + 0.4048P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.29 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.0040 (11)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1A	0.14491 (5)	0.05724 (5)	0.40164 (4)	0.05417 (16)
O1A	0.36327 (14)	-0.23933 (11)	0.28015 (10)	0.0504 (3)
H1A	0.3051	-0.3109	0.2473	0.076*
O2A	0.05490 (17)	-0.36860 (13)	0.16879 (12)	0.0655 (4)
O3A	-0.03759 (16)	-0.26279 (14)	0.06395 (11)	0.0626 (4)
N1A	0.42615 (15)	0.19802 (13)	0.45982 (10)	0.0356 (3)
C1A	0.33952 (18)	0.08303 (15)	0.40781 (12)	0.0338 (4)
C2A	0.58039 (18)	0.22445 (15)	0.46705 (12)	0.0342 (4)
C3A	0.6764 (2)	0.35052 (17)	0.51704 (13)	0.0443 (4)
H3A	0.6352	0.4153	0.5445	0.053*

C4A	0.8297 (2)	0.37743 (19)	0.52515 (15)	0.0520 (5)
H4A	0.8929	0.4611	0.5583	0.062*
C5A	0.8944 (2)	0.2818 (2)	0.48466 (15)	0.0521 (5)
H5A	0.9999	0.3017	0.4921	0.062*
C6A	0.8037 (2)	0.16025 (19)	0.43468 (14)	0.0456 (4)
H6A	0.8474	0.0973	0.4073	0.055*
C7A	0.64369 (18)	0.12781 (16)	0.42366 (12)	0.0350 (4)
C8A	0.54251 (18)	0.00498 (16)	0.36909 (12)	0.0356 (4)
H8A	0.5817	-0.0604	0.3401	0.043*
C9A	0.38875 (18)	-0.02014 (15)	0.35779 (12)	0.0321 (3)
C10A	0.27692 (18)	-0.14936 (15)	0.29730 (12)	0.0352 (4)
H10A	0.2067	-0.1775	0.3385	0.042*
C11A	0.18355 (18)	-0.14703 (15)	0.19609 (12)	0.0368 (4)
C12A	0.2087 (2)	-0.05086 (19)	0.15361 (15)	0.0547 (5)
H12A	0.1496	-0.0596	0.0889	0.066*
H12B	0.2855	0.0259	0.1883	0.066*
C13A	0.0617 (2)	-0.27074 (17)	0.14275 (14)	0.0433 (4)
C14A	-0.1553 (3)	-0.3815 (2)	0.00547 (18)	0.0770 (7)
H14A	-0.2208	-0.3645	-0.0494	0.116*
H14B	-0.2151	-0.4134	0.0493	0.116*
H14C	-0.1082	-0.4451	-0.0223	0.116*
C11B	0.34205 (5)	0.35070 (5)	0.27784 (4)	0.05394 (16)
N1B	0.52758 (16)	0.22798 (12)	0.21855 (10)	0.0386 (3)
O1B	0.71157 (13)	0.63179 (11)	0.42717 (8)	0.0422 (3)
H1B	0.7063	0.7033	0.4560	0.063*
O2B	0.45974 (15)	0.56659 (12)	0.12824 (10)	0.0530 (3)
O3B	0.59711 (17)	0.75707 (12)	0.11953 (9)	0.0553 (4)
C1B	0.51958 (18)	0.34408 (15)	0.25490 (12)	0.0342 (4)
C2B	0.66394 (19)	0.21585 (15)	0.19986 (12)	0.0357 (4)
C3B	0.6778 (2)	0.09090 (17)	0.16220 (14)	0.0482 (5)
H3B	0.5944	0.0184	0.1500	0.058*
C4B	0.8116 (3)	0.07581 (18)	0.14373 (16)	0.0560 (5)
H4B	0.8196	-0.0072	0.1191	0.067*
C5B	0.9382 (3)	0.1835 (2)	0.16113 (18)	0.0607 (6)
H5B	1.0297	0.1717	0.1483	0.073*
C6B	0.9281 (2)	0.30528 (18)	0.19672 (16)	0.0532 (5)
H6B	1.0125	0.3764	0.2074	0.064*
C7B	0.79075 (19)	0.32456 (15)	0.21766 (13)	0.0369 (4)
C8B	0.77249 (18)	0.44774 (15)	0.25652 (12)	0.0364 (4)
H8B	0.8541	0.5215	0.2686	0.044*
C9B	0.63788 (17)	0.46076 (14)	0.27667 (11)	0.0316 (3)
C10B	0.62325 (18)	0.59300 (14)	0.32479 (12)	0.0328 (4)
H10B	0.5155	0.5844	0.3250	0.039*
C11B	0.67326 (18)	0.68846 (14)	0.26558 (12)	0.0335 (4)
C12B	0.7998 (2)	0.78554 (17)	0.29927 (15)	0.0470 (4)
H12C	0.8246	0.8416	0.2595	0.056*
H12D	0.8646	0.7982	0.3629	0.056*
C13B	0.5651 (2)	0.66278 (15)	0.16500 (13)	0.0375 (4)

C14B	0.4924 (3)	0.7379 (2)	0.02265 (16)	0.0765 (7)
H14D	0.5241	0.8104	-0.0041	0.115*
H14E	0.3909	0.7304	0.0321	0.115*
H14F	0.4924	0.6599	-0.0246	0.115*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1A	0.0291 (2)	0.0501 (3)	0.0697 (3)	0.00871 (19)	0.0111 (2)	-0.0073 (2)
O1A	0.0485 (7)	0.0334 (6)	0.0626 (8)	0.0161 (6)	0.0061 (6)	0.0011 (6)
O2A	0.0628 (9)	0.0374 (8)	0.0763 (10)	-0.0048 (6)	0.0039 (8)	0.0076 (7)
O3A	0.0484 (8)	0.0571 (9)	0.0545 (8)	-0.0067 (6)	-0.0094 (7)	0.0015 (7)
N1A	0.0342 (7)	0.0321 (7)	0.0354 (7)	0.0073 (6)	0.0065 (6)	0.0029 (6)
C1A	0.0290 (8)	0.0363 (9)	0.0329 (8)	0.0085 (7)	0.0059 (7)	0.0052 (7)
C2A	0.0332 (9)	0.0356 (9)	0.0300 (8)	0.0054 (7)	0.0042 (7)	0.0093 (7)
C3A	0.0472 (11)	0.0356 (9)	0.0436 (10)	0.0040 (8)	0.0095 (8)	0.0079 (8)
C4A	0.0425 (11)	0.0462 (11)	0.0518 (11)	-0.0086 (9)	0.0031 (9)	0.0129 (9)
C5A	0.0308 (9)	0.0623 (13)	0.0565 (12)	0.0009 (9)	0.0039 (8)	0.0225 (10)
C6A	0.0335 (9)	0.0544 (11)	0.0499 (11)	0.0135 (8)	0.0089 (8)	0.0166 (9)
C7A	0.0326 (8)	0.0394 (9)	0.0323 (8)	0.0097 (7)	0.0046 (7)	0.0122 (7)
C8A	0.0359 (9)	0.0357 (9)	0.0355 (9)	0.0129 (7)	0.0089 (7)	0.0075 (7)
C9A	0.0322 (8)	0.0316 (8)	0.0301 (8)	0.0085 (7)	0.0063 (6)	0.0057 (6)
C10A	0.0358 (9)	0.0288 (8)	0.0383 (9)	0.0085 (7)	0.0101 (7)	0.0042 (7)
C11A	0.0329 (9)	0.0328 (9)	0.0377 (9)	0.0039 (7)	0.0079 (7)	0.0024 (7)
C12A	0.0561 (12)	0.0461 (11)	0.0455 (11)	-0.0004 (9)	-0.0049 (9)	0.0104 (9)
C13A	0.0383 (10)	0.0398 (10)	0.0422 (10)	0.0030 (8)	0.0108 (8)	-0.0001 (8)
C14A	0.0527 (13)	0.0717 (15)	0.0651 (14)	-0.0098 (11)	-0.0082 (11)	-0.0158 (12)
Cl1B	0.0352 (2)	0.0458 (3)	0.0787 (4)	0.0054 (2)	0.0187 (2)	0.0156 (2)
N1B	0.0413 (8)	0.0290 (7)	0.0403 (8)	0.0030 (6)	0.0077 (6)	0.0087 (6)
O1B	0.0493 (7)	0.0388 (7)	0.0343 (6)	0.0170 (6)	0.0034 (5)	0.0026 (5)
O2B	0.0555 (8)	0.0369 (7)	0.0473 (7)	0.0006 (6)	-0.0073 (6)	0.0035 (6)
O3B	0.0787 (10)	0.0391 (7)	0.0404 (7)	0.0100 (7)	0.0008 (6)	0.0144 (6)
C1B	0.0328 (8)	0.0326 (9)	0.0348 (9)	0.0054 (7)	0.0058 (7)	0.0110 (7)
C2B	0.0450 (10)	0.0278 (8)	0.0326 (8)	0.0078 (7)	0.0078 (7)	0.0093 (7)
C3B	0.0625 (12)	0.0281 (9)	0.0502 (11)	0.0083 (8)	0.0157 (9)	0.0068 (8)
C4B	0.0755 (14)	0.0349 (10)	0.0615 (13)	0.0233 (10)	0.0231 (11)	0.0080 (9)
C5B	0.0591 (13)	0.0498 (12)	0.0796 (15)	0.0261 (10)	0.0275 (11)	0.0107 (11)
C6B	0.0438 (11)	0.0393 (10)	0.0747 (14)	0.0113 (8)	0.0193 (10)	0.0088 (9)
C7B	0.0401 (9)	0.0314 (8)	0.0385 (9)	0.0105 (7)	0.0092 (7)	0.0085 (7)
C8B	0.0337 (9)	0.0264 (8)	0.0430 (9)	0.0032 (7)	0.0061 (7)	0.0065 (7)
C9B	0.0330 (8)	0.0287 (8)	0.0307 (8)	0.0065 (7)	0.0044 (7)	0.0090 (6)
C10B	0.0308 (8)	0.0291 (8)	0.0348 (9)	0.0074 (6)	0.0047 (7)	0.0051 (7)
C11B	0.0350 (9)	0.0252 (8)	0.0367 (9)	0.0089 (7)	0.0063 (7)	0.0033 (7)
C12B	0.0444 (10)	0.0370 (10)	0.0510 (11)	0.0017 (8)	0.0039 (8)	0.0124 (8)
C13B	0.0450 (10)	0.0274 (8)	0.0376 (9)	0.0121 (8)	0.0072 (8)	0.0047 (7)
C14B	0.115 (2)	0.0619 (14)	0.0432 (12)	0.0268 (14)	-0.0072 (12)	0.0172 (10)

Geometric parameters (\AA , $\text{^{\circ}}$)

C11A—C1A	1.7396 (16)	C11B—C1B	1.7459 (17)
O1A—C10A	1.4228 (19)	N1B—C1B	1.295 (2)
O1A—H1A	0.8200	N1B—C2B	1.366 (2)
O2A—C13A	1.202 (2)	O1B—C10B	1.4130 (18)
O3A—C13A	1.322 (2)	O1B—H1B	0.8200
O3A—C14A	1.447 (2)	O2B—C13B	1.195 (2)
N1A—C1A	1.295 (2)	O3B—C13B	1.331 (2)
N1A—C2A	1.369 (2)	O3B—C14B	1.441 (2)
C1A—C9A	1.415 (2)	C1B—C9B	1.410 (2)
C2A—C3A	1.404 (2)	C2B—C7B	1.406 (2)
C2A—C7A	1.410 (2)	C2B—C3B	1.408 (2)
C3A—C4A	1.358 (3)	C3B—C4B	1.351 (3)
C3A—H3A	0.9300	C3B—H3B	0.9300
C4A—C5A	1.397 (3)	C4B—C5B	1.399 (3)
C4A—H4A	0.9300	C4B—H4B	0.9300
C5A—C6A	1.353 (3)	C5B—C6B	1.358 (3)
C5A—H5A	0.9300	C5B—H5B	0.9300
C6A—C7A	1.410 (2)	C6B—C7B	1.409 (2)
C6A—H6A	0.9300	C6B—H6B	0.9300
C7A—C8A	1.406 (2)	C7B—C8B	1.410 (2)
C8A—C9A	1.357 (2)	C8B—C9B	1.359 (2)
C8A—H8A	0.9300	C8B—H8B	0.9300
C9A—C10A	1.505 (2)	C9B—C10B	1.514 (2)
C10A—C11A	1.511 (2)	C10B—C11B	1.509 (2)
C10A—H10A	0.9800	C10B—H10B	0.9800
C11A—C12A	1.315 (3)	C11B—C12B	1.313 (2)
C11A—C13A	1.485 (2)	C11B—C13B	1.481 (2)
C12A—H12A	0.9300	C12B—H12C	0.9300
C12A—H12B	0.9300	C12B—H12D	0.9300
C14A—H14A	0.9600	C14B—H14D	0.9600
C14A—H14B	0.9600	C14B—H14E	0.9600
C14A—H14C	0.9600	C14B—H14F	0.9600
C10A—O1A—H1A	109.5	C1B—N1B—C2B	117.42 (14)
C13A—O3A—C14A	116.51 (17)	C10B—O1B—H1B	109.5
C1A—N1A—C2A	117.75 (14)	C13B—O3B—C14B	115.36 (15)
N1A—C1A—C9A	126.31 (15)	N1B—C1B—C9B	126.51 (15)
N1A—C1A—Cl1A	115.01 (12)	N1B—C1B—Cl1B	114.43 (12)
C9A—C1A—Cl1A	118.68 (12)	C9B—C1B—Cl1B	119.06 (12)
N1A—C2A—C3A	119.28 (15)	N1B—C2B—C7B	121.77 (14)
N1A—C2A—C7A	121.07 (14)	N1B—C2B—C3B	119.00 (15)
C3A—C2A—C7A	119.64 (15)	C7B—C2B—C3B	119.23 (16)
C4A—C3A—C2A	119.69 (18)	C4B—C3B—C2B	120.34 (18)
C4A—C3A—H3A	120.2	C4B—C3B—H3B	119.8
C2A—C3A—H3A	120.2	C2B—C3B—H3B	119.8
C3A—C4A—C5A	121.23 (17)	C3B—C4B—C5B	120.76 (17)

C3A—C4A—H4A	119.4	C3B—C4B—H4B	119.6
C5A—C4A—H4A	119.4	C5B—C4B—H4B	119.6
C6A—C5A—C4A	120.03 (17)	C6B—C5B—C4B	120.30 (19)
C6A—C5A—H5A	120.0	C6B—C5B—H5B	119.9
C4A—C5A—H5A	120.0	C4B—C5B—H5B	119.9
C5A—C6A—C7A	120.83 (18)	C5B—C6B—C7B	120.46 (18)
C5A—C6A—H6A	119.6	C5B—C6B—H6B	119.8
C7A—C6A—H6A	119.6	C7B—C6B—H6B	119.8
C8A—C7A—C6A	123.60 (16)	C2B—C7B—C8B	117.38 (15)
C8A—C7A—C2A	117.83 (15)	C2B—C7B—C6B	118.91 (15)
C6A—C7A—C2A	118.54 (15)	C8B—C7B—C6B	123.71 (16)
C9A—C8A—C7A	121.35 (15)	C9B—C8B—C7B	121.30 (15)
C9A—C8A—H8A	119.3	C9B—C8B—H8B	119.3
C7A—C8A—H8A	119.3	C7B—C8B—H8B	119.3
C8A—C9A—C1A	115.59 (14)	C8B—C9B—C1B	115.61 (14)
C8A—C9A—C10A	122.69 (14)	C8B—C9B—C10B	120.66 (14)
C1A—C9A—C10A	121.72 (14)	C1B—C9B—C10B	123.64 (14)
O1A—C10A—C9A	107.26 (13)	O1B—C10B—C11B	112.74 (13)
O1A—C10A—C11A	109.72 (13)	O1B—C10B—C9B	106.54 (12)
C9A—C10A—C11A	113.64 (13)	C11B—C10B—C9B	111.38 (13)
O1A—C10A—H10A	108.7	O1B—C10B—H10B	108.7
C9A—C10A—H10A	108.7	C11B—C10B—H10B	108.7
C11A—C10A—H10A	108.7	C9B—C10B—H10B	108.7
C12A—C11A—C13A	121.29 (16)	C12B—C11B—C13B	122.74 (16)
C12A—C11A—C10A	125.54 (15)	C12B—C11B—C10B	123.77 (15)
C13A—C11A—C10A	113.06 (14)	C13B—C11B—C10B	113.50 (13)
C11A—C12A—H12A	120.0	C11B—C12B—H12C	120.0
C11A—C12A—H12B	120.0	C11B—C12B—H12D	120.0
H12A—C12A—H12B	120.0	H12C—C12B—H12D	120.0
O2A—C13A—O3A	123.47 (17)	O2B—C13B—O3B	123.36 (15)
O2A—C13A—C11A	122.95 (17)	O2B—C13B—C11B	123.22 (15)
O3A—C13A—C11A	113.58 (16)	O3B—C13B—C11B	113.42 (14)
O3A—C14A—H14A	109.5	O3B—C14B—H14D	109.5
O3A—C14A—H14B	109.5	O3B—C14B—H14E	109.5
H14A—C14A—H14B	109.5	H14D—C14B—H14E	109.5
O3A—C14A—H14C	109.5	O3B—C14B—H14F	109.5
H14A—C14A—H14C	109.5	H14D—C14B—H14F	109.5
H14B—C14A—H14C	109.5	H14E—C14B—H14F	109.5
C2A—N1A—C1A—C9A	0.7 (2)	C2B—N1B—C1B—C9B	0.6 (2)
C2A—N1A—C1A—Cl1A	-179.69 (11)	C2B—N1B—C1B—Cl1B	-179.34 (11)
C1A—N1A—C2A—C3A	175.82 (15)	C1B—N1B—C2B—C7B	-0.9 (2)
C1A—N1A—C2A—C7A	-3.1 (2)	C1B—N1B—C2B—C3B	178.77 (15)
N1A—C2A—C3A—C4A	179.57 (16)	N1B—C2B—C3B—C4B	-179.50 (17)
C7A—C2A—C3A—C4A	-1.5 (2)	C7B—C2B—C3B—C4B	0.2 (3)
C2A—C3A—C4A—C5A	0.0 (3)	C2B—C3B—C4B—C5B	-0.3 (3)
C3A—C4A—C5A—C6A	1.2 (3)	C3B—C4B—C5B—C6B	-0.2 (3)
C4A—C5A—C6A—C7A	-0.7 (3)	C4B—C5B—C6B—C7B	0.7 (3)

C5A—C6A—C7A—C8A	177.30 (17)	N1B—C2B—C7B—C8B	0.4 (2)
C5A—C6A—C7A—C2A	-0.8 (3)	C3B—C2B—C7B—C8B	-179.26 (15)
N1A—C2A—C7A—C8A	2.6 (2)	N1B—C2B—C7B—C6B	-179.98 (16)
C3A—C2A—C7A—C8A	-176.31 (15)	C3B—C2B—C7B—C6B	0.3 (2)
N1A—C2A—C7A—C6A	-179.18 (15)	C5B—C6B—C7B—C2B	-0.8 (3)
C3A—C2A—C7A—C6A	1.9 (2)	C5B—C6B—C7B—C8B	178.76 (19)
C6A—C7A—C8A—C9A	-177.76 (16)	C2B—C7B—C8B—C9B	0.5 (2)
C2A—C7A—C8A—C9A	0.4 (2)	C6B—C7B—C8B—C9B	-179.11 (17)
C7A—C8A—C9A—C1A	-2.5 (2)	C7B—C8B—C9B—C1B	-0.8 (2)
C7A—C8A—C9A—C10A	178.19 (15)	C7B—C8B—C9B—C10B	176.02 (14)
N1A—C1A—C9A—C8A	2.1 (2)	N1B—C1B—C9B—C8B	0.3 (2)
C11A—C1A—C9A—C8A	-177.49 (12)	C11B—C1B—C9B—C8B	-179.82 (12)
N1A—C1A—C9A—C10A	-178.61 (16)	N1B—C1B—C9B—C10B	-176.42 (15)
C11A—C1A—C9A—C10A	1.8 (2)	C11B—C1B—C9B—C10B	3.5 (2)
C8A—C9A—C10A—O1A	14.0 (2)	C8B—C9B—C10B—O1B	-70.40 (18)
C1A—C9A—C10A—O1A	-165.25 (14)	C1B—C9B—C10B—O1B	106.14 (16)
C8A—C9A—C10A—C11A	-107.46 (17)	C8B—C9B—C10B—C11B	52.92 (19)
C1A—C9A—C10A—C11A	73.3 (2)	C1B—C9B—C10B—C11B	-130.54 (15)
O1A—C10A—C11A—C12A	-109.7 (2)	O1B—C10B—C11B—C12B	10.6 (2)
C9A—C10A—C11A—C12A	10.4 (2)	C9B—C10B—C11B—C12B	-109.15 (18)
O1A—C10A—C11A—C13A	66.52 (18)	O1B—C10B—C11B—C13B	-169.19 (12)
C9A—C10A—C11A—C13A	-173.42 (13)	C9B—C10B—C11B—C13B	71.10 (17)
C14A—O3A—C13A—O2A	-2.5 (3)	C14B—O3B—C13B—O2B	1.8 (3)
C14A—O3A—C13A—C11A	177.18 (17)	C14B—O3B—C13B—C11B	-178.39 (16)
C12A—C11A—C13A—O2A	164.7 (2)	C12B—C11B—C13B—O2B	170.78 (18)
C10A—C11A—C13A—O2A	-11.7 (2)	C10B—C11B—C13B—O2B	-9.5 (2)
C12A—C11A—C13A—O3A	-15.0 (3)	C12B—C11B—C13B—O3B	-9.0 (2)
C10A—C11A—C13A—O3A	168.60 (15)	C10B—C11B—C13B—O3B	170.72 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1A—H1A···O2A	0.82	2.24	2.8372 (19)	130
O1B—H1B···C11A ⁱ	0.82	2.79	3.5040 (12)	147
O1B—H1B···N1A ⁱ	0.82	2.16	2.8609 (17)	144
C5A—H5A···O1B ⁱⁱ	0.93	2.56	3.451 (2)	162

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+1, -z+1$.