

2-[(2-Chlorophenyl)(hydroxy)methyl]-phenol

G. Prakasha,^a Jerry P. Jasinski,^{b*} Jared S. Brown,^b
H. S. Yathirajan^a and D. K. Ravishankara^c

^aDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, ^bDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, and ^cSri Mahadewhara Government First Grade College, (Affiliated to University of Mysore), Kollegal 571 440, India
Correspondence e-mail: jjjasinski@keene.edu


Received 4 July 2013; accepted 5 July 2013

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.065; wR factor = 0.213; data-to-parameter ratio = 15.7.

In the title compound, $\text{C}_{13}\text{H}_{11}\text{ClO}_2$, the dihedral angle between the mean planes of the 2-chlorophenyl and phenol rings is $87.4(9)^\circ$. The methyl hydroxy group lies nearly perpendicular to the plane of its attached benzene ring [$\text{O}-\text{C}-\text{C}-\text{C}$ torsion angle = $84.3(3)^\circ$]. The two hydroxy groups lie on the same side of the molecule and are in a slightly twisted gauche conformation [$\text{O}-\text{C}-\text{C}-\text{O}$ torsion angle = $77.1(8)^\circ$] to each other. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds between nearby methylhydroxy groups form dimers in alternating pairs aligned diagonally along the b axis. A view along the c axis reveals a hexameric aggregate mediated by a ring of six $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds generating an $R_6^6(12)$ motif loop.

Related literature

For general background to the use of benzhydrols, see: Ohkuma *et al.* (2000). For the use of the title compound in the perfume and pharmaceutical industries, see: Meguro *et al.* (1985). For related diphenylmethanol structures, see: Betz *et al.* (2011); Ferguson *et al.* (1995); Siddaraju *et al.* (2010).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{11}\text{ClO}_2$

$M_r = 234.67$

Trigonal, $R\bar{3}$
 $a = 23.4627(8)\text{ \AA}$
 $c = 11.3722(4)\text{ \AA}$
 $V = 5421.6(4)\text{ \AA}^3$
 $Z = 18$

Cu $K\alpha$ radiation
 $\mu = 2.66\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.46 \times 0.38 \times 0.24\text{ mm}$

Data collection

Agilent Xcalibur (Eos, Gemini) diffractometer
Absorption correction: multi-scan (*CrysAlis PRO* and *CrysAlis RED*; Agilent, 2012)
 $T_{\min} = 0.517$, $T_{\max} = 1.000$

11532 measured reflections
2364 independent reflections
2055 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.213$
 $S = 1.07$
2364 reflections
151 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.62\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots O1 ⁱ	0.84	1.84	2.656 (2)	163
Symmetry code: (i) $y + \frac{1}{3}, -x + y + \frac{2}{3}, -z + \frac{5}{3}$.				

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Agilent, 2012); program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL2012* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *OLEX2*.

GP thanks the UOM for research facilities to complete MSc dissertation work. JPJ acknowledges the NSF-MRI program (grant No. CHE-1039027) for funds to purchase the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5343).

References

- Agilent (2012). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies, Yarnton, England.
- Betz, R., Gerber, T., Hosten, E., Siddaraju, B. P., Yathirajan, H. S. & Ramesha, A. R. (2011). *Acta Cryst.* **E67**, o3302.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Ferguson, G., Carroll, C. D., Glidewell, C., Zakaria, C. M. & Lough, A. J. (1995). *Acta Cryst.* **B51**, 367–377.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Meguro, K., Aizawa, M., Sohda, T., Kawamatsu, Y. & Nagaoka, A. (1985). *Chem. Pharm. Bull.* **33**, 3787–3797.
- Ohkuma, T., Koizumi, M., Ikehira, H., Yokozawa, T. & Noyori, R. (2000). *Org. Lett.* **2**, 659–662.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siddaraju, B. P., Yathirajan, H. S., Narayana, B., Ng, S. W. & Tiekkink, E. R. T. (2010). *Acta Cryst.* **E66**, o2136.

supporting information

Acta Cryst. (2013). E69, o1240 [doi:10.1107/S1600536813018667]

2-[(2-Chlorophenyl)(hydroxy)methyl]phenol

G. Prakasha, Jerry P. Jasinski, Jared S. Brown, H. S. Yathirajan and D. K. Ravishankara

S1. Comment

Optically active diarylmethanols are crucial structural elements in many physiologically or/and biologically active molecules such as in the antihistamines (R)-orphenadrine and (S)-neobenodine. Benzhydrols are widely used as intermediates for the synthesis of pharmaceuticals (Ohkuma *et al.*, 2000), including drugs such as diphenhydramine, orphenadrine, diphenidol and phenyltoloxamine. The crystal structures and hydrogen bonding in some diarylmethanols have been reported (Ferguson *et al.*, 1995). The title compound, 2-[(2-chlorophenyl) hydroxymethyl]phenol, (I), is a derivative of diphenylmethanol and it is used in the perfume and pharmaceutical industries (Meguro *et al.*, 1985). Recently, the crystal structure of (2-methylphenyl) (phenyl)methanol (Siddaraju *et al.*, 2010) and 2-(2-benzylphenyl) propan-2-ol (Betz *et al.*, 2011) were reported. In view of the importance of diarylmethanols, this paper reports the crystal structure of the title compound, C₁₃H₁₁O₂Cl, (I).


In (I), the dihedral angle between the mean planes of the 2-chlorophenyl and phenol rings is 87.4 (9) $^{\circ}$ (Fig. 1). The methyl hydroxy group lies nearly perpendicular to the plane of its attached benzene ring [O1/C1/C2/C3 torsion angle = 84.3 (3) $^{\circ}$]. The two hydroxy groups lie on the same side of the molecule and are in a slightly twisted gauche conformation [O1/C1/C3/O2 angle = 77.1 (8) $^{\circ}$] to each other. In the crystal O—H \cdots O hydrogen bonds between nearby methylhydroxy groups form dimers in alternating pairs aligned diagonally along the *b* axis and contribute to packing stability (Fig. 2). A view along the *c* axis reveals a hexameric aggregate mediated by a ring of six O—H \cdots O hydrogen bonds generating an R₆(12) motif loop (Fig. 3).

S2. Experimental


The title compound was obtained as a gift sample from R. L. Fine Chem, Bengaluru, India. X-ray quality crystals were obtained from benzene solution by slow evaporation. (m.p.: 368–373 K).

S3. Refinement


H2 was located by a difference map and refined isotropically. All of the remaining H atoms were placed in their calculated positions and then refined using the riding model with Atom—H lengths of 0.95 Å or 1.00 (CH), or 0.84 $^{\circ}$ (OH). Isotropic displacement parameters for these atoms were set to 1.2 (CH) or 1.5 (OH) times *U*_{eq} of the parent atom. Idealised tetrahedral OH refined as rotating group: O1(H1).

**Figure 1**

Molecular structure of the title compound showing the atom labeling scheme and 30% probability displacement ellipsoids.

**Figure 2**

Packing diagram of the title compound viewed along the *a* axis. Dashed lines indicate O1—H1···O1 hydrogen bonds between nearby methylhydroxy groups which form dimers in alternating pairs aligned diagonally along the *b* axis. H atoms not involved in hydrogen bonding have been deleted for clarity.

**Figure 3**

Packing of the molecules along c axis displaying a hexameric aggregate mediated by a ring of six $\text{O}—\text{H}\cdots\text{O}$ hydrogen bonds generating an $\text{R}_6^6(12)$ motif loop.

2-[(2-Chlorophenyl)(hydroxy)methyl]phenol

Crystal data

$\text{C}_{13}\text{H}_{11}\text{ClO}_2$
 $M_r = 234.67$
Trigonal, $R\bar{3}$
 $a = 23.4627 (8) \text{ \AA}$
 $c = 11.3722 (4) \text{ \AA}$
 $V = 5421.6 (4) \text{ \AA}^3$
 $Z = 18$
 $F(000) = 2196$

$D_x = 1.294 \text{ Mg m}^{-3}$
Cu $K\alpha$ radiation, $\lambda = 1.5418 \text{ \AA}$
Cell parameters from 3836 reflections
 $\theta = 3.8\text{--}72.3^\circ$
 $\mu = 2.66 \text{ mm}^{-1}$
 $T = 173 \text{ K}$
Irregular, colourless
 $0.46 \times 0.38 \times 0.24 \text{ mm}$

Data collection

Agilent Xcalibur (Eos, Gemini)
diffractometer
Radiation source: Enhance (Cu) X-ray Source
Graphite monochromator
Detector resolution: 16.0416 pixels mm^{-1}
 ω scans
Absorption correction: multi-scan
(*CrysAlis PRO* and *CrysAlis RED*; Agilent,
2012)

$T_{\min} = 0.517, T_{\max} = 1.000$
11532 measured reflections
2364 independent reflections
2055 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$
 $\theta_{\max} = 72.5^\circ, \theta_{\min} = 3.8^\circ$
 $h = -28\text{--}25$
 $k = -22\text{--}28$
 $l = -12\text{--}13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.213$
 $S = 1.07$
2364 reflections
151 parameters
0 restraints

Primary atom site location: structure-invariant
direct methods
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.1357P)^2 + 8.799P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.40 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.62 \text{ e \AA}^{-3}$

Extinction correction: *SHELXL2012* (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.00038 (12)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.66585 (5)	0.59567 (4)	0.59620 (10)	0.0780 (4)
O1	0.64040 (8)	0.41759 (8)	0.77861 (16)	0.0393 (5)
H1	0.6760	0.4335	0.8159	0.059*
O2	0.56637 (10)	0.49236 (11)	0.91391 (16)	0.0436 (5)
H2	0.590 (2)	0.532 (2)	0.888 (4)	0.068 (12)*
C1	0.63717 (10)	0.46886 (11)	0.71725 (19)	0.0295 (5)
H1A	0.6573	0.5090	0.7683	0.035*
C2	0.56447 (11)	0.44658 (11)	0.7019 (2)	0.0310 (5)
C3	0.53051 (12)	0.45681 (12)	0.7923 (2)	0.0370 (6)
C4	0.46358 (14)	0.43543 (14)	0.7772 (3)	0.0480 (7)
H4	0.4400	0.4422	0.8384	0.058*
C5	0.43171 (13)	0.40498 (16)	0.6757 (3)	0.0541 (8)
H5	0.3865	0.3918	0.6661	0.065*
C6	0.46467 (14)	0.39338 (17)	0.5873 (3)	0.0558 (8)
H6	0.4421	0.3711	0.5177	0.067*
C7	0.53144 (13)	0.41460 (14)	0.6004 (2)	0.0446 (7)
H7	0.5544	0.4070	0.5392	0.054*
C8	0.67533 (10)	0.48591 (12)	0.6030 (2)	0.0329 (5)
C9	0.69119 (12)	0.54381 (13)	0.5431 (2)	0.0437 (7)
C10	0.73000 (15)	0.5615 (2)	0.4418 (3)	0.0679 (11)
H10	0.7409	0.6013	0.4021	0.081*
C11	0.75243 (16)	0.5216 (3)	0.3997 (3)	0.0783 (14)
H11	0.7792	0.5339	0.3312	0.094*
C12	0.73624 (15)	0.4639 (2)	0.4566 (3)	0.0695 (12)
H12	0.7513	0.4360	0.4264	0.083*
C13	0.69824 (13)	0.44599 (15)	0.5574 (2)	0.0458 (7)
H13	0.6876	0.4060	0.5961	0.055*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0763 (7)	0.0518 (5)	0.1047 (9)	0.0312 (4)	-0.0183 (5)	0.0120 (4)
O1	0.0278 (8)	0.0368 (9)	0.0499 (11)	0.0137 (7)	-0.0014 (7)	0.0132 (7)
O2	0.0534 (11)	0.0509 (12)	0.0340 (9)	0.0318 (10)	0.0080 (8)	-0.0075 (8)
C1	0.0277 (11)	0.0279 (10)	0.0316 (11)	0.0130 (8)	-0.0020 (8)	-0.0008 (8)
C2	0.0282 (11)	0.0317 (11)	0.0341 (11)	0.0158 (9)	0.0014 (8)	0.0026 (9)

C3	0.0403 (13)	0.0361 (12)	0.0416 (13)	0.0243 (10)	0.0078 (10)	0.0066 (10)
C4	0.0406 (14)	0.0513 (16)	0.0607 (17)	0.0294 (12)	0.0158 (12)	0.0136 (13)
C5	0.0283 (12)	0.0592 (18)	0.073 (2)	0.0208 (12)	0.0050 (12)	0.0138 (15)
C6	0.0332 (14)	0.0679 (19)	0.0545 (17)	0.0164 (13)	-0.0109 (12)	-0.0024 (14)
C7	0.0318 (12)	0.0575 (16)	0.0386 (13)	0.0179 (11)	-0.0008 (10)	-0.0053 (11)
C8	0.0229 (10)	0.0350 (12)	0.0339 (12)	0.0093 (9)	-0.0032 (8)	-0.0030 (9)
C9	0.0319 (12)	0.0434 (14)	0.0393 (13)	0.0065 (10)	-0.0076 (10)	0.0084 (10)
C10	0.0386 (15)	0.083 (2)	0.0433 (16)	0.0008 (16)	-0.0057 (13)	0.0225 (16)
C11	0.0346 (15)	0.135 (4)	0.0338 (15)	0.0182 (19)	0.0038 (12)	0.0009 (19)
C12	0.0385 (15)	0.118 (3)	0.0506 (18)	0.0378 (18)	-0.0045 (13)	-0.032 (2)
C13	0.0336 (12)	0.0563 (16)	0.0476 (14)	0.0226 (12)	-0.0031 (10)	-0.0144 (12)

Geometric parameters (\AA , $^{\circ}$)

C11—C9	1.710 (3)	C5—C6	1.376 (5)
O1—H1	0.8400	C6—H6	0.9500
O1—C1	1.425 (3)	C6—C7	1.394 (4)
O2—H2	0.87 (5)	C7—H7	0.9500
O2—C3	1.617 (3)	C8—C9	1.394 (4)
C1—H1A	1.0000	C8—C13	1.392 (4)
C1—C2	1.524 (3)	C9—C10	1.396 (4)
C1—C8	1.514 (3)	C10—H10	0.9500
C2—C3	1.393 (3)	C10—C11	1.369 (6)
C2—C7	1.383 (4)	C11—H11	0.9500
C3—C4	1.400 (4)	C11—C12	1.372 (7)
C4—H4	0.9500	C12—H12	0.9500
C4—C5	1.367 (5)	C12—C13	1.382 (4)
C5—H5	0.9500	C13—H13	0.9500
C1—O1—H1	109.5	C7—C6—H6	120.2
C3—O2—H2	99 (3)	C2—C7—C6	120.5 (3)
O1—C1—H1A	108.1	C2—C7—H7	119.7
O1—C1—C2	106.83 (17)	C6—C7—H7	119.7
O1—C1—C8	111.67 (19)	C9—C8—C1	120.8 (2)
C2—C1—H1A	108.1	C13—C8—C1	121.2 (2)
C8—C1—H1A	108.1	C13—C8—C9	118.0 (2)
C8—C1—C2	113.87 (18)	C8—C9—C11	120.2 (2)
C3—C2—C1	119.7 (2)	C8—C9—C10	120.6 (3)
C7—C2—C1	120.6 (2)	C10—C9—C11	119.2 (3)
C7—C2—C3	119.7 (2)	C9—C10—H10	119.9
C2—C3—O2	121.7 (2)	C11—C10—C9	120.1 (3)
C2—C3—C4	118.9 (2)	C11—C10—H10	119.9
C4—C3—O2	119.4 (2)	C10—C11—H11	120.1
C3—C4—H4	119.6	C10—C11—C12	119.9 (3)
C5—C4—C3	120.9 (3)	C12—C11—H11	120.1
C5—C4—H4	119.6	C11—C12—H12	119.7
C4—C5—H5	119.8	C11—C12—C13	120.6 (4)
C4—C5—C6	120.4 (2)	C13—C12—H12	119.7

C6—C5—H5	119.8	C8—C13—H13	119.6
C5—C6—H6	120.2	C12—C13—C8	120.7 (3)
C5—C6—C7	119.6 (3)	C12—C13—H13	119.6
Cl1—C9—C10—C11	178.1 (2)	C3—C2—C7—C6	0.9 (4)
O1—C1—C2—C3	84.3 (3)	C3—C4—C5—C6	1.6 (5)
O1—C1—C2—C7	−93.8 (3)	C4—C5—C6—C7	−1.8 (5)
O1—C1—C8—C9	−165.2 (2)	C5—C6—C7—C2	0.6 (5)
O1—C1—C8—C13	11.7 (3)	C7—C2—C3—O2	178.5 (2)
O2—C3—C4—C5	−179.8 (2)	C7—C2—C3—C4	−1.2 (4)
C1—C2—C3—O2	0.4 (3)	C8—C1—C2—C3	−151.9 (2)
C1—C2—C3—C4	−179.3 (2)	C8—C1—C2—C7	29.9 (3)
C1—C2—C7—C6	179.0 (3)	C8—C9—C10—C11	0.6 (4)
C1—C8—C9—Cl1	−1.7 (3)	C9—C8—C13—C12	0.8 (4)
C1—C8—C9—C10	175.8 (2)	C9—C10—C11—C12	0.6 (5)
C1—C8—C13—C12	−176.3 (2)	C10—C11—C12—C13	−1.0 (5)
C2—C1—C8—C9	73.7 (3)	C11—C12—C13—C8	0.3 (4)
C2—C1—C8—C13	−109.4 (2)	C13—C8—C9—Cl1	−178.72 (19)
C2—C3—C4—C5	0.0 (4)	C13—C8—C9—C10	−1.2 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···O1 ⁱ	0.84	1.84	2.656 (2)	163

Symmetry code: (i) $y+1/3, -x+y+2/3, -z+5/3$.