

Amitraz

Sangjin Lee,^a Tae Ho Kim,^{a*} Yong Woon Shin,^b Youngeun Jeon^a and Jineun Kim^{a*}

^aDepartment of Chemistry and Research Institute of Natural Sciences, Gyeongsang National University, Jinju 660-701, Republic of Korea, and ^bTest and Analytical Laboratory, Korea Food and Drug Administration, 123-7 Yongdang-dong, Busan 608-829, Republic of Korea
Correspondence e-mail: jekim@gnu.ac.kr, thkim@gnu.ac.kr


Received 8 July 2013; accepted 17 July 2013

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.062; wR factor = 0.182; data-to-parameter ratio = 16.2.

In the asymmetric unit of the title compound [systematic name: *N'*-(2,4-dimethylphenyl)-*N*-(2,4-dimethylphenyl)-carboximidoyl]-*N*-methylmethanimidamide], C₁₉H₂₃N₃, which is a formamidine pesticide, there are two independent and conformationally similar molecules, with the dihedral angle between the mean planes of the 2,4-dimethylbenzene rings in each molecule being 41.63 (6) and 42.09 (5)°. The crystal structure is stabilized by a C–H···N hydrogen bond, as well as weak intermolecular C–H···π and π–π interactions [ring centroid separation = 3.7409 (15) Å], giving one-dimensional chains extending down the b direction.

Related literature

For the toxicity and insecticidal properties of the title compound, see: Del Pino *et al.* (2013); Hollingworth (1976). For a related crystal structure, see: Peoples *et al.* (2012).



Experimental

Crystal data

C₁₉H₂₃N₃
 $M_r = 293.40$

Monoclinic, $P2_1/c$
 $a = 11.9362(11)$ Å

Data collection

Bruker APEXII CCD-dectecto diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2006)
 $R_{\text{int}} = 0.058$
 $T_{\min} = 0.973$, $T_{\max} = 0.973$

18091 measured reflections
6593 independent reflections
3615 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.058$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.062$
 $wR(F^2) = 0.182$
 $S = 0.98$
6593 reflections

407 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.26$ e Å⁻³
 $\Delta\rho_{\min} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and *Cg4* are the centroids of the C2–C8 and C31–C38 rings.

D–H···A	D–H	H···A	D···A	D–H···A
C29–H29C···N3 ⁱ	0.98	2.49	3.334 (4)	144
C17–H17A···Cg4 ⁱ	0.98	2.92	3.806 (3)	151
C24–H24A···Cg1 ⁱⁱ	0.98	2.85	3.828 (3)	175
C33–H33B···Cg4 ⁱⁱⁱ	0.98	2.88	3.625 (3)	134
Symmetry codes:	(i)	$-x, y - \frac{1}{2}, -z + \frac{3}{2}$	(ii)	$-x, y + \frac{1}{2}, -z + \frac{3}{2}$
		$-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$		(iii)

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2006); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (grant No. 2012R1A1B3003337).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2270).

References

- Bruker (2006). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Del Pino, J., Martinez, M. A., Castellano, V., Ramos, E., Martinez-Larranaga, M. R. & Anadon, A. (2013). *Toxicology*, **308**, 88–95.
- Hollingworth, R. M. (1976). *Environ. Health Perspect.* **14**, 57–69.
- Peoples, B. C., De la Vega, G., Valdebenito, C., Quijada, R., Ibañez, A., Valderrama, M. & Rojas, R. (2012). *J. Organomet. Chem.* **790**, 147–153.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2013). E69, o1300 [doi:10.1107/S1600536813019764]

Amitraz

Sangjin Lee, Tae Ho Kim, Yong Woon Shin, Youngeun Jeon and Jineun Kim

S1. Comment

The title compound ($C_{19}H_{23}N_3$) (common name amitraz) is one of the most widely used formamidine acaricide/insecticide in veterinary medicine and agriculture for use on ectoparasites (Del Pino *et al.*, 2013; Hollingworth, 1976). However, until now its crystal structure has not been reported.

In this compound (Scheme 1, Fig. 1), there are two independent and conformationally similar molecules (*A* and *B*) in the asymmetric unit, with the dihedral angle between the mean planes of the 2,4-dimethylphenyl rings in each of 41.63 (6) and 42.09 (5) $^{\circ}$, respectively. All bond lengths and bond angles are normal and comparable to those observed in the crystal structures of a similar compound (Peoples *et al.*, 2012).


In the crystal structure (Fig. 2), an intermolecular C—H \cdots N hydrogen bond between a methyl C29 H atom and the N3 acceptor of the *B* molecule, as well as weak intermolecular C—H \cdots π interactions (Table 1), are present. There is also a π — π interaction between one of the *A*-molecule 2,4-dimethylphenyl rings (defined by atoms C2–C8) [ring centroid separation 3.7409 (15) Å].

S2. Experimental


The title compound was purchased from the Dr Ehrenstorfer GmbH Company. Slow evaporation of a solution in CH_2Cl_2 gave single crystals suitable for X-ray analysis.

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H = 0.95 Å, $U_{iso} = 1.2U_{eq}(C)$ for Csp^2 —H and C—H = 0.98 Å, $U_{iso} = 1.5U_{eq}(C)$ for CH_3 groups.

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as small spheres of arbitrary radius.

**Figure 2**

Crystal packing of the title compound in the unit cell, with weak intermolecular C—H···N hydrogen bonds shown as dashed lines.

N'-(2,4-Dimethylphenyl)-*N*-[*N*-(2,4-dimethylphenyl)carboximidoyl]-*N*-methylmethanimidamide

Crystal data

$C_{19}H_{23}N_3$
 $M_r = 293.40$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 11.9362 (11)$ Å
 $b = 7.5110 (7)$ Å
 $c = 37.514 (3)$ Å
 $\beta = 91.650 (2)^\circ$
 $V = 3361.9 (5)$ Å³
 $Z = 8$

$F(000) = 1264$
 $D_x = 1.159 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2173 reflections
 $\theta = 2.7\text{--}23.1^\circ$
 $\mu = 0.07 \text{ mm}^{-1}$
 $T = 173$ K
Block, colourless
 $0.40 \times 0.40 \times 0.40$ mm

Data collection

Bruker APEXII CCD-dectecto
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2006)
 $T_{\min} = 0.973$, $T_{\max} = 0.973$

18091 measured reflections
6593 independent reflections
3615 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.058$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.1^\circ$
 $h = -14 \rightarrow 14$
 $k = -9 \rightarrow 9$
 $l = -46 \rightarrow 36$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.062$
 $wR(F^2) = 0.182$
 $S = 0.98$
6593 reflections
407 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0907P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.22503 (17)	0.8667 (3)	0.54029 (6)	0.0318 (5)
N2	0.07334 (17)	0.8469 (3)	0.57777 (5)	0.0303 (5)
N3	-0.06461 (17)	0.8046 (3)	0.61902 (5)	0.0316 (5)
N4	0.02203 (18)	0.7540 (3)	0.88836 (6)	0.0349 (6)
N5	0.16767 (17)	0.7179 (3)	0.84902 (6)	0.0335 (6)
N6	0.30082 (18)	0.6971 (3)	0.80512 (6)	0.0342 (6)
C1	0.3145 (2)	0.6882 (4)	0.48038 (7)	0.0393 (7)
H1A	0.2808	0.7846	0.4661	0.059*
H1B	0.2551	0.6175	0.4910	0.059*
H1C	0.3591	0.6118	0.4651	0.059*
C2	0.3887 (2)	0.7661 (3)	0.50947 (7)	0.0306 (6)
C3	0.5052 (2)	0.7534 (3)	0.50820 (7)	0.0344 (7)
H3	0.5367	0.6911	0.4889	0.041*
C4	0.5770 (2)	0.8273 (3)	0.53382 (7)	0.0326 (7)
C5	0.7027 (2)	0.8131 (4)	0.53071 (8)	0.0440 (8)
H5A	0.7231	0.8379	0.5061	0.066*

H5B	0.7271	0.6927	0.5373	0.066*
H5C	0.7394	0.8997	0.5467	0.066*
C6	0.5292 (2)	0.9178 (3)	0.56183 (7)	0.0349 (7)
H6	0.5763	0.9703	0.5798	0.042*
C7	0.4140 (2)	0.9327 (3)	0.56400 (7)	0.0322 (6)
H7	0.3829	0.9970	0.5831	0.039*
C8	0.3435 (2)	0.8542 (3)	0.53838 (7)	0.0288 (6)
C9	0.1853 (2)	0.8345 (3)	0.57073 (7)	0.0304 (6)
H9	0.2356	0.8002	0.5896	0.036*
C10	-0.0063 (2)	0.9115 (4)	0.55032 (7)	0.0377 (7)
H10A	-0.0510	0.8115	0.5410	0.057*
H10B	0.0348	0.9660	0.5309	0.057*
H10C	-0.0559	1.0002	0.5607	0.057*
C11	0.0382 (2)	0.8003 (3)	0.61073 (7)	0.0317 (6)
H11	0.0924	0.7631	0.6282	0.038*
C12	-0.0910 (2)	0.7724 (3)	0.65503 (7)	0.0288 (6)
C13	-0.1938 (2)	0.6914 (3)	0.66229 (7)	0.0320 (6)
C14	-0.2710 (2)	0.6305 (4)	0.63229 (7)	0.0417 (7)
H14A	-0.3297	0.5543	0.6418	0.063*
H14B	-0.2281	0.5631	0.6149	0.063*
H14C	-0.3054	0.7343	0.6206	0.063*
C15	-0.2228 (2)	0.6670 (4)	0.69766 (7)	0.0354 (7)
H15	-0.2925	0.6122	0.7025	0.042*
C16	-0.1540 (2)	0.7193 (4)	0.72608 (7)	0.0372 (7)
C17	-0.1871 (3)	0.6882 (4)	0.76443 (7)	0.0497 (8)
H17A	-0.2529	0.6102	0.7647	0.075*
H17B	-0.2050	0.8024	0.7755	0.075*
H17C	-0.1248	0.6319	0.7777	0.075*
C18	-0.0534 (2)	0.8020 (4)	0.71861 (7)	0.0393 (7)
H18	-0.0050	0.8402	0.7377	0.047*
C19	-0.0226 (2)	0.8295 (4)	0.68357 (7)	0.0366 (7)
H19	0.0460	0.8881	0.6790	0.044*
C20	-0.0414 (3)	0.9162 (4)	0.95372 (7)	0.0493 (8)
H20A	-0.0774	0.9872	0.9721	0.074*
H20B	0.0204	0.9843	0.9438	0.074*
H20C	-0.0121	0.8058	0.9644	0.074*
C21	-0.1258 (2)	0.8719 (3)	0.92448 (7)	0.0325 (6)
C22	-0.2386 (2)	0.9098 (4)	0.92838 (7)	0.0367 (7)
H22	-0.2612	0.9612	0.9502	0.044*
C23	-0.3199 (2)	0.8764 (4)	0.90212 (7)	0.0345 (7)
C24	-0.4417 (2)	0.9186 (4)	0.90765 (8)	0.0463 (8)
H24A	-0.4476	1.0309	0.9208	0.069*
H24B	-0.4757	0.8226	0.9213	0.069*
H24C	-0.4809	0.9297	0.8844	0.069*
C25	-0.2856 (2)	0.7988 (4)	0.87098 (7)	0.0395 (7)
H25	-0.3391	0.7734	0.8524	0.047*
C26	-0.1744 (2)	0.7576 (4)	0.86650 (7)	0.0386 (7)
H26	-0.1530	0.7006	0.8451	0.046*

C27	-0.0928 (2)	0.7972 (4)	0.89247 (7)	0.0322 (6)
C28	0.0609 (2)	0.7700 (4)	0.85751 (7)	0.0343 (7)
H28	0.0142	0.8205	0.8393	0.041*
C29	0.2435 (2)	0.6444 (4)	0.87668 (7)	0.0399 (7)
H29A	0.2636	0.7378	0.8940	0.060*
H29B	0.3114	0.5997	0.8657	0.060*
H29C	0.2060	0.5466	0.8889	0.060*
C30	0.2031 (2)	0.7375 (3)	0.81483 (7)	0.0323 (6)
H30	0.1516	0.7841	0.7974	0.039*
C31	0.3252 (2)	0.7255 (3)	0.76890 (7)	0.0311 (6)
C32	0.4296 (2)	0.8009 (3)	0.76116 (7)	0.0327 (7)
C33	0.5113 (2)	0.8437 (4)	0.79118 (7)	0.0435 (8)
H33A	0.4770	0.9280	0.8075	0.065*
H33B	0.5790	0.8968	0.7815	0.065*
H33C	0.5313	0.7342	0.8041	0.065*
C34	0.4541 (2)	0.8329 (4)	0.72587 (8)	0.0398 (7)
H34	0.5238	0.8867	0.7207	0.048*
C35	0.3809 (3)	0.7896 (4)	0.69786 (8)	0.0422 (8)
C36	0.4117 (3)	0.8261 (5)	0.65963 (8)	0.0645 (10)
H36A	0.3974	0.7195	0.6452	0.097*
H36B	0.4914	0.8572	0.6588	0.097*
H36C	0.3664	0.9251	0.6502	0.097*
C37	0.2804 (2)	0.7109 (4)	0.70560 (8)	0.0420 (8)
H37	0.2298	0.6779	0.6867	0.050*
C38	0.2520 (2)	0.6791 (4)	0.74065 (7)	0.0384 (7)
H38	0.1821	0.6252	0.7455	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0276 (13)	0.0354 (13)	0.0327 (13)	-0.0026 (10)	0.0062 (10)	0.0000 (10)
N2	0.0257 (12)	0.0392 (13)	0.0261 (13)	0.0016 (10)	0.0039 (10)	0.0012 (10)
N3	0.0300 (13)	0.0350 (13)	0.0300 (13)	-0.0006 (10)	0.0066 (10)	0.0023 (10)
N4	0.0367 (14)	0.0377 (13)	0.0304 (13)	-0.0014 (11)	0.0025 (11)	0.0006 (11)
N5	0.0275 (13)	0.0385 (13)	0.0344 (14)	0.0012 (10)	-0.0009 (10)	0.0061 (11)
N6	0.0298 (13)	0.0364 (14)	0.0366 (14)	-0.0012 (10)	0.0025 (11)	0.0031 (11)
C1	0.0416 (17)	0.0466 (18)	0.0300 (16)	-0.0029 (14)	0.0044 (13)	-0.0033 (13)
C2	0.0316 (16)	0.0324 (15)	0.0280 (15)	-0.0010 (12)	0.0048 (12)	0.0042 (12)
C3	0.0379 (17)	0.0352 (16)	0.0307 (16)	0.0013 (13)	0.0089 (13)	0.0028 (13)
C4	0.0290 (15)	0.0313 (15)	0.0376 (17)	-0.0014 (12)	0.0033 (12)	0.0069 (13)
C5	0.0303 (16)	0.0447 (18)	0.057 (2)	0.0010 (14)	0.0034 (14)	0.0034 (15)
C6	0.0358 (16)	0.0348 (16)	0.0342 (17)	-0.0048 (13)	0.0006 (13)	0.0033 (13)
C7	0.0353 (16)	0.0362 (16)	0.0253 (15)	-0.0022 (13)	0.0062 (12)	0.0002 (12)
C8	0.0272 (14)	0.0312 (15)	0.0283 (16)	-0.0002 (12)	0.0051 (12)	0.0035 (12)
C9	0.0251 (15)	0.0335 (16)	0.0328 (16)	0.0007 (12)	0.0033 (12)	-0.0010 (12)
C10	0.0311 (16)	0.0494 (18)	0.0326 (16)	0.0064 (13)	-0.0002 (12)	0.0049 (14)
C11	0.0290 (16)	0.0383 (16)	0.0281 (15)	0.0026 (12)	0.0037 (12)	0.0004 (12)
C12	0.0237 (14)	0.0313 (15)	0.0317 (16)	0.0040 (12)	0.0039 (11)	0.0046 (12)

C13	0.0265 (15)	0.0297 (15)	0.0399 (17)	0.0041 (12)	0.0038 (12)	0.0038 (12)
C14	0.0350 (17)	0.0422 (18)	0.0478 (19)	-0.0071 (13)	0.0009 (14)	0.0069 (14)
C15	0.0310 (16)	0.0327 (16)	0.0428 (18)	0.0001 (12)	0.0095 (13)	0.0068 (13)
C16	0.0416 (17)	0.0339 (16)	0.0366 (17)	0.0085 (14)	0.0106 (14)	0.0065 (13)
C17	0.060 (2)	0.053 (2)	0.0377 (18)	0.0053 (16)	0.0165 (15)	0.0080 (15)
C18	0.0379 (17)	0.0470 (18)	0.0330 (17)	0.0050 (14)	0.0017 (13)	-0.0020 (13)
C19	0.0265 (15)	0.0461 (18)	0.0374 (17)	0.0001 (13)	0.0061 (13)	0.0016 (14)
C20	0.059 (2)	0.050 (2)	0.0384 (19)	0.0076 (16)	-0.0076 (15)	-0.0096 (15)
C21	0.0400 (17)	0.0324 (15)	0.0250 (15)	-0.0004 (13)	-0.0010 (12)	-0.0006 (12)
C22	0.0463 (18)	0.0322 (16)	0.0318 (16)	0.0014 (13)	0.0066 (14)	-0.0011 (13)
C23	0.0370 (17)	0.0328 (16)	0.0340 (17)	0.0002 (13)	0.0037 (13)	0.0068 (13)
C24	0.0432 (19)	0.0472 (19)	0.049 (2)	0.0020 (15)	0.0092 (15)	0.0045 (15)
C25	0.0348 (17)	0.0534 (19)	0.0304 (17)	-0.0037 (14)	-0.0001 (13)	0.0006 (14)
C26	0.0363 (17)	0.0537 (19)	0.0259 (16)	-0.0008 (14)	0.0055 (13)	-0.0052 (13)
C27	0.0321 (16)	0.0328 (15)	0.0319 (16)	0.0014 (12)	0.0022 (12)	0.0045 (12)
C28	0.0293 (16)	0.0348 (16)	0.0385 (17)	0.0005 (12)	-0.0029 (13)	0.0028 (13)
C29	0.0317 (16)	0.0450 (18)	0.0426 (18)	0.0012 (13)	-0.0034 (13)	0.0101 (14)
C30	0.0323 (16)	0.0323 (16)	0.0322 (16)	-0.0047 (12)	-0.0019 (12)	0.0022 (12)
C31	0.0273 (15)	0.0294 (15)	0.0364 (17)	0.0040 (12)	-0.0009 (12)	0.0000 (12)
C32	0.0313 (16)	0.0284 (15)	0.0385 (17)	0.0042 (12)	0.0012 (13)	-0.0026 (12)
C33	0.0327 (17)	0.0488 (19)	0.049 (2)	-0.0061 (14)	-0.0013 (14)	-0.0027 (15)
C34	0.0400 (18)	0.0331 (16)	0.0470 (19)	0.0034 (13)	0.0109 (15)	0.0037 (14)
C35	0.054 (2)	0.0390 (17)	0.0339 (18)	0.0128 (15)	0.0068 (15)	-0.0006 (14)
C36	0.090 (3)	0.066 (2)	0.038 (2)	0.018 (2)	0.0131 (18)	0.0051 (17)
C37	0.0427 (18)	0.0452 (18)	0.0375 (18)	0.0141 (15)	-0.0092 (14)	-0.0054 (14)
C38	0.0281 (16)	0.0424 (17)	0.0446 (19)	0.0030 (13)	-0.0011 (13)	-0.0046 (14)

Geometric parameters (\AA , $^{\circ}$)

N1—C9	1.272 (3)	C16—C17	1.521 (3)
N1—C8	1.421 (3)	C17—H17A	0.9800
N2—C11	1.363 (3)	C17—H17B	0.9800
N2—C9	1.372 (3)	C17—H17C	0.9800
N2—C10	1.463 (3)	C18—C19	1.391 (3)
N3—C11	1.275 (3)	C18—H18	0.9500
N3—C12	1.417 (3)	C19—H19	0.9500
N4—C28	1.265 (3)	C20—C21	1.506 (4)
N4—C27	1.422 (3)	C20—H20A	0.9800
N5—C30	1.370 (3)	C20—H20B	0.9800
N5—C28	1.379 (3)	C20—H20C	0.9800
N5—C29	1.465 (3)	C21—C22	1.388 (4)
N6—C30	1.269 (3)	C21—C27	1.393 (3)
N6—C31	1.414 (3)	C22—C23	1.385 (4)
C1—C2	1.504 (3)	C22—H22	0.9500
C1—H1A	0.9800	C23—C25	1.378 (4)
C1—H1B	0.9800	C23—C24	1.508 (4)
C1—H1C	0.9800	C24—H24A	0.9800
C2—C8	1.393 (3)	C24—H24B	0.9800

C2—C3	1.396 (3)	C24—H24C	0.9800
C3—C4	1.385 (4)	C25—C26	1.378 (4)
C3—H3	0.9500	C25—H25	0.9500
C4—C6	1.388 (4)	C26—C27	1.389 (3)
C4—C5	1.512 (3)	C26—H26	0.9500
C5—H5A	0.9800	C28—H28	0.9500
C5—H5B	0.9800	C29—H29A	0.9800
C5—H5C	0.9800	C29—H29B	0.9800
C6—C7	1.384 (4)	C29—H29C	0.9800
C6—H6	0.9500	C30—H30	0.9500
C7—C8	1.390 (3)	C31—C38	1.399 (4)
C7—H7	0.9500	C31—C32	1.406 (4)
C9—H9	0.9500	C32—C34	1.385 (4)
C10—H10A	0.9800	C32—C33	1.503 (4)
C10—H10B	0.9800	C33—H33A	0.9800
C10—H10C	0.9800	C33—H33B	0.9800
C11—H11	0.9500	C33—H33C	0.9800
C12—C19	1.395 (4)	C34—C35	1.386 (4)
C12—C13	1.404 (3)	C34—H34	0.9500
C13—C15	1.393 (3)	C35—C37	1.377 (4)
C13—C14	1.505 (4)	C35—C36	1.516 (4)
C14—H14A	0.9800	C36—H36A	0.9800
C14—H14B	0.9800	C36—H36B	0.9800
C14—H14C	0.9800	C36—H36C	0.9800
C15—C16	1.384 (4)	C37—C38	1.388 (4)
C15—H15	0.9500	C37—H37	0.9500
C16—C18	1.388 (4)	C38—H38	0.9500
C9—N1—C8	115.5 (2)	C16—C18—H18	119.6
C11—N2—C9	118.9 (2)	C19—C18—H18	119.6
C11—N2—C10	120.9 (2)	C18—C19—C12	121.0 (3)
C9—N2—C10	120.2 (2)	C18—C19—H19	119.5
C11—N3—C12	118.1 (2)	C12—C19—H19	119.5
C28—N4—C27	117.3 (2)	C21—C20—H20A	109.5
C30—N5—C28	119.9 (2)	C21—C20—H20B	109.5
C30—N5—C29	120.2 (2)	H20A—C20—H20B	109.5
C28—N5—C29	119.9 (2)	C21—C20—H20C	109.5
C30—N6—C31	117.1 (2)	H20A—C20—H20C	109.5
C2—C1—H1A	109.5	H20B—C20—H20C	109.5
C2—C1—H1B	109.5	C22—C21—C27	118.2 (3)
H1A—C1—H1B	109.5	C22—C21—C20	120.6 (2)
C2—C1—H1C	109.5	C27—C21—C20	121.1 (3)
H1A—C1—H1C	109.5	C23—C22—C21	123.3 (3)
H1B—C1—H1C	109.5	C23—C22—H22	118.3
C8—C2—C3	117.9 (3)	C21—C22—H22	118.3
C8—C2—C1	121.1 (2)	C25—C23—C22	117.4 (3)
C3—C2—C1	121.0 (2)	C25—C23—C24	121.1 (3)
C4—C3—C2	123.1 (2)	C22—C23—C24	121.5 (3)

C4—C3—H3	118.5	C23—C24—H24A	109.5
C2—C3—H3	118.5	C23—C24—H24B	109.5
C3—C4—C6	117.5 (2)	H24A—C24—H24B	109.5
C3—C4—C5	120.9 (2)	C23—C24—H24C	109.5
C6—C4—C5	121.5 (3)	H24A—C24—H24C	109.5
C4—C5—H5A	109.5	H24B—C24—H24C	109.5
C4—C5—H5B	109.5	C26—C25—C23	120.6 (3)
H5A—C5—H5B	109.5	C26—C25—H25	119.7
C4—C5—H5C	109.5	C23—C25—H25	119.7
H5A—C5—H5C	109.5	C25—C26—C27	121.6 (3)
H5B—C5—H5C	109.5	C25—C26—H26	119.2
C7—C6—C4	121.0 (3)	C27—C26—H26	119.2
C7—C6—H6	119.5	C26—C27—C21	118.8 (2)
C4—C6—H6	119.5	C26—C27—N4	122.2 (2)
C6—C7—C8	120.5 (2)	C21—C27—N4	118.9 (2)
C6—C7—H7	119.7	N4—C28—N5	123.4 (3)
C8—C7—H7	119.7	N4—C28—H28	118.3
C7—C8—C2	120.0 (2)	N5—C28—H28	118.3
C7—C8—N1	121.4 (2)	N5—C29—H29A	109.5
C2—C8—N1	118.6 (2)	N5—C29—H29B	109.5
N1—C9—N2	123.4 (3)	H29A—C29—H29B	109.5
N1—C9—H9	118.3	N5—C29—H29C	109.5
N2—C9—H9	118.3	H29A—C29—H29C	109.5
N2—C10—H10A	109.5	H29B—C29—H29C	109.5
N2—C10—H10B	109.5	N6—C30—N5	123.6 (3)
H10A—C10—H10B	109.5	N6—C30—H30	118.2
N2—C10—H10C	109.5	N5—C30—H30	118.2
H10A—C10—H10C	109.5	C38—C31—C32	118.8 (3)
H10B—C10—H10C	109.5	C38—C31—N6	123.3 (2)
N3—C11—N2	122.6 (2)	C32—C31—N6	117.9 (2)
N3—C11—H11	118.7	C34—C32—C31	118.7 (3)
N2—C11—H11	118.7	C34—C32—C33	121.9 (3)
C19—C12—C13	118.7 (2)	C31—C32—C33	119.4 (2)
C19—C12—N3	122.5 (2)	C32—C33—H33A	109.5
C13—C12—N3	118.6 (2)	C32—C33—H33B	109.5
C15—C13—C12	119.0 (3)	H33A—C33—H33B	109.5
C15—C13—C14	120.6 (2)	C32—C33—H33C	109.5
C12—C13—C14	120.4 (2)	H33A—C33—H33C	109.5
C13—C14—H14A	109.5	H33B—C33—H33C	109.5
C13—C14—H14B	109.5	C32—C34—C35	122.5 (3)
H14A—C14—H14B	109.5	C32—C34—H34	118.7
C13—C14—H14C	109.5	C35—C34—H34	118.7
H14A—C14—H14C	109.5	C37—C35—C34	118.4 (3)
H14B—C14—H14C	109.5	C37—C35—C36	120.9 (3)
C16—C15—C13	122.6 (3)	C34—C35—C36	120.7 (3)
C16—C15—H15	118.7	C35—C36—H36A	109.5
C13—C15—H15	118.7	C35—C36—H36B	109.5
C15—C16—C18	118.0 (3)	H36A—C36—H36B	109.5

C15—C16—C17	121.4 (3)	C35—C36—H36C	109.5
C18—C16—C17	120.6 (3)	H36A—C36—H36C	109.5
C16—C17—H17A	109.5	H36B—C36—H36C	109.5
C16—C17—H17B	109.5	C35—C37—C38	120.8 (3)
H17A—C17—H17B	109.5	C35—C37—H37	119.6
C16—C17—H17C	109.5	C38—C37—H37	119.6
H17A—C17—H17C	109.5	C37—C38—C31	120.7 (3)
H17B—C17—H17C	109.5	C37—C38—H38	119.7
C16—C18—C19	120.7 (3)	C31—C38—H38	119.7
C8—C2—C3—C4	-1.4 (4)	C27—C21—C22—C23	0.2 (4)
C1—C2—C3—C4	178.2 (2)	C20—C21—C22—C23	178.3 (3)
C2—C3—C4—C6	-0.1 (4)	C21—C22—C23—C25	1.2 (4)
C2—C3—C4—C5	-178.6 (2)	C21—C22—C23—C24	179.6 (3)
C3—C4—C6—C7	0.2 (4)	C22—C23—C25—C26	-0.3 (4)
C5—C4—C6—C7	178.7 (2)	C24—C23—C25—C26	-178.7 (3)
C4—C6—C7—C8	1.1 (4)	C23—C25—C26—C27	-2.0 (4)
C6—C7—C8—C2	-2.7 (4)	C25—C26—C27—C21	3.4 (4)
C6—C7—C8—N1	179.4 (2)	C25—C26—C27—N4	179.4 (3)
C3—C2—C8—C7	2.7 (4)	C22—C21—C27—C26	-2.5 (4)
C1—C2—C8—C7	-176.8 (2)	C20—C21—C27—C26	179.5 (3)
C3—C2—C8—N1	-179.3 (2)	C22—C21—C27—N4	-178.6 (2)
C1—C2—C8—N1	1.1 (4)	C20—C21—C27—N4	3.3 (4)
C9—N1—C8—C7	-46.5 (3)	C28—N4—C27—C26	37.4 (4)
C9—N1—C8—C2	135.5 (3)	C28—N4—C27—C21	-146.6 (3)
C8—N1—C9—N2	177.4 (2)	C27—N4—C28—N5	-174.0 (2)
C11—N2—C9—N1	176.4 (2)	C30—N5—C28—N4	179.1 (2)
C10—N2—C9—N1	-4.4 (4)	C29—N5—C28—N4	-1.8 (4)
C12—N3—C11—N2	-173.2 (2)	C31—N6—C30—N5	-179.8 (2)
C9—N2—C11—N3	-178.5 (2)	C28—N5—C30—N6	178.2 (2)
C10—N2—C11—N3	2.3 (4)	C29—N5—C30—N6	-0.9 (4)
C11—N3—C12—C19	35.7 (4)	C30—N6—C31—C38	-44.3 (4)
C11—N3—C12—C13	-149.0 (2)	C30—N6—C31—C32	137.1 (3)
C19—C12—C13—C15	-1.5 (4)	C38—C31—C32—C34	2.6 (4)
N3—C12—C13—C15	-177.0 (2)	N6—C31—C32—C34	-178.8 (2)
C19—C12—C13—C14	179.0 (2)	C38—C31—C32—C33	-177.2 (2)
N3—C12—C13—C14	3.6 (4)	N6—C31—C32—C33	1.5 (4)
C12—C13—C15—C16	-0.1 (4)	C31—C32—C34—C35	-1.6 (4)
C14—C13—C15—C16	179.4 (2)	C33—C32—C34—C35	178.2 (3)
C13—C15—C16—C18	1.2 (4)	C32—C34—C35—C37	-0.4 (4)
C13—C15—C16—C17	-179.0 (2)	C32—C34—C35—C36	-179.6 (3)
C15—C16—C18—C19	-0.6 (4)	C34—C35—C37—C38	1.3 (4)
C17—C16—C18—C19	179.6 (3)	C36—C35—C37—C38	-179.4 (3)
C16—C18—C19—C12	-1.0 (4)	C35—C37—C38—C31	-0.3 (4)
C13—C12—C19—C18	2.1 (4)	C32—C31—C38—C37	-1.6 (4)
N3—C12—C19—C18	177.3 (2)	N6—C31—C38—C37	179.8 (2)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg4 are the centroids of which rings?

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
C29—H29C···N3 ⁱ	0.98	2.49	3.334 (4)	144
C17—H17A···Cg4 ⁱ	0.98	2.92	3.806 (3)	151
C24—H24A···Cg1 ⁱⁱ	0.98	2.85	3.828 (3)	175
C33—H33B···Cg4 ⁱⁱⁱ	0.98	2.88	3.625 (3)	134

Symmetry codes: (i) $-x, y-1/2, -z+3/2$; (ii) $-x, y+1/2, -z+3/2$; (iii) $-x+1, y+1/2, -z+3/2$.