

4'-(1*H*-Imidazol-2-yl)-3'-[*(1H*-indol-3-yl)-carbonyl]-1'-methyl-2-oxospiro[indoline-3,2'-pyrrolidine]-3'-carbonitrile 0.15-hydrate

S. Antony Inglebert,^a Yuvaraj Arun,^b K. Sethusankar^{c*} and Paramasivam T. Perumal^b

^aSri Ram Engineering College, Chennai 602 024, India, ^bOrganic Chemistry Division, Central Leather Research Institute, Adyar, Chennai 600 020, India, and ^cDepartment of Physics, RKM Vivekananda College (Autonomous), Chennai 600 004, India
Correspondence e-mail: ksethusankar@yahoo.co.in


Received 1 August 2013; accepted 19 August 2013

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; H-atom completeness 99%; disorder in solvent or counterion; R factor = 0.040; wR factor = 0.110; data-to-parameter ratio = 15.1.

In the title compound, $\text{C}_{25}\text{H}_{20}\text{N}_6\text{O}_2 \cdot 0.15\text{H}_2\text{O}$, the dihedral angles between the least-squares planes of the indole and pyrrolidine rings and between the oxindole and imidazole rings are $77.66(7)$ and $45.31(7)^\circ$, respectively. The pyrrolidine ring and the fused five-membered pyrrolidine ring of the oxindole moiety exhibit twisted conformations. The amide N atom is involved in both intra- and intermolecular hydrogen bonding, having a bifurcated character. The molecular structure is characterized by an intramolecular N—H···O hydrogen bond, which generates an *S*(7) ring motif while an intermolecular N—H···O hydrogen bond links the organic and solvent water molecules. In the crystal, N—H···N hydrogen bonds generate a zigzag chain running parallel to *c*-axis direction. The H atoms of the solvent water molecule were not located.

Related literature

For background to indole derivatives and their biological activity, see: Rudrangi *et al.* (2011). For puckering parameters, see: Cremer & Pople (1975). For bond-length data, see: Allen *et al.* (1987). For graph-set notation, see: Bernstein *et al.* (1995). For a related structure, see: Inglebert *et al.* (2013).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{20}\text{N}_6\text{O}_2 \cdot 0.15\text{H}_2\text{O}$	$V = 2100.6(15)\text{ \AA}^3$
$M_r = 439.17$	$Z = 4$
Monoclinic, $P2_1/n$	$\text{Mo } K\alpha$ radiation
$a = 8.650(5)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 16.952(5)\text{ \AA}$	$T = 295\text{ K}$
$c = 14.438(5)\text{ \AA}$	$0.35 \times 0.30 \times 0.25\text{ mm}$
$\beta = 97.161(5)^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer	22198 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2008)	4819 independent reflections
$T_{\min} = 0.968$, $T_{\max} = 0.977$	3757 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.110$	$\Delta\rho_{\text{max}} = 0.21\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.18\text{ e \AA}^{-3}$
4819 reflections	
320 parameters	
3 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N1—H1A···N5 ⁱ	0.89 (1)	2.13 (1)	2.9889 (19)	164 (2)
N6—H6A···O1W	0.90 (1)	1.98 (2)	2.714 (8)	138 (2)
N6—H6A···O1	0.90 (1)	2.57 (2)	3.064 (2)	116 (2)

Symmetry code: (i) $x + \frac{1}{2}$, $-y + \frac{1}{2}$, $z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

The authors gratefully acknowledge Dr Babu Varghese, SAIF, IIT, Chennai, India, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2411).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2008). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Inglebert, S. A., Arun, Y., Sethusankar, K. & Perumal, P. T. (2013). *Acta Cryst. E69*, o1328–o1329.
- Rudrangi, S. R. S., Bontha, V. K., Manda, V. R. & Bethi, S. (2011). *Asian J. Res. Chem.* **4**, 335–338.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supporting information

Acta Cryst. (2013). E69, o1481 [doi:10.1107/S1600536813023246]

4'-(1*H*-imidazol-2-yl)-3'-(1*H*-indol-3-yl)carbonyl]-1'-methyl-2-oxospiro-[indoline-3,2'-pyrrolidine]-3'-carbonitrile 0.15-hydrate

S. Antony Inglebert, Yuvaraj Arun, K. Sethusankar and Paramasivam T. Perumal

S1. Comment

Oxindoles are endogenous aromatic organic compounds that are found in the tissues and body fluids of mammals and in the natural products of some plants. Oxindoles exhibit an extensive range of biological effects, which include anticancer, anti-inflammatory, antiviral, antibacterial, antihypertensive and anticonvulsant activities. Oxindoles are also used to inhibit the replication of HIV and combat the infections that are caused by drug-resistant, drug-sensitive and mutant strains of HIV (Rudrangi *et al.*, 2011).

The title compound consists of a pyrrolidine ring connected to an imidazole at C11, an oxindole ring system at C15, a methyl group at N3 and an indole unit at C10 *via* carbonyl group. In addition, the asymmetric unit contains a 0.15 occupancy water molecule. The H atoms of the partial occupancy water molecules are neither located nor included in the refinement. The title structure exhibits structural similarities with the previously reported structure (Inglebert *et al.*, 2013).

The pyrrolidine ring adopts a twisted conformation with puckering parameters $q(2) = 0.4351 (14)\text{\AA}$ and $\phi(2) = 309.66 (19)^\circ$. Pyrrole ring in the oxindole unit also having twisted conformation with puckering parameters $q(2) = 0.0931 (15)\text{\AA}$ and $\phi(2) = 304.2 (9)^\circ$. The least square plane of oxindole unit makes dihedral angles of $78.50 (4)^\circ$ and $44.28 (5)^\circ$ with the pyrrolidine and imidazole, respectively. The indole unit is essentially planar - maximum deviation = $0.0138 (17)\text{\AA}$ for the C2 atom] and is oriented at a least square plane that makes dihedral angles of $51.42 (4)^\circ$, $39.92 (4)^\circ$ and $70.15 (4)^\circ$, with the oxindole unit, pyrrolidine and imidazole rings, respectively.

The cyano bond distance C13≡N2 agrees well with the reported value of $1.138 (7)\text{\AA}$ (Allen *et al.*, 1987). The sum of the angle around atom N3 ($340.14 (35)^\circ$) is in accordance with sp^3 hybridization. The amide N atom shows bifurcated intramolecular hydrogen bond (N—H···O) with an O atom of the carbonyl group and an intermolecular hydrogen (N—H···O) bond with the 0.15 occupancy solvent water molecule. In addition, the classical intermolecular N—H···N hydrogen bonds generate a zigzag chain running parallel to *c* axis.

S2. Experimental

A mixture of isatin (1 mmol), sarcosine (1.2 mmol) and 3-(1*H*-imidazol-2-yl)-2-(1*H*-indole-3-carbonyl)acrylonitrile (1 mmol) were refluxed in ethanol (30 ml). After completion of the reaction as evidenced by *TLC* analysis, the reaction mixture was poured into ice–water, the resulting solid was filtered off and purified by column chromatography using ethyl acetate : petroleum ether (6 : 4) as an eluent to afford pure product.

S3. Refinement

Positions of hydrogen atoms were localized from the difference electron density maps and their distances were geometrically constrained. The H atoms bound to the C atoms were treated as riding atoms with $d(\text{C}—\text{H}) = 0.93\text{\AA}$ for

aromatic H, $d(C—H) = 0.97\text{\AA}$ for methylene H with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $d(C—H) = 0.96\text{\AA}$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms. The rotation angles for methyl groups were optimized by least squares. The N bonded H atoms were refined freely.





Figure 1


The asymmetric unit of the title compound with the atom numbering scheme. Displacement ellipsoids drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius. H atoms of water molecule are not found.

**Figure 2**

The packing diagram of the title compound viewed along the *a* axis. H atoms have omitted for clarity.

4'-(1*H*-Imidazol-2-yl)-3'-[*(1H*-indol-3-yl)carbonyl]-1'-methyl-2-oxospiro[indoline-3,2'-pyrrolidine]-3'-carbonitrile 0.15-hydrate

Crystal data



M_r = 439.17

Monoclinic, P2₁/n

Hall symbol: -P 2yn

a = 8.650 (5) Å

b = 16.952 (5) Å

c = 14.438 (5) Å

β = 97.161 (5)°

V = 2100.6 (15) Å³

Z = 4

F(000) = 916.8

D_x = 1.389 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 4819 reflections

θ = 2.4–27.5°

μ = 0.09 mm⁻¹

T = 295 K

Block, colourless

0.35 × 0.30 × 0.25 mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and φ scans

Absorption correction: multi-scan

(SADABS; Bruker, 2008)

*T*_{min} = 0.968, *T*_{max} = 0.977

22198 measured reflections

4819 independent reflections

3757 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.024

θ_{max} = 27.5°, θ_{min} = 2.4°

h = -11→10

k = -21→22

l = -18→18

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.040$$

$$wR(F^2) = 0.110$$

$$S = 1.03$$

4819 reflections

320 parameters

3 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0524P)^2 + 0.5356P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$	Occ. (<1)
O1	0.37937 (14)	0.07447 (6)	0.49050 (7)	0.0487 (3)	
O2	0.60471 (12)	0.18182 (6)	0.64371 (8)	0.0489 (3)	
N3	0.33938 (14)	0.30031 (7)	0.63558 (8)	0.0366 (3)	
N2	0.66582 (14)	0.25656 (8)	0.52028 (9)	0.0406 (3)	
N1	0.38442 (14)	0.21594 (7)	0.22422 (8)	0.0380 (3)	
C15	0.40978 (15)	0.26985 (7)	0.55702 (9)	0.0315 (3)	
C10	0.28828 (14)	0.20355 (7)	0.52052 (9)	0.0292 (3)	
N6	0.12101 (15)	0.03867 (7)	0.61150 (9)	0.0432 (3)	
N5	-0.02773 (15)	0.14222 (8)	0.61946 (11)	0.0508 (3)	
C1	0.42119 (15)	0.09829 (8)	0.29638 (9)	0.0318 (3)	
C16	0.44630 (15)	0.33152 (8)	0.48826 (9)	0.0340 (3)	
C11	0.25875 (15)	0.16641 (8)	0.61571 (9)	0.0326 (3)	
H11	0.3466	0.1309	0.6343	0.039*	
C6	0.43006 (16)	0.13884 (8)	0.21219 (9)	0.0352 (3)	
C9	0.34886 (14)	0.13897 (8)	0.45599 (9)	0.0313 (3)	
C8	0.36716 (15)	0.15545 (8)	0.36003 (9)	0.0316 (3)	
C12	0.27540 (17)	0.23523 (8)	0.68508 (10)	0.0399 (3)	
H12A	0.1749	0.2494	0.7034	0.048*	
H12B	0.3452	0.2213	0.7406	0.048*	
C7	0.34692 (16)	0.22545 (8)	0.31089 (9)	0.0359 (3)	
H7	0.3123	0.2724	0.3345	0.043*	
C21	0.59667 (16)	0.31921 (8)	0.46655 (10)	0.0368 (3)	
C2	0.46065 (17)	0.01848 (8)	0.30176 (10)	0.0386 (3)	
H2	0.4569	-0.0096	0.3568	0.046*	

C22	0.57128 (15)	0.22878 (8)	0.58113 (10)	0.0359 (3)	
C13	0.14541 (16)	0.24026 (8)	0.47441 (9)	0.0361 (3)	
C3	0.50525 (19)	-0.01783 (9)	0.22397 (11)	0.0478 (4)	
H3	0.5299	-0.0712	0.2266	0.057*	
C23	0.11522 (16)	0.11799 (8)	0.61483 (9)	0.0351 (3)	
C19	0.5689 (2)	0.42550 (10)	0.36056 (12)	0.0552 (4)	
H19	0.6079	0.4569	0.3160	0.066*	
N4	0.03167 (16)	0.26634 (9)	0.43965 (10)	0.0581 (4)	
C5	0.47677 (19)	0.10239 (10)	0.13400 (10)	0.0467 (4)	
H5	0.4824	0.1302	0.0790	0.056*	
C20	0.66026 (19)	0.36546 (10)	0.40259 (11)	0.0476 (4)	
H20	0.7607	0.3566	0.3883	0.057*	
C4	0.5143 (2)	0.02372 (10)	0.14131 (11)	0.0512 (4)	
H4	0.5462	-0.0023	0.0903	0.061*	
C17	0.35815 (18)	0.39336 (8)	0.44843 (11)	0.0424 (3)	
H17	0.2594	0.4036	0.4648	0.051*	
C18	0.4207 (2)	0.44015 (9)	0.38310 (12)	0.0524 (4)	
H18	0.3625	0.4815	0.3544	0.063*	
C25	-0.1158 (2)	0.07419 (10)	0.61833 (14)	0.0584 (5)	
H25	-0.2227	0.0728	0.6207	0.070*	
C14	0.4324 (2)	0.35557 (10)	0.69609 (12)	0.0555 (4)	
H14A	0.5180	0.3281	0.7306	0.083*	
H14B	0.3690	0.3790	0.7387	0.083*	
H14C	0.4718	0.3961	0.6589	0.083*	
C24	-0.0261 (2)	0.01074 (10)	0.61340 (12)	0.0513 (4)	
H24	-0.0576	-0.0418	0.6116	0.062*	
O1W	0.2610 (10)	-0.0987 (6)	0.5729 (7)	0.082 (3)	0.15
H2A	0.7632 (12)	0.2407 (9)	0.5205 (12)	0.052 (5)*	
H1A	0.393 (2)	0.2564 (8)	0.1861 (11)	0.061 (5)*	
H6A	0.2060 (17)	0.0112 (11)	0.6021 (15)	0.075 (6)*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0721 (8)	0.0369 (6)	0.0403 (5)	0.0139 (5)	0.0190 (5)	0.0060 (4)
O2	0.0363 (6)	0.0479 (6)	0.0605 (7)	0.0013 (5)	-0.0018 (5)	0.0177 (5)
N3	0.0386 (6)	0.0330 (6)	0.0384 (6)	-0.0033 (5)	0.0063 (5)	-0.0054 (5)
N2	0.0267 (6)	0.0464 (7)	0.0486 (7)	0.0010 (5)	0.0047 (5)	0.0038 (5)
N1	0.0448 (7)	0.0351 (6)	0.0344 (6)	0.0017 (5)	0.0053 (5)	0.0064 (5)
C15	0.0275 (6)	0.0294 (6)	0.0374 (6)	-0.0001 (5)	0.0035 (5)	0.0001 (5)
C10	0.0257 (6)	0.0300 (6)	0.0324 (6)	0.0002 (5)	0.0053 (5)	0.0000 (5)
N6	0.0462 (8)	0.0346 (7)	0.0494 (7)	-0.0044 (6)	0.0088 (6)	-0.0035 (5)
N5	0.0386 (7)	0.0406 (7)	0.0776 (9)	-0.0088 (6)	0.0247 (6)	-0.0085 (6)
C1	0.0292 (6)	0.0354 (7)	0.0315 (6)	-0.0020 (5)	0.0064 (5)	0.0007 (5)
C16	0.0333 (7)	0.0297 (7)	0.0389 (7)	-0.0044 (5)	0.0041 (5)	-0.0004 (5)
C11	0.0324 (7)	0.0348 (7)	0.0316 (6)	-0.0025 (5)	0.0077 (5)	-0.0014 (5)
C6	0.0350 (7)	0.0360 (7)	0.0350 (7)	-0.0007 (6)	0.0059 (5)	0.0028 (5)
C9	0.0294 (6)	0.0312 (7)	0.0339 (6)	0.0005 (5)	0.0060 (5)	0.0006 (5)

C8	0.0295 (6)	0.0323 (7)	0.0334 (6)	-0.0004 (5)	0.0059 (5)	0.0003 (5)
C12	0.0425 (8)	0.0419 (8)	0.0364 (7)	-0.0086 (6)	0.0098 (6)	-0.0073 (6)
C7	0.0355 (7)	0.0358 (7)	0.0361 (7)	0.0022 (6)	0.0027 (5)	-0.0003 (5)
C21	0.0346 (7)	0.0364 (7)	0.0393 (7)	-0.0054 (6)	0.0043 (5)	-0.0027 (6)
C2	0.0435 (8)	0.0353 (7)	0.0384 (7)	0.0014 (6)	0.0103 (6)	0.0040 (6)
C22	0.0288 (7)	0.0349 (7)	0.0428 (7)	-0.0029 (5)	-0.0006 (5)	0.0002 (6)
C13	0.0313 (7)	0.0401 (8)	0.0371 (7)	0.0006 (6)	0.0054 (5)	-0.0046 (6)
C3	0.0597 (10)	0.0378 (8)	0.0474 (8)	0.0095 (7)	0.0127 (7)	-0.0012 (6)
C23	0.0393 (8)	0.0335 (7)	0.0343 (6)	-0.0058 (6)	0.0117 (5)	-0.0038 (5)
C19	0.0660 (11)	0.0504 (10)	0.0516 (9)	-0.0112 (8)	0.0166 (8)	0.0112 (7)
N4	0.0386 (7)	0.0737 (10)	0.0595 (8)	0.0149 (7)	-0.0037 (6)	-0.0056 (7)
C5	0.0568 (9)	0.0520 (9)	0.0331 (7)	0.0037 (7)	0.0129 (6)	0.0051 (6)
C20	0.0454 (9)	0.0515 (9)	0.0480 (8)	-0.0071 (7)	0.0141 (7)	0.0021 (7)
C4	0.0645 (11)	0.0526 (10)	0.0391 (8)	0.0101 (8)	0.0165 (7)	-0.0054 (7)
C17	0.0436 (8)	0.0324 (7)	0.0513 (8)	0.0022 (6)	0.0066 (6)	0.0023 (6)
C18	0.0642 (11)	0.0359 (8)	0.0564 (9)	0.0003 (7)	0.0048 (8)	0.0115 (7)
C25	0.0441 (9)	0.0502 (10)	0.0849 (13)	-0.0177 (8)	0.0239 (8)	-0.0101 (9)
C14	0.0714 (11)	0.0435 (9)	0.0515 (9)	-0.0174 (8)	0.0078 (8)	-0.0134 (7)
C24	0.0586 (10)	0.0394 (8)	0.0569 (9)	-0.0187 (8)	0.0116 (7)	-0.0048 (7)
O1W	0.063 (5)	0.070 (6)	0.102 (7)	0.036 (5)	-0.028 (5)	-0.039 (5)

Geometric parameters (\AA , $^{\circ}$)

O1—C9	1.2168 (16)	C11—H11	0.9800
O2—C22	1.2119 (17)	C6—C5	1.390 (2)
N3—C15	1.4478 (17)	C9—C8	1.4413 (18)
N3—C14	1.4538 (19)	C8—C7	1.3826 (19)
N3—C12	1.4604 (18)	C12—H12A	0.9700
N2—C22	1.3570 (19)	C12—H12B	0.9700
N2—C21	1.4043 (19)	C7—H7	0.9300
N2—H2A	0.884 (9)	C21—C20	1.377 (2)
N1—C7	1.3410 (18)	C2—C3	1.377 (2)
N1—C6	1.3825 (19)	C2—H2	0.9300
N1—H1A	0.888 (9)	C13—N4	1.1367 (19)
C15—C16	1.5022 (18)	C3—C4	1.396 (2)
C15—C22	1.561 (2)	C3—H3	0.9300
C15—C10	1.5835 (18)	C19—C20	1.382 (2)
C10—C13	1.4671 (19)	C19—C18	1.383 (3)
C10—C11	1.5615 (18)	C19—H19	0.9300
C10—C9	1.5693 (18)	C5—C4	1.373 (2)
N6—C23	1.3467 (19)	C5—H5	0.9300
N6—C24	1.361 (2)	C20—H20	0.9300
N6—H6A	0.895 (9)	C4—H4	0.9300
N5—C23	1.313 (2)	C17—C18	1.393 (2)
N5—C25	1.381 (2)	C17—H17	0.9300
C1—C2	1.395 (2)	C18—H18	0.9300
C1—C6	1.4068 (18)	C25—C24	1.333 (2)
C1—C8	1.4522 (18)	C25—H25	0.9300

C16—C17	1.3790 (19)	C14—H14A	0.9600
C16—C21	1.391 (2)	C14—H14B	0.9600
C11—C23	1.4871 (19)	C14—H14C	0.9600
C11—C12	1.5325 (19)	C24—H24	0.9300
C15—N3—C14	116.43 (12)	H12A—C12—H12B	108.8
C15—N3—C12	109.63 (11)	N1—C7—C8	110.23 (12)
C14—N3—C12	114.08 (12)	N1—C7—H7	124.9
C22—N2—C21	111.75 (12)	C8—C7—H7	124.9
C22—N2—H2A	122.9 (11)	C20—C21—C16	121.90 (14)
C21—N2—H2A	124.9 (11)	C20—C21—N2	128.52 (14)
C7—N1—C6	109.69 (11)	C16—C21—N2	109.56 (12)
C7—N1—H1A	122.3 (12)	C3—C2—C1	118.65 (13)
C6—N1—H1A	127.3 (12)	C3—C2—H2	120.7
N3—C15—C16	114.40 (11)	C1—C2—H2	120.7
N3—C15—C22	115.88 (11)	O2—C22—N2	127.13 (13)
C16—C15—C22	101.64 (10)	O2—C22—C15	125.67 (13)
N3—C15—C10	100.80 (10)	N2—C22—C15	107.20 (12)
C16—C15—C10	117.35 (11)	N4—C13—C10	177.44 (16)
C22—C15—C10	107.23 (10)	C2—C3—C4	121.51 (14)
C13—C10—C11	110.45 (11)	C2—C3—H3	119.2
C13—C10—C9	110.56 (10)	C4—C3—H3	119.2
C11—C10—C9	110.16 (10)	N5—C23—N6	110.71 (12)
C13—C10—C15	109.67 (11)	N5—C23—C11	128.15 (13)
C11—C10—C15	99.66 (10)	N6—C23—C11	121.11 (13)
C9—C10—C15	115.84 (10)	C20—C19—C18	121.66 (15)
C23—N6—C24	107.91 (13)	C20—C19—H19	119.2
C23—N6—H6A	124.1 (14)	C18—C19—H19	119.2
C24—N6—H6A	127.6 (14)	C4—C5—C6	117.31 (13)
C23—N5—C25	105.06 (14)	C4—C5—H5	121.3
C2—C1—C6	118.86 (12)	C6—C5—H5	121.3
C2—C1—C8	135.04 (12)	C21—C20—C19	117.34 (15)
C6—C1—C8	106.10 (12)	C21—C20—H20	121.3
C17—C16—C21	120.26 (13)	C19—C20—H20	121.3
C17—C16—C15	130.83 (13)	C5—C4—C3	121.18 (14)
C21—C16—C15	108.91 (12)	C5—C4—H4	119.4
C23—C11—C12	115.86 (11)	C3—C4—H4	119.4
C23—C11—C10	116.30 (11)	C16—C17—C18	118.30 (15)
C12—C11—C10	104.83 (11)	C16—C17—H17	120.9
C23—C11—H11	106.4	C18—C17—H17	120.9
C12—C11—H11	106.4	C19—C18—C17	120.45 (15)
C10—C11—H11	106.4	C19—C18—H18	119.8
N1—C6—C5	129.68 (13)	C17—C18—H18	119.8
N1—C6—C1	107.83 (12)	C24—C25—N5	110.53 (15)
C5—C6—C1	122.49 (13)	C24—C25—H25	124.7
O1—C9—C8	121.56 (12)	N5—C25—H25	124.7
O1—C9—C10	117.08 (11)	N3—C14—H14A	109.5
C8—C9—C10	121.35 (11)	N3—C14—H14B	109.5

C7—C8—C9	129.61 (12)	H14A—C14—H14B	109.5
C7—C8—C1	106.14 (11)	N3—C14—H14C	109.5
C9—C8—C1	124.17 (12)	H14A—C14—H14C	109.5
N3—C12—C11	105.41 (11)	H14B—C14—H14C	109.5
N3—C12—H12A	110.7	C25—C24—N6	105.79 (14)
C11—C12—H12A	110.7	C25—C24—H24	127.1
N3—C12—H12B	110.7	N6—C24—H24	127.1
C11—C12—H12B	110.7		
C14—N3—C15—C16	−62.11 (16)	C15—N3—C12—C11	−19.05 (14)
C12—N3—C15—C16	166.52 (11)	C14—N3—C12—C11	−151.66 (13)
C14—N3—C15—C22	55.69 (16)	C23—C11—C12—N3	−139.92 (12)
C12—N3—C15—C22	−75.69 (14)	C10—C11—C12—N3	−10.31 (14)
C14—N3—C15—C10	171.01 (12)	C6—N1—C7—C8	0.44 (16)
C12—N3—C15—C10	39.64 (13)	C9—C8—C7—N1	176.60 (13)
N3—C15—C10—C13	72.92 (12)	C1—C8—C7—N1	−0.36 (15)
C16—C15—C10—C13	−51.98 (14)	C17—C16—C21—C20	−2.7 (2)
C22—C15—C10—C13	−165.45 (11)	C15—C16—C21—C20	177.84 (13)
N3—C15—C10—C11	−43.01 (11)	C17—C16—C21—N2	175.99 (12)
C16—C15—C10—C11	−167.91 (11)	C15—C16—C21—N2	−3.47 (15)
C22—C15—C10—C11	78.62 (12)	C22—N2—C21—C20	175.24 (14)
N3—C15—C10—C9	−161.10 (10)	C22—N2—C21—C16	−3.34 (16)
C16—C15—C10—C9	74.00 (14)	C6—C1—C2—C3	−0.7 (2)
C22—C15—C10—C9	−39.47 (14)	C8—C1—C2—C3	178.51 (15)
N3—C15—C16—C17	−45.93 (19)	C21—N2—C22—O2	−170.64 (14)
C22—C15—C16—C17	−171.58 (14)	C21—N2—C22—C15	8.35 (15)
C10—C15—C16—C17	71.86 (19)	N3—C15—C22—O2	44.68 (19)
N3—C15—C16—C21	133.45 (12)	C16—C15—C22—O2	169.35 (14)
C22—C15—C16—C21	7.80 (14)	C10—C15—C22—O2	−66.94 (17)
C10—C15—C16—C21	−108.75 (13)	N3—C15—C22—N2	−134.34 (12)
C13—C10—C11—C23	46.36 (16)	C16—C15—C22—N2	−9.67 (13)
C9—C10—C11—C23	−76.07 (14)	C10—C15—C22—N2	114.04 (12)
C15—C10—C11—C23	161.69 (11)	C1—C2—C3—C4	1.2 (2)
C13—C10—C11—C12	−82.99 (13)	C25—N5—C23—N6	−0.39 (18)
C9—C10—C11—C12	154.58 (11)	C25—N5—C23—C11	−178.58 (14)
C15—C10—C11—C12	32.35 (12)	C24—N6—C23—N5	0.46 (17)
C7—N1—C6—C5	179.16 (15)	C24—N6—C23—C11	178.79 (12)
C7—N1—C6—C1	−0.33 (16)	C12—C11—C23—N5	41.7 (2)
C2—C1—C6—N1	179.51 (12)	C10—C11—C23—N5	−82.13 (18)
C8—C1—C6—N1	0.10 (15)	C12—C11—C23—N6	−136.32 (14)
C2—C1—C6—C5	0.0 (2)	C10—C11—C23—N6	99.85 (15)
C8—C1—C6—C5	−179.43 (13)	N1—C6—C5—C4	−179.14 (15)
C13—C10—C9—O1	−129.88 (13)	C1—C6—C5—C4	0.3 (2)
C11—C10—C9—O1	−7.51 (16)	C16—C21—C20—C19	0.3 (2)
C15—C10—C9—O1	104.60 (14)	N2—C21—C20—C19	−178.08 (15)
C13—C10—C9—C8	50.01 (16)	C18—C19—C20—C21	1.5 (3)
C11—C10—C9—C8	172.37 (11)	C6—C5—C4—C3	0.2 (3)
C15—C10—C9—C8	−75.52 (15)	C2—C3—C4—C5	−0.9 (3)

O1—C9—C8—C7	−176.01 (14)	C21—C16—C17—C18	3.1 (2)
C10—C9—C8—C7	4.1 (2)	C15—C16—C17—C18	−177.58 (14)
O1—C9—C8—C1	0.5 (2)	C20—C19—C18—C17	−1.1 (3)
C10—C9—C8—C1	−179.41 (11)	C16—C17—C18—C19	−1.3 (2)
C2—C1—C8—C7	−179.12 (15)	C23—N5—C25—C24	0.2 (2)
C6—C1—C8—C7	0.15 (14)	N5—C25—C24—N6	0.1 (2)
C2—C1—C8—C9	3.7 (2)	C23—N6—C24—C25	−0.32 (19)
C6—C1—C8—C9	−177.02 (12)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1 <i>A</i> ···N5 ⁱ	0.89 (1)	2.13 (1)	2.9889 (19)	164 (2)
N6—H6 <i>A</i> ···O1 <i>W</i>	0.90 (1)	1.98 (2)	2.714 (8)	138 (2)
N6—H6 <i>A</i> ···O1	0.90 (1)	2.57 (2)	3.064 (2)	116 (2)

Symmetry code: (i) $x+1/2, -y+1/2, z-1/2$.