

N-(2-Hydroxyphenyl)-4-methylbenzene-sulfonamide

Shaaban K. Mohamed,^{a,b} Mehmet Akkurt,^c Benson M. Kariuki,^d Ali M. Ali^e and Mustafa R. Albayati^{f*}

^aChemistry and Environmental Division, Manchester Metropolitan University, Manchester M1 5GD, England, ^bChemistry Department, Faculty of Science, Minia University, 61519 El-Minia, Egypt, ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, ^dSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT, Wales, ^eDepartment of Chemistry, Faculty of Science, Sohag University, 82524 Sohag, Egypt, and ^fKirkuk University, College of Science, Department of Chemistry, Kirkuk, Iraq
Correspondence e-mail: shaabankamel@yahoo.com


Received 9 December 2013; accepted 9 December 2013

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.030; wR factor = 0.082; data-to-parameter ratio = 13.8.

In the title compound, $\text{C}_{13}\text{H}_{13}\text{NO}_3\text{S}$, the dihedral angle between the benzene rings is $64.15(7)^\circ$ and the $\text{C}-\text{S}-\text{N}-\text{C}$ torsion angle is $-57.18(12)^\circ$. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond closes an $S(5)$ ring. In the crystal, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into $C(8)$ chains propagating in [100]. Weak $\text{C}-\text{H}\cdots\pi$ interactions are also observed.

Related literature

For background to the biological activity of sulfonamide compounds, see: Ozbek *et al.* (2007); El-Sayed *et al.* (2011). For related structures, see: Gowda *et al.* (2008a,b,c).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{13}\text{NO}_3\text{S}$

$M_r = 263.31$

Monoclinic, $P2_1/c$

$a = 7.6780(1)\text{ \AA}$

$b = 15.4747(3)\text{ \AA}$

$c = 10.7250(2)\text{ \AA}$

$\beta = 104.333(2)^\circ$

$V = 1234.62(4)\text{ \AA}^3$

$Z = 4$

$\text{Cu } K\alpha$ radiation

$\mu = 2.34\text{ mm}^{-1}$

$T = 120\text{ K}$

$0.35 \times 0.16 \times 0.13\text{ mm}$

Data collection

Oxford Diffraction SuperNova

(Dual, Cu at zero, Atlas)
diffractometer

Absorption correction: multi-scan
(*CrysAlis PRO*; Oxford)

Diffractometer, 2013)
 $T_{\min} = 0.494$, $T_{\max} = 0.750$
4355 measured reflections
2377 independent reflections
2248 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.011$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.082$

$S = 1.06$

2377 reflections

172 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.38\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ and $Cg2$ are the centroids of the C1–C6 and C8–C13 benzene rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}1-\text{H}1\text{N}\cdots\text{O}3$	0.83 (2)	2.22 (2)	2.6420 (16)	111.6 (17)
$\text{O}3-\text{H}1\text{O}\cdots\text{O}2^i$	0.86 (2)	1.94 (2)	2.7852 (15)	172 (2)
$\text{C}3-\text{H}3\cdots\text{C}g2^{ii}$	0.95	2.92	3.8022 (16)	155
$\text{C}7-\text{H}7\text{C}\cdots\text{C}g1^{iii}$	0.98	2.85	3.5937 (17)	134

Symmetry codes: (i) $x - 1, y, z$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $-x, -y + 1, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

We thank Manchester Metropolitan University, Erciyes University and Cardiff University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB7173).

References

- El-Sayed, N. S., El-Bendary, E. R., El-Ashry, S. M. & El-Kerdawy, M. M. (2011). *Eur. J. Med. Chem.* **46**, 3714–3720.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2008a). *Acta Cryst. E* **64**, o1691.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2008b). *Acta Cryst. E* **64**, o1825.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2008c). *Acta Cryst. E* **64**, o2190.
- Oxford Diffraction (2013). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Ozbek, N., Katircioğlu, H., Karacan, N. & Baykal, T. (2007). *Bioorg. Med. Chem.* **15**, 5105–5109.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2014). E70, o54 [https://doi.org/10.1107/S1600536813033394]

N-(2-Hydroxyphenyl)-4-methylbenzenesulfonamide

Shaaban K. Mohamed, Mehmet Akkurt, Benson M. Kariuki, Ali M. Ali and Mustafa R. Albayati

S1. Comment

The biological activities of sulphonamide compounds are well documented, for example as antimicrobial (Ozbek *et al.*, 2007) and anticancer (El-Sayed *et al.*, 2011) agents. Further to our interest in related compounds with potential biactivity, we now report the synthesis and crystal structure of the title compound.

The benzene rings (C1–C6 and C8–C13) of the title compound (I) in Fig. 1 make a dihedral angle of 64.15 (7) $^{\circ}$ with each other. The bridge C1—S1—N1—C8 torsion angle between the benzene rings is -57.18 (12) $^{\circ}$. The O1—S1—O2 and C1—S1—N1 angles are 119.78 (6) and 107.97 (6) $^{\circ}$, respectively. The bond lengths and angles are similar to those in related structures (Gowda *et al.*, 2008*a,b,c*).


The molecular conformation features an N—H \cdots O hydrogen bond which forms an S(5) ring (Fig. 2). In the crystal, molecules are linked by O—H \cdots O hydrogen bonds into C(8) chains along [100] (Figs. 2 and 3). Weak C—H \cdots π interactions are also observed (Table 1).

S2. Experimental


A mixture of 2-aminophenol (109 mg, 1 mmol) and *p*-toluenesulfonyl chloride (190 mg, 1 mmol) in 10 ml dioxane with addition of few drops of triethylamine as a catalyst, was refluxed for 4 h. The reaction mixture was left to cool at ambient temperature where the solid product was deposited, collected by filtration and recrystallized from ethanol in 91% yield. Brown needles were grown from ethanol solution over 3 days at room temperature. *M.p.* 391 K.

S3. Refinement


The H atoms of the NH and OH groups were found from difference Fourier maps and refined freely. The C-bound H atoms were positioned geometrically, with C—H = 0.95 and 0.98 Å and refined as riding with $U_{\text{iso}}(\text{H}) = 1.U_{\text{eq}}(\text{C})$ for the methyl H atoms and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for the other H atoms.

**Figure 1**

View of the title compound with displacement ellipsoids for non-H atoms drawn at the 50% probability level.

**Figure 2**

View of the hydrogen bonds along the a axis direction of the title compound. H bonds are shown as dashed lines.

**Figure 3**

View of the molecular packing along the *a* axis of the title compound. H bonds are shown as dashed lines.

N-(2-Hydroxyphenyl)-4-methylbenzenesulfonamide

Crystal data



$$M_r = 263.31$$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$$a = 7.6780 (1) \text{ \AA}$$

$$b = 15.4747 (3) \text{ \AA}$$

$$c = 10.7250 (2) \text{ \AA}$$

$$\beta = 104.333 (2)^\circ$$

$$V = 1234.62 (4) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 552$$

$$D_x = 1.417 \text{ Mg m}^{-3}$$

Cu $K\alpha$ radiation, $\lambda = 1.54180 \text{ \AA}$

Cell parameters from 2248 reflections

$$\theta = 5.1\text{--}73.2^\circ$$

$$\mu = 2.34 \text{ mm}^{-1}$$

$$T = 120 \text{ K}$$

Needle, brown

$$0.35 \times 0.16 \times 0.13 \text{ mm}$$

Data collection

Oxford Diffraction SuperNova (Dual, Cu at zero, Atlas) diffractometer	$T_{\min} = 0.494, T_{\max} = 0.750$ 4355 measured reflections 2377 independent reflections 2248 reflections with $I > 2\sigma(I)$
Radiation source: SuperNova (Cu) X-ray Source	$R_{\text{int}} = 0.011$
Mirror monochromator	$\theta_{\max} = 73.2^\circ, \theta_{\min} = 5.1^\circ$
ω scans	$h = -8 \rightarrow 9$
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2013)	$k = -19 \rightarrow 12$ $l = -11 \rightarrow 13$

Refinement

Refinement on F^2	Hydrogen site location: mixed
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.030$	$W = 1/[\Sigma^2(FO^2) + (0.0424P)^2 + 0.5981P]$ where $P = (FO^2 + 2FC^2)/3$
$wR(F^2) = 0.082$	$(\Delta/\sigma)_{\max} = 0.001$
$S = 1.06$	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
2377 reflections	$\Delta\rho_{\min} = -0.38 \text{ e } \text{\AA}^{-3}$
172 parameters	
0 restraints	

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors.

Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating - R -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.51287 (4)	0.65871 (2)	0.83597 (3)	0.0203 (1)
O1	0.57618 (13)	0.61174 (7)	0.95358 (10)	0.0281 (3)
O2	0.63687 (13)	0.70958 (7)	0.78545 (10)	0.0267 (3)
O3	0.00824 (14)	0.71635 (7)	0.81974 (10)	0.0264 (3)
N1	0.36177 (15)	0.72621 (8)	0.86360 (11)	0.0208 (3)
C1	0.40408 (17)	0.58685 (9)	0.71468 (13)	0.0196 (4)
C2	0.3833 (2)	0.60851 (9)	0.58584 (14)	0.0239 (4)
C3	0.2974 (2)	0.55124 (10)	0.49170 (14)	0.0268 (4)
C4	0.23172 (19)	0.47243 (9)	0.52309 (14)	0.0247 (4)
C5	0.2521 (2)	0.45305 (10)	0.65261 (15)	0.0278 (4)
C6	0.3373 (2)	0.50922 (10)	0.74888 (14)	0.0253 (4)
C7	0.1447 (2)	0.40915 (11)	0.42031 (17)	0.0343 (5)
C8	0.25492 (18)	0.77847 (9)	0.76315 (13)	0.0192 (3)
C9	0.3290 (2)	0.83739 (9)	0.69307 (15)	0.0243 (4)
C10	0.2174 (2)	0.89020 (9)	0.60237 (15)	0.0273 (4)
C11	0.0320 (2)	0.88327 (10)	0.58080 (14)	0.0265 (4)
C12	-0.04291 (19)	0.82447 (10)	0.65038 (14)	0.0234 (4)

C13	0.06818 (18)	0.77298 (9)	0.74268 (13)	0.0202 (3)
H1N	0.301 (3)	0.7023 (13)	0.9076 (19)	0.035 (5)*
H1O	-0.106 (3)	0.7191 (15)	0.807 (2)	0.051 (6)*
H2	0.42760	0.66200	0.56290	0.0290*
H3	0.28280	0.56590	0.40370	0.0320*
H5	0.20630	0.39990	0.67550	0.0330*
H6	0.35000	0.49500	0.83680	0.0300*
H7A	0.22220	0.35830	0.42470	0.0510*
H7B	0.12710	0.43650	0.33560	0.0510*
H7C	0.02810	0.39140	0.43350	0.0510*
H9	0.45580	0.84160	0.70710	0.0290*
H10	0.26800	0.93100	0.55520	0.0330*
H11	-0.04390	0.91890	0.51810	0.0320*
H12	-0.16980	0.81950	0.63480	0.0280*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0117 (2)	0.0246 (2)	0.0243 (2)	0.0002 (1)	0.0036 (1)	-0.0013 (1)
O1	0.0198 (5)	0.0360 (6)	0.0254 (5)	0.0033 (4)	-0.0005 (4)	0.0019 (4)
O2	0.0140 (5)	0.0298 (5)	0.0380 (6)	-0.0027 (4)	0.0096 (4)	-0.0031 (4)
O3	0.0146 (5)	0.0354 (6)	0.0298 (5)	-0.0019 (4)	0.0068 (4)	0.0060 (4)
N1	0.0150 (5)	0.0253 (6)	0.0229 (6)	-0.0006 (5)	0.0065 (5)	-0.0022 (5)
C1	0.0154 (6)	0.0207 (6)	0.0233 (7)	0.0027 (5)	0.0059 (5)	-0.0003 (5)
C2	0.0258 (7)	0.0208 (7)	0.0269 (7)	0.0013 (5)	0.0101 (6)	0.0031 (5)
C3	0.0308 (8)	0.0275 (7)	0.0232 (7)	0.0035 (6)	0.0089 (6)	0.0008 (6)
C4	0.0197 (6)	0.0247 (7)	0.0302 (7)	0.0040 (6)	0.0071 (6)	-0.0042 (6)
C5	0.0280 (8)	0.0210 (7)	0.0359 (8)	-0.0025 (6)	0.0107 (6)	0.0019 (6)
C6	0.0264 (7)	0.0254 (7)	0.0247 (7)	0.0003 (6)	0.0074 (6)	0.0047 (6)
C7	0.0288 (8)	0.0334 (8)	0.0404 (9)	-0.0004 (7)	0.0081 (7)	-0.0130 (7)
C8	0.0171 (6)	0.0196 (6)	0.0211 (6)	-0.0004 (5)	0.0052 (5)	-0.0056 (5)
C9	0.0206 (7)	0.0230 (7)	0.0319 (7)	-0.0026 (5)	0.0112 (6)	-0.0047 (6)
C10	0.0317 (8)	0.0223 (7)	0.0313 (8)	-0.0011 (6)	0.0144 (6)	0.0009 (6)
C11	0.0292 (8)	0.0243 (7)	0.0260 (7)	0.0048 (6)	0.0069 (6)	0.0001 (6)
C12	0.0178 (6)	0.0268 (7)	0.0255 (7)	0.0018 (5)	0.0051 (5)	-0.0041 (6)
C13	0.0186 (6)	0.0213 (6)	0.0220 (6)	-0.0023 (5)	0.0076 (5)	-0.0050 (5)

Geometric parameters (\AA , ^\circ)

S1—O1	1.4325 (11)	C8—C9	1.390 (2)
S1—O2	1.4405 (11)	C9—C10	1.391 (2)
S1—N1	1.6417 (12)	C10—C11	1.389 (2)
S1—C1	1.7574 (14)	C11—C12	1.389 (2)
O3—C13	1.3606 (18)	C12—C13	1.387 (2)
O3—H1O	0.86 (2)	C2—H2	0.9500
N1—C8	1.4318 (18)	C3—H3	0.9500
N1—H1N	0.83 (2)	C5—H5	0.9500
C1—C6	1.391 (2)	C6—H6	0.9500

C1—C2	1.392 (2)	C7—H7A	0.9800
C2—C3	1.382 (2)	C7—H7B	0.9800
C3—C4	1.393 (2)	C7—H7C	0.9800
C4—C7	1.502 (2)	C9—H9	0.9500
C4—C5	1.392 (2)	C10—H10	0.9500
C5—C6	1.384 (2)	C11—H11	0.9500
C8—C13	1.398 (2)	C12—H12	0.9500
O1—S1—O2	119.78 (6)	C11—C12—C13	119.77 (14)
O1—S1—N1	105.36 (6)	O3—C13—C8	115.57 (12)
O1—S1—C1	109.04 (6)	O3—C13—C12	124.25 (13)
O2—S1—N1	106.42 (6)	C8—C13—C12	120.17 (13)
O2—S1—C1	107.74 (6)	C1—C2—H2	120.00
N1—S1—C1	107.97 (6)	C3—C2—H2	120.00
C13—O3—H1O	111.0 (15)	C2—C3—H3	119.00
S1—N1—C8	121.50 (9)	C4—C3—H3	119.00
C8—N1—H1N	112.5 (15)	C4—C5—H5	119.00
S1—N1—H1N	110.0 (15)	C6—C5—H5	119.00
S1—C1—C6	119.35 (11)	C1—C6—H6	121.00
S1—C1—C2	119.94 (11)	C5—C6—H6	121.00
C2—C1—C6	120.71 (13)	C4—C7—H7A	109.00
C1—C2—C3	119.17 (13)	C4—C7—H7B	109.00
C2—C3—C4	121.39 (14)	C4—C7—H7C	109.00
C5—C4—C7	120.78 (13)	H7A—C7—H7B	109.00
C3—C4—C5	118.17 (13)	H7A—C7—H7C	109.00
C3—C4—C7	121.03 (13)	H7B—C7—H7C	109.00
C4—C5—C6	121.67 (14)	C8—C9—H9	120.00
C1—C6—C5	118.87 (13)	C10—C9—H9	120.00
N1—C8—C9	122.85 (13)	C9—C10—H10	120.00
N1—C8—C13	117.26 (12)	C11—C10—H10	120.00
C9—C8—C13	119.73 (13)	C10—C11—H11	120.00
C8—C9—C10	119.99 (14)	C12—C11—H11	120.00
C9—C10—C11	119.98 (14)	C11—C12—H12	120.00
C10—C11—C12	120.33 (14)	C13—C12—H12	120.00
O1—S1—N1—C8	-173.59 (11)	C2—C3—C4—C7	-177.69 (15)
O2—S1—N1—C8	58.25 (12)	C2—C3—C4—C5	1.0 (2)
C1—S1—N1—C8	-57.18 (12)	C3—C4—C5—C6	-1.0 (2)
O1—S1—C1—C2	-158.83 (12)	C7—C4—C5—C6	177.79 (15)
O2—S1—C1—C2	-27.38 (14)	C4—C5—C6—C1	-0.1 (2)
N1—S1—C1—C2	87.19 (13)	N1—C8—C9—C10	-175.85 (13)
O1—S1—C1—C6	22.08 (14)	C13—C8—C9—C10	-0.4 (2)
O2—S1—C1—C6	153.53 (12)	N1—C8—C13—O3	-1.43 (18)
N1—S1—C1—C6	-91.91 (13)	N1—C8—C13—C12	177.40 (13)
S1—N1—C8—C9	-59.30 (17)	C9—C8—C13—O3	-177.10 (13)
S1—N1—C8—C13	125.18 (12)	C9—C8—C13—C12	1.7 (2)
S1—C1—C6—C5	-179.91 (12)	C8—C9—C10—C11	-0.8 (2)
S1—C1—C2—C3	-180.00 (12)	C9—C10—C11—C12	0.7 (2)

C6—C1—C2—C3	−0.9 (2)	C10—C11—C12—C13	0.6 (2)
C2—C1—C6—C5	1.0 (2)	C11—C12—C13—O3	176.92 (13)
C1—C2—C3—C4	−0.1 (2)	C11—C12—C13—C8	−1.8 (2)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C1—C6 and C8—C13 benzene rings, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O3	0.83 (2)	2.22 (2)	2.6420 (16)	111.6 (17)
O3—H1O···O2 ⁱ	0.86 (2)	1.94 (2)	2.7852 (15)	172 (2)
C9—H9···O2	0.95	2.50	3.0531 (18)	117
C3—H3···Cg2 ⁱⁱ	0.95	2.92	3.8022 (16)	155
C7—H7C···Cg1 ⁱⁱⁱ	0.98	2.85	3.5937 (17)	134

Symmetry codes: (i) $x-1, y, z$; (ii) $x, -y+3/2, z-1/2$; (iii) $-x, -y+1, -z+1$.