

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl (2Z)-2-[(2Z)-3-[(cyclopentylidene)amino]-4-oxo-2-phenylimino-1,3-thiazolidin-5-ylidene]acetate

Joel T. Mague,^a Mehmet Akkurt,^b Shaaban K. Mohamed,^{c,d}
Alaa A. Hassan^d and Mustafa R. Albayati^{e*}

^aDepartment of Chemistry, Tulane University, New Orleans, LA 70118, USA,^bDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri,Turkey, ^cChemistry and Environmental Division, Manchester MetropolitanUniversity, Manchester M1 5GD, England, ^dChemistry Department, Faculty ofScience, Minia University, 61519 El-Minia, Egypt, and ^eKirkuk University, College of

Science, Department of Chemistry, Kirkuk, Iraq

Correspondence e-mail: shaabankamel@yahoo.com


Received 19 February 2014; accepted 20 February 2014

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å;
R factor = 0.029; wR factor = 0.075; data-to-parameter ratio = 13.5.

In the title compound, $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_3\text{S}$, the thiazole ring is nearly planar [maximum deviation = 0.015 (1) Å for the ring N atom] and the cyclopentane ring has a twist conformation. The molecular conformation is stabilized by a hypervalent interaction between the S atom and the ester group carbonyl O atom, with an $\text{S}\cdots\text{O}$ distance of 2.7931 (10) Å. In the crystal, $\text{C}-\text{H}\cdots\text{O}$ interactions generate chains of molecules propagating along [110] and $\pi-\pi$ stacking interactions [centroid-centroid distance = 3.4677 (7) Å] between the thiazole rings organize these chains into (001) layers.

Related literature

For the synthesis and similar structures, see: Akkurt *et al.* (2009); Li *et al.* (2011); Mague *et al.* (2013); Mohamed *et al.* (2013a,b); Pomés Hernández *et al.* (1996); Sundar *et al.* (2003). For the general biological significance of thiazolidinone scaffold compounds, see: Pfützner *et al.* (2007); Schianca *et al.* (2012); Jain *et al.* (2012); Lant (1986); Rock *et al.* (1991).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_3\text{S}$ $M_r = 343.41$ Monoclinic, $P2_1/c$ $a = 9.9684$ (2) Å $b = 9.9657$ (2) Å $c = 16.9818$ (3) Å $\beta = 105.9290$ (6)° $V = 1622.23$ (5) Å³ $Z = 4$ Cu $K\alpha$ radiation $\mu = 1.96$ mm⁻¹ $T = 100$ K $0.17 \times 0.16 \times 0.09$ mm

Data collection

Bruker D8 VENTURE PHOTON

100 CMOS diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2013)

 $T_{\min} = 0.76$, $T_{\max} = 0.84$

17510 measured reflections

2951 independent reflections

2769 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.023$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.029$ $wR(F^2) = 0.075$ $S = 1.08$

2951 reflections

218 parameters

61 restraints

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.30$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|-------|-------------|-------------|---------------|
| $\text{C10}-\text{H10B}\cdots\text{O1}^i$ | 0.99 | 2.57 | 3.2889 (17) | 130 |
| $\text{C11}-\text{H11A}\cdots\text{O1}^i$ | 0.99 | 2.58 | 3.2547 (16) | 125 |

Symmetry code: (i) $-x + 1, -y + 2, -z + 1$.

Data collection: APEX2 (Bruker, 2013); cell refinement: SAINT (Bruker, 2013); data reduction: SAINT; program(s) used to solve structure: SHELXS2013 (Sheldrick, 2008); program(s) used to refine structure: SHELXL2013 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: WinGX (Farrugia, 2012) and PLATON (Spek, 2009).

Manchester Metropolitan University, Tulane University and Erciyes University are gratefully acknowledged for supporting this study as is the support of NSF-MRI grant No. 1228232 for the purchase of the diffractometer.

Supporting information for this paper is available from the IUCr electronic archives (Reference: GK2605).

References

- Akkurt, M., Karaca, S., Cihan, G., Çapan, G. & Büyükgüngör, O. (2009). *Acta Cryst.* **E65**, o1009–o1010.
- Bruker (2013). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Jain, A. K., Vaidya, A., Ravichandran, V., Kashaw, S. K. & Agrawal, R. K. (2012). *Bioorg. Med. Chem.* **20**, 3378–95.
- Lant, A. (1986). *Drugs*, **31**, 40–55.
- Li, Y.-M., Li, J.-Y., Wang, Q., Hao, A.-Y. & Sun, T. (2011). *Acta Cryst.* **E67**, o3303.
- Mague, J. T., Akkurt, M., Mohamed, S. K., Hassan, A. A. & Albayati, M. R. (2013). *Acta Cryst.* **E69**, o1401–o1402.
- Mohamed, S. K., Akkurt, M., Mague, J. T., Hassan, A. A. & Albayati, M. R. (2013b). *Acta Cryst.* **E69**, o1844–o1845.
- Mohamed, S. K., Mague, J. T., Akkurt, M., Hassan, A. A. & Albayati, M. R. (2013a). *Acta Cryst.* **E69**, o1553–o1554.
- Pfützner, A., Weber, M. M. & Forst, T. (2007). *Expert Opin. Pharmacother.* **8**, 1985–98.
- Pomés Hernández, R., Duque Rodríguez, J., Novoa de Armas, H. & Toscano, R. A. (1996). *Acta Cryst.* **C52**, 1731–1733.
- Rock, D. R., McLean, M. J., Macdonald, R. L., Catterall, W. A. & Taylor, C. P. (1991). *Epilepsy Res.* **8**, 197–203.
- Schianca, G. P. C., Sola, D., Rossi, L., Fra, G. P. & Bartoli, E. (2012). *ISRN Endocrinology*, Article ID 601380, 1–6.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sundar, T. V., Parthasarathi, V., García-Granda, S., Jain, A. & Pardasani, R. T. (2003). *Acta Cryst.* **E59**, o1967–o1969.

supporting information

Acta Cryst. (2014). E70, o366–o367 [doi:10.1107/S1600536814004048]

Methyl (2Z)-2-[(2Z)-3-[(cyclopentylidene)amino]-4-oxo-2-phenylimino-1,3-thiazolidin-5-ylidene]acetate

Joel T. Mague, Mehmet Akkurt, Shaaban K. Mohamed, Alaa A. Hassan and Mustafa R. Albayati

S1. Comment

Diversity in the biological response profile of thiazolidinone and analogous scaffolds has attracted much attention to the exploration of this skeleton for a variety of therapeutic applications. Thus, the successful pharmaceutical applications of pioglitazone as a hypoglycemic agent (Pfützner *et al.*, 2007; Schianca *et al.*, 2012), thiazolidomycin activity against streptomyces species (Jain *et al.*, 2012), etozoline as an antihypertensive (Lant, 1986), and ralitoline as a potent anti-convulsant (Rock *et al.*, 1991) have established the wide spectrum potential of the thiazolidinone moiety. As part of our ongoing program of drug design and discovery, we report the synthesis and crystal structure of the title compound.

As shown in Fig. 1, the thiazole ring (S1/N1/C1–C3) of the title compound (I) is nearly planar with a maximum deviation of 0.015 (1) Å for N1. The cyclopentane ring (C7–C11) is twisted around the C9–C10 bond. The dihedral angle between the thiazole and phenyl rings is 53.84 (6)°. The C1–C3–C4–C5, C3–C4–C5–O2, C3–C4–C5–O3, C4–C5–O3–C6 and O2–C5–O3–C6 torsion angles are 178.90 (12), 6.7 (2), -174.42 (12), -178.11 (11) and 2.99 (19)°, respectively. All the bond lengths in (I) are comparable to those observed in similar structures (Akkurt *et al.*, 2009; Li *et al.*, 2011; Mague *et al.*, 2013; Mohamed *et al.*, 2013*a,b*; Pomés Hernández *et al.*, 1996; Sundar *et al.*, 2003).

Molecular conformation of (I) is stabilized by a short intramolecular (hypervalent) contact between the S1 atom and the ester group carbonyl O2 atom of 2.7931 (10) Å. In the crystal packing, the C—H···O interactions (Table 1, Fig. 2) with H···O distance of 2.57 - 2.58 Å generate chains of molecules propagating along the [110] direction. Furthermore, π - π stacking interactions [$Cg1 \cdots Cg1$ (1 - x , 1 - y , 1 - z) = 3.4677 (7) Å; where $Cg1$ is a centroid of the S1/N1/C1–C3 ring] between the thiazole rings organize these chains into the (001) layers.

S2. Experimental

A solution of 1 mmol (233 mg) 2-cyclopentylidene-*N*-phenylhydrazinecarbothioamide in 15 ml ethanol was added dropwise to a solution of 1 mmol (142 mg) dimethyl but-2-ynedioate in 10 ml ethanol. The reaction mixture was stirred and refluxed at 351 K. The reaction progress was monitored by TLC until completion. On cooling a solid yellow product was precipitated, filtered off under vacuum and recrystallized from ethanol to furnish block-shaped yellow crystals (m.p. 541–543 K).

S3. Refinement

All H atoms were positioned geometrically and treated as riding atoms, with C—H = 0.95 Å (aromatic H), 0.98 Å (methyl H) and 0.99 Å (methylene H), with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{iso}}(\text{C})$ for methyl H atoms and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{iso}}(\text{C})$ for the others. The components of the displacement parameters in the direction of the bond between non-hydrogen atoms were restrained to be equal within an effective standard deviation of 0.01 (DELU instruction).




Figure 1

Perspective view of the title compound with 50% probability displacement ellipsoids.




Figure 2

Crystal packing of the title compound viewed down the *a* axis. Only hydrogen atoms involved in C-H...O interactions (dashed lines) are shown.

Methyl (2*Z*)-2-[(2*Z*)-3-[(cyclopentylidene)amino]-4-oxo-2-phenylimino-1,3-thiazolidin-5-ylidene]acetate

Crystal data

$C_{17}H_{17}N_3O_3S$

$M_r = 343.41$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 9.9684\ (2)\ \text{\AA}$

$b = 9.9657\ (2)\ \text{\AA}$

$c = 16.9818\ (3)\ \text{\AA}$

$\beta = 105.9290\ (6)^\circ$

$V = 1622.23\ (5)\ \text{\AA}^3$

$Z = 4$

$F(000) = 720$

$D_x = 1.406\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54178\ \text{\AA}$

Cell parameters from 9943 reflections

$\theta = 4.4\text{--}68.3^\circ$

$\mu = 1.96\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Block, yellow

$0.17 \times 0.16 \times 0.09\ \text{mm}$

Data collection

Bruker D8 VENTURE PHOTON 100 CMOS
diffractometer
Radiation source: INCOATEC I μ S micro-focus
source
Mirror monochromator
Detector resolution: 10.4167 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2013)

$T_{\min} = 0.76$, $T_{\max} = 0.84$
17510 measured reflections
2951 independent reflections
2769 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\max} = 68.2^\circ$, $\theta_{\min} = 4.6^\circ$
 $h = -12 \rightarrow 12$
 $k = -11 \rightarrow 12$
 $l = -20 \rightarrow 20$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.075$
 $S = 1.08$
2951 reflections
218 parameters
61 restraints

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.038P)^2 + 0.6771P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.30 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.21 \text{ e } \text{Å}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|--------------|----------------------------------|
| S1 | 0.49073 (3) | 0.41667 (3) | 0.35998 (2) | 0.0187 (1) |
| O1 | 0.66547 (10) | 0.71951 (9) | 0.49052 (6) | 0.0244 (3) |
| O2 | 0.72709 (10) | 0.25889 (9) | 0.37200 (6) | 0.0258 (3) |
| O3 | 0.93931 (9) | 0.30447 (9) | 0.45742 (6) | 0.0238 (3) |
| N1 | 0.44614 (11) | 0.65036 (10) | 0.41830 (6) | 0.0190 (3) |
| N2 | 0.36915 (11) | 0.75447 (11) | 0.44449 (7) | 0.0214 (3) |
| N3 | 0.24379 (11) | 0.55652 (11) | 0.33548 (7) | 0.0204 (3) |
| C1 | 0.58785 (13) | 0.63691 (13) | 0.44889 (8) | 0.0193 (3) |
| C2 | 0.37387 (13) | 0.54855 (12) | 0.36801 (8) | 0.0184 (3) |
| C3 | 0.63174 (13) | 0.50466 (13) | 0.42232 (7) | 0.0186 (3) |
| C4 | 0.76427 (13) | 0.46352 (13) | 0.44650 (8) | 0.0202 (4) |
| C5 | 0.80432 (13) | 0.33216 (13) | 0.42077 (8) | 0.0207 (4) |
| C6 | 0.98665 (15) | 0.17461 (14) | 0.43752 (9) | 0.0276 (4) |
| C7 | 0.36894 (13) | 0.86381 (13) | 0.40585 (8) | 0.0202 (4) |
| C8 | 0.43371 (17) | 0.89572 (14) | 0.33790 (9) | 0.0298 (4) |
| C9 | 0.41616 (18) | 1.04928 (15) | 0.32757 (11) | 0.0364 (5) |
| C10 | 0.28241 (16) | 1.07780 (14) | 0.35110 (9) | 0.0300 (4) |

| | | | | |
|------|---------------|--------------|-------------|------------|
| C11 | 0.29241 (14) | 0.98521 (13) | 0.42434 (8) | 0.0241 (4) |
| C12 | 0.17495 (13) | 0.45669 (13) | 0.27910 (8) | 0.0209 (3) |
| C13 | 0.05398 (14) | 0.39885 (14) | 0.28986 (9) | 0.0249 (4) |
| C14 | -0.01893 (15) | 0.30450 (15) | 0.23424 (9) | 0.0283 (4) |
| C15 | 0.02709 (15) | 0.26783 (14) | 0.16740 (9) | 0.0291 (4) |
| C16 | 0.14570 (15) | 0.32725 (16) | 0.15559 (9) | 0.0300 (4) |
| C17 | 0.21879 (14) | 0.42310 (14) | 0.21052 (8) | 0.0252 (4) |
| H4 | 0.83320 | 0.51970 | 0.48060 | 0.0240* |
| H6A | 0.94260 | 0.10380 | 0.46180 | 0.0410* |
| H6B | 1.08820 | 0.16880 | 0.45940 | 0.0410* |
| H6C | 0.96120 | 0.16350 | 0.37790 | 0.0410* |
| H8A | 0.38460 | 0.84850 | 0.28680 | 0.0360* |
| H8B | 0.53350 | 0.87020 | 0.35310 | 0.0360* |
| H9A | 0.49630 | 1.09710 | 0.36430 | 0.0440* |
| H9B | 0.40730 | 1.07640 | 0.27030 | 0.0440* |
| H10A | 0.19940 | 1.05620 | 0.30550 | 0.0360* |
| H10B | 0.27760 | 1.17310 | 0.36660 | 0.0360* |
| H11A | 0.34520 | 1.02850 | 0.47600 | 0.0290* |
| H11B | 0.19850 | 0.96050 | 0.42870 | 0.0290* |
| H13 | 0.02130 | 0.42400 | 0.33530 | 0.0300* |
| H14 | -0.10100 | 0.26480 | 0.24210 | 0.0340* |
| H15 | -0.02240 | 0.20230 | 0.12990 | 0.0350* |
| H16 | 0.17740 | 0.30250 | 0.10970 | 0.0360* |
| H17 | 0.29840 | 0.46550 | 0.20120 | 0.0300* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|------------|------------|-------------|------------|-------------|
| S1 | 0.0170 (2) | 0.0147 (2) | 0.0233 (2) | 0.0017 (1) | 0.0036 (1) | -0.0023 (1) |
| O1 | 0.0223 (5) | 0.0191 (5) | 0.0305 (5) | -0.0023 (4) | 0.0053 (4) | -0.0060 (4) |
| O2 | 0.0216 (5) | 0.0233 (5) | 0.0304 (5) | 0.0028 (4) | 0.0037 (4) | -0.0056 (4) |
| O3 | 0.0179 (5) | 0.0226 (5) | 0.0294 (5) | 0.0051 (4) | 0.0040 (4) | -0.0028 (4) |
| N1 | 0.0194 (5) | 0.0141 (5) | 0.0238 (5) | 0.0017 (4) | 0.0063 (4) | -0.0014 (4) |
| N2 | 0.0222 (6) | 0.0160 (5) | 0.0274 (6) | 0.0036 (4) | 0.0092 (4) | -0.0026 (4) |
| N3 | 0.0181 (5) | 0.0181 (5) | 0.0245 (6) | 0.0016 (4) | 0.0051 (4) | 0.0009 (4) |
| C1 | 0.0200 (6) | 0.0176 (6) | 0.0211 (6) | 0.0003 (5) | 0.0072 (5) | 0.0012 (5) |
| C2 | 0.0210 (6) | 0.0146 (6) | 0.0206 (6) | 0.0016 (5) | 0.0076 (5) | 0.0014 (5) |
| C3 | 0.0210 (6) | 0.0161 (6) | 0.0191 (6) | -0.0013 (5) | 0.0063 (5) | 0.0000 (5) |
| C4 | 0.0188 (6) | 0.0193 (7) | 0.0218 (6) | -0.0009 (5) | 0.0043 (5) | -0.0006 (5) |
| C5 | 0.0178 (6) | 0.0219 (7) | 0.0223 (6) | 0.0016 (5) | 0.0054 (5) | 0.0016 (5) |
| C6 | 0.0247 (7) | 0.0240 (7) | 0.0329 (7) | 0.0085 (6) | 0.0060 (6) | -0.0022 (6) |
| C7 | 0.0170 (6) | 0.0182 (7) | 0.0239 (6) | -0.0004 (5) | 0.0032 (5) | -0.0041 (5) |
| C8 | 0.0371 (8) | 0.0209 (7) | 0.0371 (8) | 0.0065 (6) | 0.0197 (7) | 0.0048 (6) |
| C9 | 0.0487 (10) | 0.0221 (8) | 0.0451 (9) | 0.0068 (7) | 0.0243 (8) | 0.0076 (7) |
| C10 | 0.0359 (8) | 0.0185 (7) | 0.0347 (8) | 0.0058 (6) | 0.0084 (6) | 0.0005 (6) |
| C11 | 0.0253 (7) | 0.0172 (7) | 0.0302 (7) | 0.0034 (5) | 0.0084 (5) | -0.0032 (5) |
| C12 | 0.0180 (6) | 0.0174 (6) | 0.0247 (6) | 0.0037 (5) | 0.0015 (5) | 0.0024 (5) |
| C13 | 0.0220 (7) | 0.0241 (7) | 0.0281 (7) | 0.0000 (5) | 0.0062 (6) | 0.0009 (6) |

| | | | | | | |
|-----|------------|------------|------------|-------------|-------------|-------------|
| C14 | 0.0235 (7) | 0.0249 (7) | 0.0336 (8) | -0.0032 (6) | 0.0029 (6) | 0.0028 (6) |
| C15 | 0.0286 (7) | 0.0230 (7) | 0.0295 (7) | -0.0002 (6) | -0.0025 (6) | -0.0017 (6) |
| C16 | 0.0280 (7) | 0.0351 (8) | 0.0242 (7) | 0.0060 (6) | 0.0024 (6) | -0.0035 (6) |
| C17 | 0.0200 (7) | 0.0279 (7) | 0.0262 (7) | 0.0016 (5) | 0.0038 (5) | 0.0005 (6) |

Geometric parameters (Å, °)

| | | | |
|-----------|-------------|-------------|-----------|
| S1—C2 | 1.7861 (13) | C13—C14 | 1.388 (2) |
| S1—C3 | 1.7461 (13) | C14—C15 | 1.385 (2) |
| O1—C1 | 1.2148 (16) | C15—C16 | 1.385 (2) |
| O2—C5 | 1.2091 (16) | C16—C17 | 1.393 (2) |
| O3—C5 | 1.3471 (16) | C4—H4 | 0.9500 |
| O3—C6 | 1.4489 (17) | C6—H6A | 0.9800 |
| N1—N2 | 1.4320 (15) | C6—H6B | 0.9800 |
| N1—C1 | 1.3713 (17) | C6—H6C | 0.9800 |
| N1—C2 | 1.3929 (16) | C8—H8A | 0.9900 |
| N2—C7 | 1.2717 (17) | C8—H8B | 0.9900 |
| N3—C2 | 1.2648 (18) | C9—H9A | 0.9900 |
| N3—C12 | 1.4201 (17) | C9—H9B | 0.9900 |
| C1—C3 | 1.4968 (18) | C10—H10A | 0.9900 |
| C3—C4 | 1.3358 (19) | C10—H10B | 0.9900 |
| C4—C5 | 1.4696 (18) | C11—H11A | 0.9900 |
| C7—C8 | 1.503 (2) | C11—H11B | 0.9900 |
| C7—C11 | 1.5086 (19) | C13—H13 | 0.9500 |
| C8—C9 | 1.545 (2) | C14—H14 | 0.9500 |
| C9—C10 | 1.520 (2) | C15—H15 | 0.9500 |
| C10—C11 | 1.530 (2) | C16—H16 | 0.9500 |
| C12—C13 | 1.393 (2) | C17—H17 | 0.9500 |
| C12—C17 | 1.3924 (19) | | |
| C2—S1—C3 | 91.01 (6) | O3—C6—H6A | 109.00 |
| C5—O3—C6 | 115.04 (10) | O3—C6—H6B | 109.00 |
| N2—N1—C1 | 122.50 (10) | O3—C6—H6C | 109.00 |
| N2—N1—C2 | 119.15 (11) | H6A—C6—H6B | 109.00 |
| C1—N1—C2 | 117.82 (11) | H6A—C6—H6C | 110.00 |
| N1—N2—C7 | 112.69 (11) | H6B—C6—H6C | 109.00 |
| C2—N3—C12 | 119.89 (11) | C7—C8—H8A | 111.00 |
| O1—C1—N1 | 125.42 (12) | C7—C8—H8B | 111.00 |
| O1—C1—C3 | 125.44 (12) | C9—C8—H8A | 111.00 |
| N1—C1—C3 | 109.14 (11) | C9—C8—H8B | 111.00 |
| S1—C2—N1 | 110.15 (9) | H8A—C8—H8B | 109.00 |
| S1—C2—N3 | 128.67 (10) | C8—C9—H9A | 111.00 |
| N1—C2—N3 | 121.19 (12) | C8—C9—H9B | 111.00 |
| S1—C3—C1 | 111.82 (9) | C10—C9—H9A | 111.00 |
| S1—C3—C4 | 126.58 (10) | C10—C9—H9B | 111.00 |
| C1—C3—C4 | 121.59 (12) | H9A—C9—H9B | 109.00 |
| C3—C4—C5 | 120.56 (12) | C9—C10—H10A | 111.00 |
| O2—C5—O3 | 124.08 (12) | C9—C10—H10B | 111.00 |

| | | | |
|---------------|--------------|-----------------|--------------|
| O2—C5—C4 | 124.60 (12) | C11—C10—H10A | 111.00 |
| O3—C5—C4 | 111.32 (11) | C11—C10—H10B | 111.00 |
| N2—C7—C8 | 129.59 (12) | H10A—C10—H10B | 109.00 |
| N2—C7—C11 | 120.60 (12) | C7—C11—H11A | 111.00 |
| C8—C7—C11 | 109.79 (11) | C7—C11—H11B | 111.00 |
| C7—C8—C9 | 103.75 (12) | C10—C11—H11A | 111.00 |
| C8—C9—C10 | 103.62 (13) | C10—C11—H11B | 111.00 |
| C9—C10—C11 | 103.58 (12) | H11A—C11—H11B | 109.00 |
| C7—C11—C10 | 103.79 (11) | C12—C13—H13 | 120.00 |
| N3—C12—C13 | 118.49 (12) | C14—C13—H13 | 120.00 |
| N3—C12—C17 | 121.86 (12) | C13—C14—H14 | 120.00 |
| C13—C12—C17 | 119.47 (12) | C15—C14—H14 | 120.00 |
| C12—C13—C14 | 120.09 (13) | C14—C15—H15 | 120.00 |
| C13—C14—C15 | 120.46 (14) | C16—C15—H15 | 120.00 |
| C14—C15—C16 | 119.58 (14) | C15—C16—H16 | 120.00 |
| C15—C16—C17 | 120.43 (14) | C17—C16—H16 | 120.00 |
| C12—C17—C16 | 119.90 (13) | C12—C17—H17 | 120.00 |
| C3—C4—H4 | 120.00 | C16—C17—H17 | 120.00 |
| C5—C4—H4 | 120.00 | | |
| | | | |
| C3—S1—C2—N1 | -1.19 (9) | O1—C1—C3—C4 | 3.2 (2) |
| C3—S1—C2—N3 | 179.13 (13) | N1—C1—C3—S1 | 1.81 (13) |
| C2—S1—C3—C1 | -0.35 (10) | N1—C1—C3—C4 | -177.24 (12) |
| C2—S1—C3—C4 | 178.65 (12) | S1—C3—C4—C5 | 0.00 (19) |
| C6—O3—C5—O2 | -2.99 (19) | C1—C3—C4—C5 | 178.90 (12) |
| C6—O3—C5—C4 | 178.11 (11) | C3—C4—C5—O2 | 6.7 (2) |
| C1—N1—N2—C7 | 84.66 (15) | C3—C4—C5—O3 | -174.42 (12) |
| C2—N1—N2—C7 | -103.83 (13) | N2—C7—C8—C9 | -171.37 (15) |
| N2—N1—C1—O1 | -11.7 (2) | C11—C7—C8—C9 | 10.31 (16) |
| N2—N1—C1—C3 | 168.71 (10) | N2—C7—C11—C10 | -164.51 (13) |
| C2—N1—C1—O1 | 176.71 (13) | C8—C7—C11—C10 | 14.00 (15) |
| C2—N1—C1—C3 | -2.90 (15) | C7—C8—C9—C10 | -30.79 (16) |
| N2—N1—C2—S1 | -169.23 (8) | C8—C9—C10—C11 | 39.85 (15) |
| N2—N1—C2—N3 | 10.49 (18) | C9—C10—C11—C7 | -33.10 (14) |
| C1—N1—C2—S1 | 2.68 (14) | N3—C12—C13—C14 | 177.64 (13) |
| C1—N1—C2—N3 | -177.61 (12) | C17—C12—C13—C14 | 2.6 (2) |
| N1—N2—C7—C8 | 1.4 (2) | N3—C12—C17—C16 | -178.17 (13) |
| N1—N2—C7—C11 | 179.58 (11) | C13—C12—C17—C16 | -3.2 (2) |
| C12—N3—C2—S1 | -5.62 (19) | C12—C13—C14—C15 | -0.5 (2) |
| C12—N3—C2—N1 | 174.72 (11) | C13—C14—C15—C16 | -0.9 (2) |
| C2—N3—C12—C13 | 130.97 (14) | C14—C15—C16—C17 | 0.2 (2) |
| C2—N3—C12—C17 | -54.06 (18) | C15—C16—C17—C12 | 1.9 (2) |
| O1—C1—C3—S1 | -177.80 (11) | | |

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-----------------------------------|-------|-------------|-------------|---------------|
| C10—H10B \cdots O1 ⁱ | 0.99 | 2.57 | 3.2889 (17) | 130 |

| | | | | |
|----------------------------|------|------|-------------|-----|
| C11—H11A···O1 ⁱ | 0.99 | 2.58 | 3.2547 (16) | 125 |
|----------------------------|------|------|-------------|-----|

Symmetry code: (i) $-x+1, -y+2, -z+1$.