

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Hydroxy-5-methoxy-*N*,1-dimethyl-2-oxo-*N*-[4-(trifluoromethyl)phenyl]-1,2-dihydroquinoline-3-carboxamide

Emmanuel S. Akinboye,^a Ray J. Butcher,^{b*} Sema Ozturk Yildirim^c and John T. Isaacs^a

^aThe Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, 1650 Orleans Street, Baltimore, MD 21287, USA, ^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington DC 20059, USA, and ^cDepartment of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey

Correspondence e-mail: rbutcher99@yahoo.com


Received 25 January 2014; accepted 10 February 2014

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.132; data-to-parameter ratio = 13.2.

The title compound, $\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_4$, named tasquinimod, is a second-generation oral quinoline-3-carboxamide analogue, which is currently in phase III clinical trials for the treatment of metastatic prostate cancer. The quinoline unit is almost planar (r.m.s. deviation of fitted atoms = 0.0075 Å). The carboxamide side chain, substituted at position 3, is tilted by 88.07 (7)° to the quinoline plane. Both the methyl and carbonyl groups of this carboxamide side chain are in a *syn* conformation. The 4-(trifluoromethyl)phenyl plane is inclined at 50.62 (17)° to the plane of the carboxamide side chain, and at 87.14 (4)° to the plane of the quinoline ring system. The 4-hydroxy H atom acts as a double proton donor in an intramolecular hydrogen bond to the 5-position methoxy O atom and in an intermolecular contact to the 2-oxo group, generating a chain along [010] in the crystal structure.

Related literature

For background to the activity of the second generation quinolone-3-carboxamide analogues roquinimex (also known as linomide, systematic name: 4-hydroxy-*N*,1-dimethyl-2-oxo-*N*-phenyl-1,2-dihydroquinoline-3-carboxamide) and tasquinimod (systematic name: 4-hydroxy-5-methoxy-*N*,1-dimethyl-2-oxo-*N*-[4-(trifluoromethyl)phenyl]-1,2-dihydroquinoline-3-carboxamide), see: Isaacs (2010). For similar structures, see: Dasari & Srikrishnan (2002); Jönsson *et al.* (2004); Jansson *et al.* (2006).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_4$

$M_r = 406.36$

Monoclinic, $P2_1/c$

$a = 10.8643$ (9) Å

$b = 10.6705$ (9) Å

$c = 15.8062$ (13) Å

$\beta = 105.696$ (9)°

$V = 1764.1$ (3) Å³

$Z = 4$

Cu $K\alpha$ radiation

$\mu = 1.10$ mm⁻¹

$T = 298$ K

$0.35 \times 0.24 \times 0.12$ mm

Data collection

Agilent Xcalibur (Ruby, Gemini) diffractometer

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2012)

$T_{\text{min}} = 0.741$, $T_{\text{max}} = 1.000$

6559 measured reflections

3539 independent reflections

3010 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.132$

$S = 1.06$

3539 reflections

269 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.35$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|----------|-------------|-------------|---------------|
| $\text{O2}-\text{H2O}\cdots\text{O4}$ | 0.89 (3) | 1.82 (3) | 2.5538 (16) | 138 (2) |
| $\text{O2}-\text{H2O}\cdots\text{O3}^i$ | 0.89 (3) | 2.28 (3) | 2.8591 (17) | 123 (2) |

Symmetry code: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This study was partially supported by a sponsored research agreement between The Johns Hopkins University School of Medicine (J. Isaacs, PI) and Active Biotech Research AB. RJB acknowledges the NSF-MRI program (grant No. CHE0619278) for funds to purchase the X-ray diffractometer as well as the Howard University Nanoscience Facility for access to liquid nitrogen.

Supporting information for this paper is available from the IUCr electronic archives (Reference: KP2464).

References

Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.

- Dasari, K. B. & Srikrishnan, T. (2002). *J. Chem. Crystallogr.* **32**, 499–504.
Isaacs, J. T. (2010). *Expert Opin. Investig. Drugs*, **19**, 1235–1243.
Jansson, K., Fristedt, T., Olsson, A., Svensson, B. & Jönsson, S. (2006). *J. Org. Chem.* **71**, 1658–1667.
Jönsson, S., Andersson, G., Fex, T., Fristedt, T., Hedlund, G., Jansson, K., Abramo, L., Fritzson, I., Pekarski, O., Runström, A., Sandin, H., Thuvesson, I. & Björk, A. (2004). *J. Med. Chem.* **47**, 2075–2088.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2014). E70, o297–o298 [doi:10.1107/S1600536814003031]

4-Hydroxy-5-methoxy-*N*,1-dimethyl-2-oxo-*N*-[4-(trifluoromethyl)phenyl]-1,2-dihydroquinoline-3-carboxamide

Emmanuel S. Akinboye, Ray J. Butcher, Sema Ozturk Yildirim and John T. Isaacs

S1. Comment

Roquinimex (also known as linomide: 4-hydroxy-*N*, 1-dimethyl-2-oxo-*N* phenyl-1,2-dihydroquinoline-3-carboxamide) is an orally active molecule that is therapeutic against solid malignancies *via* tumor selective anti-angiogenic abilities (Isaacs, 2010). In clinical trials, however, it had dose-limiting unacceptable side effects. From a library of second generation quinolone-3-carboxamide analogues, an analogue termed tasquinimod (*i.e.*, 4-hydroxy-5-methoxy-*N*,1-dimethyl-2-oxo-*N*-[4-(trifluoromethyl) phenyl]-1,2-dihydroquinoline-3-carboxamide) was identified being more than 30 times more potent than an anti-angiogenic agent without the toxic effects of roquinimex (Isaacs, 2010). Presently, tasquinimod is undergoing phase III clinical trials as monotherapy for castration resistant metastatic prostate cancer. Ultimately, tasquinimod will be optimally used in combination with other therapeutic agents. To identify the optimal combinational regime, the identification of the molecular target(s) involved in its mechanism of action is critical. Similar structures have been previously reported (Dasari *et al.*, 2002; Jönsson *et al.*, 2004; Jansson *et al.*, 2006).

As a lead for defining pharmacophoric features of tasquinimod, its three-dimensional structure was determined by X-ray structure analysis. The molecular structure shows that quinoline ring is almost planar (rms deviation of fitted atoms = 0.0075 Å). The carboxamide side chain at position 3, is tilted by 88.07 (7)° to the quinoline plane. Both, the methyl and carbonyl groups of this carboxamide side chain are in a *syn* conformation. The *N*-[4-(trifluoromethyl)phenyl] plane is inclined at 50.62 (17)° to the plane of the carboxamide side chain, and 87.14 (4)° to the plane of the quinoline ring. The 4-hydroxy proton forms intramolecular hydrogen bond with the 5-position methoxy oxygen being planar to the quinoline ring (Fig. 1, Table 1). In the crystal packing molecules are linked by an intermolecular hydrogen bond between 4-hydroxy and 2-oxo groups generating a chain in the direction [010] (Fig. 2, Table 1).

S2. Experimental

The preparation of the titled compound was reported by Jönsson *et al.* (2004).

To obtain crystal of this compound, a saturated solution of a pure sample of the title compound was made in 5 mL of acetonitrile (HPLC grade) at room temperature. The solution was filtered, and 2.5 mL of this saturated solution was transferred into a transparent 2 dram glass vial. Crystallization of this sample was by vapor diffusion in diethyl ether at room temperature over a period of 48 h. Crystal growth was observed within 24 h, but the experiment was left to stand for 48 h. A gentle vacuum filtration of the sample followed by air drying afforded white crystals of the compound for X-ray diffraction studies.

S3. Refinement

H atoms were positioned geometrically and refined using the riding model, with C–H distance of 0.93–0.96 Å, with U_{iso} (H) = 1.20 U_{eq} (C) or 1.50 U_{eq} (C) for methyl H atoms. Hydrogen atoms involved in hydrogen bonding were refined

isotropically.




Figure 1

ORTEP diagram of the title compound showing the atom numbering and the atomic displacement parameters drawn at the 30% probability level. Intramolecular hydrogen bonding is shown by dashed lines.




Figure 2

Crystal packing of the title compound viewed along the *a* axis. Hydrogen bonding is shown by dashed lines.

4-Hydroxy-5-methoxy-*N*,1-dimethyl-2-oxo-*N*-[4-(trifluoromethyl)phenyl]-1,2-dihydroquinoline-3-carboxamide

Crystal data

$C_{20}H_{17}F_3N_2O_4$

$M_r = 406.36$

Monoclinic, $P2_1/c$

$a = 10.8643(9) \text{ \AA}$

$b = 10.6705(9) \text{ \AA}$

$c = 15.8062(13) \text{ \AA}$

$\beta = 105.696(9)^\circ$

$V = 1764.1(3) \text{ \AA}^3$

$Z = 4$

$F(000) = 840$

$D_x = 1.530 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$

Cell parameters from 2882 reflections

$\theta = 2.9\text{--}75.7^\circ$

$\mu = 1.10 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Prism, colourless

$0.35 \times 0.24 \times 0.12 \text{ mm}$

Data collection

Agilent Xcalibur (Ruby, Gemini) diffractometer

Radiation source: Enhance (Cu) X-ray Source

Graphite monochromator

Detector resolution: 10.5081 pixels mm⁻¹

ω scans

Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2012)

$T_{\min} = 0.741$, $T_{\max} = 1.000$

6559 measured reflections

3539 independent reflections

3010 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 75.9^\circ$, $\theta_{\min} = 4.2^\circ$

$h = -8 \rightarrow 13$

$k = -12 \rightarrow 13$

$l = -19 \rightarrow 19$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.132$

$S = 1.06$

3539 reflections

269 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.079P)^2 + 0.3326P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|---------------|--------------|--------------|----------------------------------|
| F1 | -0.01967 (12) | 0.99424 (13) | 0.21797 (7) | 0.0584 (4) |
| F2 | 0.02599 (11) | 0.89869 (12) | 0.34160 (8) | 0.0513 (3) |
| F3 | 0.15877 (10) | 1.02840 (11) | 0.31447 (8) | 0.0480 (3) |
| O1 | 0.42263 (14) | 0.38668 (14) | 0.11242 (8) | 0.0479 (3) |
| O2 | 0.59427 (11) | 0.58973 (11) | 0.24904 (7) | 0.0341 (3) |
| H2O | 0.665 (2) | 0.627 (3) | 0.2800 (17) | 0.056 (7)* |
| O3 | 0.26656 (11) | 0.31456 (11) | 0.25976 (7) | 0.0374 (3) |
| O4 | 0.77617 (11) | 0.61692 (12) | 0.38907 (7) | 0.0404 (3) |
| N1 | 0.29733 (14) | 0.55256 (14) | 0.11795 (8) | 0.0358 (3) |
| N2 | 0.43399 (12) | 0.33345 (13) | 0.38337 (8) | 0.0305 (3) |
| C1 | 0.24402 (16) | 0.64495 (15) | 0.16254 (10) | 0.0339 (3) |
| C2 | 0.11175 (16) | 0.66235 (17) | 0.13832 (11) | 0.0405 (4) |
| H2A | 0.0598 | 0.6100 | 0.0966 | 0.049* |
| C3 | 0.05753 (16) | 0.75701 (17) | 0.17604 (11) | 0.0404 (4) |
| H3A | -0.0306 | 0.7682 | 0.1597 | 0.049* |

| | | | | |
|------|--------------|--------------|--------------|------------|
| C4 | 0.13541 (15) | 0.83560 (15) | 0.23863 (10) | 0.0338 (3) |
| C5 | 0.26668 (15) | 0.81696 (15) | 0.26418 (10) | 0.0334 (3) |
| H5A | 0.3184 | 0.8678 | 0.3072 | 0.040* |
| C6 | 0.32089 (14) | 0.72269 (15) | 0.22568 (10) | 0.0329 (3) |
| H6A | 0.4091 | 0.7115 | 0.2422 | 0.039* |
| C7 | 0.07592 (15) | 0.93888 (17) | 0.27743 (10) | 0.0365 (4) |
| C8 | 0.25048 (19) | 0.54745 (18) | 0.02153 (10) | 0.0442 (4) |
| H8A | 0.3207 | 0.5308 | -0.0029 | 0.066* |
| H8B | 0.2122 | 0.6263 | -0.0002 | 0.066* |
| H8C | 0.1880 | 0.4820 | 0.0048 | 0.066* |
| C9 | 0.38280 (16) | 0.46314 (16) | 0.15651 (10) | 0.0350 (3) |
| C10 | 0.43151 (15) | 0.45503 (15) | 0.25502 (9) | 0.0313 (3) |
| C11 | 0.54421 (15) | 0.50991 (15) | 0.29655 (9) | 0.0303 (3) |
| C12 | 0.60720 (14) | 0.47863 (15) | 0.38701 (9) | 0.0291 (3) |
| C13 | 0.72526 (15) | 0.53184 (16) | 0.43456 (10) | 0.0322 (3) |
| C14 | 0.78219 (15) | 0.49785 (16) | 0.52027 (10) | 0.0346 (3) |
| H14A | 0.8594 | 0.5336 | 0.5510 | 0.042* |
| C15 | 0.72218 (16) | 0.40882 (16) | 0.56024 (10) | 0.0362 (3) |
| H15A | 0.7602 | 0.3862 | 0.6183 | 0.043* |
| C16 | 0.60834 (15) | 0.35357 (15) | 0.51625 (10) | 0.0331 (3) |
| H16A | 0.5708 | 0.2938 | 0.5441 | 0.040* |
| C17 | 0.54900 (14) | 0.38831 (14) | 0.42858 (9) | 0.0297 (3) |
| C18 | 0.36910 (14) | 0.36439 (15) | 0.29723 (10) | 0.0313 (3) |
| C19 | 0.37877 (16) | 0.23443 (16) | 0.42576 (10) | 0.0361 (3) |
| H19A | 0.3656 | 0.2656 | 0.4796 | 0.054* |
| H19B | 0.4361 | 0.1642 | 0.4382 | 0.054* |
| H19C | 0.2984 | 0.2085 | 0.3873 | 0.054* |
| C20 | 0.88302 (17) | 0.6906 (2) | 0.43465 (12) | 0.0461 (4) |
| H20A | 0.9008 | 0.7526 | 0.3956 | 0.069* |
| H20B | 0.9563 | 0.6375 | 0.4554 | 0.069* |
| H20C | 0.8639 | 0.7314 | 0.4837 | 0.069* |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|------------|------------|------------|-------------|-------------|-------------|
| F1 | 0.0625 (7) | 0.0633 (8) | 0.0377 (5) | 0.0276 (6) | -0.0067 (5) | -0.0050 (5) |
| F2 | 0.0575 (6) | 0.0533 (7) | 0.0479 (6) | -0.0070 (5) | 0.0224 (5) | -0.0005 (5) |
| F3 | 0.0453 (5) | 0.0401 (6) | 0.0594 (6) | -0.0072 (5) | 0.0155 (5) | -0.0122 (5) |
| O1 | 0.0600 (8) | 0.0492 (8) | 0.0284 (5) | 0.0130 (6) | 0.0014 (5) | -0.0064 (5) |
| O2 | 0.0374 (6) | 0.0372 (6) | 0.0250 (5) | -0.0019 (5) | 0.0036 (4) | 0.0034 (4) |
| O3 | 0.0362 (6) | 0.0355 (6) | 0.0356 (6) | -0.0014 (5) | 0.0012 (4) | -0.0019 (5) |
| O4 | 0.0403 (6) | 0.0441 (7) | 0.0314 (5) | -0.0102 (5) | 0.0005 (5) | 0.0059 (5) |
| N1 | 0.0438 (7) | 0.0350 (7) | 0.0228 (6) | -0.0013 (6) | -0.0009 (5) | 0.0007 (5) |
| N2 | 0.0337 (6) | 0.0302 (6) | 0.0261 (6) | 0.0003 (5) | 0.0056 (5) | 0.0003 (5) |
| C1 | 0.0404 (8) | 0.0312 (7) | 0.0255 (7) | -0.0025 (6) | 0.0010 (6) | 0.0050 (6) |
| C2 | 0.0399 (8) | 0.0364 (8) | 0.0361 (8) | -0.0045 (7) | -0.0056 (6) | -0.0006 (6) |
| C3 | 0.0342 (7) | 0.0389 (9) | 0.0403 (8) | -0.0032 (7) | -0.0034 (6) | 0.0011 (7) |
| C4 | 0.0369 (8) | 0.0323 (8) | 0.0285 (7) | -0.0033 (6) | 0.0028 (6) | 0.0038 (6) |

| | | | | | | |
|-----|-------------|-------------|------------|-------------|-------------|-------------|
| C5 | 0.0368 (7) | 0.0328 (8) | 0.0261 (7) | -0.0057 (6) | 0.0011 (6) | 0.0029 (6) |
| C6 | 0.0330 (7) | 0.0339 (8) | 0.0273 (7) | -0.0029 (6) | 0.0005 (5) | 0.0041 (6) |
| C7 | 0.0340 (7) | 0.0415 (9) | 0.0298 (7) | -0.0032 (6) | 0.0013 (6) | 0.0030 (6) |
| C8 | 0.0584 (10) | 0.0421 (9) | 0.0245 (7) | -0.0007 (8) | -0.0020 (7) | 0.0010 (6) |
| C9 | 0.0402 (8) | 0.0346 (8) | 0.0260 (7) | -0.0023 (6) | 0.0017 (6) | -0.0022 (6) |
| C10 | 0.0363 (7) | 0.0301 (7) | 0.0240 (7) | 0.0034 (6) | 0.0020 (5) | -0.0003 (5) |
| C11 | 0.0353 (7) | 0.0303 (7) | 0.0237 (6) | 0.0037 (6) | 0.0054 (5) | 0.0008 (5) |
| C12 | 0.0325 (7) | 0.0297 (7) | 0.0226 (6) | 0.0040 (6) | 0.0030 (5) | -0.0015 (5) |
| C13 | 0.0341 (7) | 0.0320 (8) | 0.0284 (7) | 0.0019 (6) | 0.0050 (6) | 0.0004 (6) |
| C14 | 0.0326 (7) | 0.0373 (8) | 0.0286 (7) | 0.0001 (6) | -0.0010 (6) | -0.0008 (6) |
| C15 | 0.0421 (8) | 0.0380 (8) | 0.0243 (7) | 0.0035 (7) | 0.0018 (6) | 0.0026 (6) |
| C16 | 0.0397 (8) | 0.0322 (7) | 0.0260 (7) | 0.0027 (6) | 0.0068 (6) | 0.0021 (6) |
| C17 | 0.0331 (7) | 0.0302 (7) | 0.0243 (6) | 0.0035 (6) | 0.0053 (5) | -0.0022 (5) |
| C18 | 0.0337 (7) | 0.0299 (7) | 0.0274 (7) | 0.0030 (6) | 0.0034 (6) | -0.0034 (6) |
| C19 | 0.0399 (8) | 0.0349 (8) | 0.0324 (7) | -0.0041 (6) | 0.0080 (6) | 0.0011 (6) |
| C20 | 0.0399 (8) | 0.0497 (11) | 0.0430 (9) | -0.0116 (8) | 0.0013 (7) | 0.0059 (8) |

Geometric parameters (Å, °)

| | | | |
|------------|-------------|-------------|-------------|
| F1—C7 | 1.3351 (19) | C5—H5A | 0.9300 |
| F2—C7 | 1.343 (2) | C6—H6A | 0.9300 |
| F3—C7 | 1.334 (2) | C8—H8A | 0.9600 |
| O1—C9 | 1.225 (2) | C8—H8B | 0.9600 |
| O2—C11 | 1.3440 (19) | C8—H8C | 0.9600 |
| O2—H2O | 0.89 (3) | C9—C10 | 1.505 (2) |
| O3—C18 | 1.232 (2) | C10—C11 | 1.356 (2) |
| O4—C13 | 1.365 (2) | C10—C18 | 1.444 (2) |
| O4—C20 | 1.426 (2) | C11—C12 | 1.447 (2) |
| N1—C9 | 1.356 (2) | C12—C17 | 1.409 (2) |
| N1—C1 | 1.423 (2) | C12—C13 | 1.418 (2) |
| N1—C8 | 1.4713 (19) | C13—C14 | 1.377 (2) |
| N2—C17 | 1.390 (2) | C14—C15 | 1.396 (2) |
| N2—C18 | 1.3930 (19) | C14—H14A | 0.9300 |
| N2—C19 | 1.464 (2) | C15—C16 | 1.377 (2) |
| C1—C6 | 1.390 (2) | C15—H15A | 0.9300 |
| C1—C2 | 1.396 (2) | C16—C17 | 1.411 (2) |
| C2—C3 | 1.383 (3) | C16—H16A | 0.9300 |
| C2—H2A | 0.9300 | C19—H19A | 0.9600 |
| C3—C4 | 1.395 (2) | C19—H19B | 0.9600 |
| C3—H3A | 0.9300 | C19—H19C | 0.9600 |
| C4—C5 | 1.387 (2) | C20—H20A | 0.9600 |
| C4—C7 | 1.491 (2) | C20—H20B | 0.9600 |
| C5—C6 | 1.387 (2) | C20—H20C | 0.9600 |
| C11—O2—H2O | 113.5 (17) | N1—C9—C10 | 120.79 (14) |
| C13—O4—C20 | 119.44 (13) | C11—C10—C18 | 122.75 (13) |
| C9—N1—C1 | 125.87 (12) | C11—C10—C9 | 119.65 (14) |
| C9—N1—C8 | 116.27 (14) | C18—C10—C9 | 116.03 (13) |

| | | | |
|-------------|--------------|-----------------|--------------|
| C1—N1—C8 | 117.75 (14) | O2—C11—C10 | 116.88 (13) |
| C17—N2—C18 | 123.35 (13) | O2—C11—C12 | 122.92 (14) |
| C17—N2—C19 | 119.43 (12) | C10—C11—C12 | 120.16 (14) |
| C18—N2—C19 | 117.18 (13) | C17—C12—C13 | 118.84 (13) |
| C6—C1—C2 | 119.45 (16) | C17—C12—C11 | 117.81 (14) |
| C6—C1—N1 | 121.55 (15) | C13—C12—C11 | 123.33 (15) |
| C2—C1—N1 | 118.89 (14) | O4—C13—C14 | 123.74 (14) |
| C3—C2—C1 | 120.39 (15) | O4—C13—C12 | 115.12 (13) |
| C3—C2—H2A | 119.8 | C14—C13—C12 | 121.14 (15) |
| C1—C2—H2A | 119.8 | C13—C14—C15 | 118.94 (14) |
| C2—C3—C4 | 119.84 (15) | C13—C14—H14A | 120.5 |
| C2—C3—H3A | 120.1 | C15—C14—H14A | 120.5 |
| C4—C3—H3A | 120.1 | C16—C15—C14 | 121.90 (14) |
| C5—C4—C3 | 119.93 (16) | C16—C15—H15A | 119.1 |
| C5—C4—C7 | 120.80 (14) | C14—C15—H15A | 119.1 |
| C3—C4—C7 | 119.27 (15) | C15—C16—C17 | 119.54 (15) |
| C6—C5—C4 | 120.09 (14) | C15—C16—H16A | 120.2 |
| C6—C5—H5A | 120.0 | C17—C16—H16A | 120.2 |
| C4—C5—H5A | 120.0 | N2—C17—C12 | 120.19 (13) |
| C5—C6—C1 | 120.28 (15) | N2—C17—C16 | 120.19 (14) |
| C5—C6—H6A | 119.9 | C12—C17—C16 | 119.63 (14) |
| C1—C6—H6A | 119.9 | O3—C18—N2 | 121.36 (15) |
| F3—C7—F1 | 107.22 (15) | O3—C18—C10 | 122.93 (14) |
| F3—C7—F2 | 105.04 (13) | N2—C18—C10 | 115.69 (13) |
| F1—C7—F2 | 106.06 (14) | N2—C19—H19A | 109.5 |
| F3—C7—C4 | 113.18 (14) | N2—C19—H19B | 109.5 |
| F1—C7—C4 | 112.25 (13) | H19A—C19—H19B | 109.5 |
| F2—C7—C4 | 112.52 (15) | N2—C19—H19C | 109.5 |
| N1—C8—H8A | 109.5 | H19A—C19—H19C | 109.5 |
| N1—C8—H8B | 109.5 | H19B—C19—H19C | 109.5 |
| H8A—C8—H8B | 109.5 | O4—C20—H20A | 109.5 |
| N1—C8—H8C | 109.5 | O4—C20—H20B | 109.5 |
| H8A—C8—H8C | 109.5 | H20A—C20—H20B | 109.5 |
| H8B—C8—H8C | 109.5 | O4—C20—H20C | 109.5 |
| O1—C9—N1 | 121.14 (14) | H20A—C20—H20C | 109.5 |
| O1—C9—C10 | 118.07 (15) | H20B—C20—H20C | 109.5 |
| | | | |
| C9—N1—C1—C6 | 53.1 (2) | O2—C11—C12—C17 | -176.45 (13) |
| C8—N1—C1—C6 | -131.00 (17) | C10—C11—C12—C17 | 1.4 (2) |
| C9—N1—C1—C2 | -130.73 (18) | O2—C11—C12—C13 | 1.9 (2) |
| C8—N1—C1—C2 | 45.2 (2) | C10—C11—C12—C13 | 179.78 (14) |
| C6—C1—C2—C3 | 0.7 (3) | C20—O4—C13—C14 | -10.4 (2) |
| N1—C1—C2—C3 | -175.56 (15) | C20—O4—C13—C12 | 169.82 (15) |
| C1—C2—C3—C4 | 0.0 (3) | C17—C12—C13—O4 | 178.95 (13) |
| C2—C3—C4—C5 | -1.3 (3) | C11—C12—C13—O4 | 0.6 (2) |
| C2—C3—C4—C7 | 178.77 (15) | C17—C12—C13—C14 | -0.9 (2) |
| C3—C4—C5—C6 | 1.9 (2) | C11—C12—C13—C14 | -179.19 (15) |
| C7—C4—C5—C6 | -178.19 (14) | O4—C13—C14—C15 | -179.40 (15) |

| | | | |
|-----------------|--------------|-----------------|--------------|
| C4—C5—C6—C1 | -1.2 (2) | C12—C13—C14—C15 | 0.4 (2) |
| C2—C1—C6—C5 | -0.1 (2) | C13—C14—C15—C16 | 0.4 (3) |
| N1—C1—C6—C5 | 176.04 (14) | C14—C15—C16—C17 | -0.8 (3) |
| C5—C4—C7—F3 | 19.4 (2) | C18—N2—C17—C12 | -1.2 (2) |
| C3—C4—C7—F3 | -160.76 (15) | C19—N2—C17—C12 | 176.37 (13) |
| C5—C4—C7—F1 | 140.90 (16) | C18—N2—C17—C16 | 179.06 (14) |
| C3—C4—C7—F1 | -39.2 (2) | C19—N2—C17—C16 | -3.4 (2) |
| C5—C4—C7—F2 | -99.54 (18) | C13—C12—C17—N2 | -179.28 (13) |
| C3—C4—C7—F2 | 80.34 (18) | C11—C12—C17—N2 | -0.8 (2) |
| C1—N1—C9—O1 | 179.39 (16) | C13—C12—C17—C16 | 0.5 (2) |
| C8—N1—C9—O1 | 3.4 (2) | C11—C12—C17—C16 | 178.94 (14) |
| C1—N1—C9—C10 | -0.6 (2) | C15—C16—C17—N2 | -179.92 (14) |
| C8—N1—C9—C10 | -176.58 (15) | C15—C16—C17—C12 | 0.3 (2) |
| O1—C9—C10—C11 | 83.9 (2) | C17—N2—C18—O3 | -179.18 (14) |
| N1—C9—C10—C11 | -96.11 (19) | C19—N2—C18—O3 | 3.2 (2) |
| O1—C9—C10—C18 | -82.2 (2) | C17—N2—C18—C10 | 2.4 (2) |
| N1—C9—C10—C18 | 97.77 (18) | C19—N2—C18—C10 | -175.13 (13) |
| C18—C10—C11—O2 | 177.94 (13) | C11—C10—C18—O3 | 179.83 (15) |
| C9—C10—C11—O2 | 12.8 (2) | C9—C10—C18—O3 | -14.5 (2) |
| C18—C10—C11—C12 | -0.1 (2) | C11—C10—C18—N2 | -1.8 (2) |
| C9—C10—C11—C12 | -165.21 (14) | C9—C10—C18—N2 | 163.83 (13) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H... <i>A</i> | <i>D</i> —H | H... <i>A</i> | <i>D</i> ... <i>A</i> | <i>D</i> —H... <i>A</i> |
|--------------------------|-------------|---------------|-----------------------|-------------------------|
| O2—H2O...O4 | 0.89 (3) | 1.82 (3) | 2.5538 (16) | 138 (2) |
| O2—H2O...O3 ⁱ | 0.89 (3) | 2.28 (3) | 2.8591 (17) | 123 (2) |

Symmetry code: (i) $-x+1, y+1/2, -z+1/2$.