

tert-Butyl N-[2-(*N*-isobutyl-4-methoxybenzenesulfonamido)ethyl]carbamate

Xiao-Guang Bai and Ju-Xian Wang*

Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, People's Republic of China
Correspondence e-mail: imbjxwang@gmail.com


Received 15 April 2014; accepted 23 April 2014

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.034; wR factor = 0.089; data-to-parameter ratio = 14.9.

The title compound, $C_{18}H_{30}N_2O_5S$, was synthesized by the reaction of *tert*-butyl 2-(isobutylamino)ethylcarbamate with *p*-methoxyphenylsulfonyl chloride. In the molecule, two intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds are observed. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds involving the imino group N atom and the ester group O atom into chains running parallel to the b axis. The chains are further connected by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming layers parallel to the bc plane.

Related literature

For potential HIV-1 protease inhibitors, see: Surleraux *et al.* (2005); Ghosh *et al.* (2006, 2011); Guo *et al.* (2010). For the structure of the methoxy analogue, see: Chatziefthimiou *et al.* (2006)



Experimental

Crystal data

$C_{18}H_{30}N_2O_5S$
 $M_r = 386.50$
Monoclinic, $P2_1/c$

$\beta = 92.497(3)\text{ }^\circ$
 $V = 2053.46(9)\text{ \AA}^3$
 $Z = 4$
Cu $K\alpha$ radiation

$\mu = 1.65\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.28 \times 0.16 \times 0.14\text{ mm}$

Data collection

Agilent Xcalibur (Atlas, Gemini ultra) diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2013)
 $T_{\min} = 0.776$, $T_{\max} = 1.000$

11865 measured reflections
3658 independent reflections
3122 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.089$
 $S = 1.05$
3658 reflections
245 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.17\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C8—H8A \cdots O2	0.97	2.43	2.9106 (19)	110
C13—H13A \cdots O3	0.97	2.48	3.107 (2)	122
C3—H3 \cdots O4 ⁱ	0.93	2.59	3.402 (2)	147
N2—H2A \cdots O4 ⁱⁱ	0.82 (2)	2.38 (2)	3.190 (2)	171 (2)

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $x, y + 1, z$.

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC*.

We are grateful for the financial support from the National Natural Science Foundation of China (No. 81302644).

Supporting information for this paper is available from the IUCr electronic archives (Reference: RZ5120).

References

- Agilent (2013). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
- Chatziefthimiou, S. D., Lazarou, Y. G., Hadjoudis, E., Dziembowska, T. & Mavridis, I. M. (2006). *J. Phys. Chem. B*, **110**, 23701–23709.
- Ghosh, A. K., Chapsal, B. D., Parham, G. L., Steffey, M., Agniswamy, J., Wang, Y.-F., Amano, M., Weber, I. T. & Mitsuya, H. (2011). *J. Med. Chem.* **54**, 5890–5901.
- Ghosh, A. K., Sridhar, P. R., Leshchenko, S., Hussain, A. K., Li, J., Kovalevsky, A. Y., Walters, D. E., Wedekind, J. E., Grum-Tokars, V., Das, D., Koh, Y., Maeda, K., Gatanaga, H., Weber, I. T. & Mitsuya, H. (2006). *J. Med. Chem.* **49**, 5252–5261.
- Guo, X., Bai, X. G., Li, Y. L. & Wang, Y. C. (2010). *Acta Cryst. E* **66**, o1108.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Surleraux, D. L. N. G., Tahri, A., Verschueren, W. G., Pille, G. M. E., de Kock, H. A., Jonckers, T. H. M., Peeters, A., De Meyer, S., Azijn, H., Pauwels, R., de Bethune, M.-P., King, N. M., Prabu-Jeyabalan, M., Schiffer, C. A. & Wigerinck, P. B. T. P. (2005). *J. Med. Chem.* **48**, 1813–1822.

supporting information

Acta Cryst. (2014). E70, o674 [doi:10.1107/S1600536814009143]

tert-Butyl N-[2-(*N*-isobutyl-4-methoxybenzenesulfonamido)ethyl]carbamate

Xiao-Guang Bai and Ju-Xian Wang

S1. Comment

As a part of our ongoing project aimed at the development of potential HIV-1 protease inhibitors (Surleraux *et al.*, 2005; Ghosh *et al.*, 2006, 2011; Guo *et al.*, 2010), we have synthesized the title compound and report its crystal structure herein.


The molecular structure of the title compound is illustrated in Fig. 1. Bond distances and angles are similar to those found in the methoxy analogue (Chatziefthimiou *et al.*, 2006). The molecular conformation is stabilized by two intramolecular C—H···O hydrogen bonds (Table 1). In the crystal, the molecules are linked into chains by intermolecular N—H···O hydrogen bonds (Table 1) parallel to the *b* axis (Fig. 2), which are further connected to form layers parallel to the *bc* plane by C—H···O hydrogen bonds (Table 1).

S2. Experimental


To a solution of *tert*-butyl 2-(isobutylamino)ethylcarbamate (1.13 g, 5.2 mmol) and "N,N-diisopropylethylamine (1.34 g, 10.4 mmol) in dichloromethane (10 ml) was added dropwise a solution of *p*-methoxyphenylsulfonyl chloride (1.18 g, 5.7 mmol) in dichloromethane (3 ml) over a period of 10 min at room temperature. The reaction mixture was stirred for 5 h at the same temperature and concentrated under reduced pressure. *tert*-Butyl 2-(*N*-isobutyl-4-methoxyphenylsulfonamido)-ethylcarbamate was obtained as a white solid by flash chromatography (40 g silica gel, petroleum ether/AcOEt, 1:10 v/v). The yield is 42%. Colourless block crystals suitable for X-ray diffraction were obtained in 3 day by slow evaporation of a petroleum ether/AcOEt (4:1 v/v) solution.

S3. Refinement

All H atoms could be detected in a difference Fourier map. The H atom bonded to N2 was refined freely, all other Hatoms were placed in calculated positions and refined using a riding motion approximation, with C—H = 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

**Figure 1**

The molecular structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

Partial packing diagram of the title compound showing the formation of a molecular chain through N—H···O hydrogen bonds. Hydrogen atoms not involved in hydrogen bonding (dashed lines) are omitted.

tert*-Butyl N-[2-(*N*-isobutyl-4-methoxybenzenesulfonamido)ethyl]carbamateCrystal data*

C₁₈H₃₀N₂O₅S
*M*_r = 386.50
 Monoclinic, *P*2₁/*c*
 Hall symbol: -P 2ybc
a = 19.2484 (5) Å
b = 5.29088 (12) Å
c = 20.1825 (6) Å
 β = 92.497 (3) $^\circ$
V = 2053.46 (9) Å³
Z = 4

F(000) = 832
*D*_x = 1.250 Mg m⁻³
 Cu *K* α radiation, λ = 1.54184 Å
 Cell parameters from 5271 reflections
 θ = 4.4–67.1 $^\circ$
 μ = 1.65 mm⁻¹
T = 293 K
 Block, colorless
 0.28 × 0.16 × 0.14 mm

Data collection

Agilent Xcalibur (Atlas, Gemini ultra)
 diffractometer
 Radiation source: Enhance Ultra (Cu) X-ray
 Source
 Mirror monochromator
 Detector resolution: 10.4713 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan
 (*CrysAlis PRO*; Agilent, 2013)

*T*_{min} = 0.776, *T*_{max} = 1.000
 11865 measured reflections
 3658 independent reflections
 3122 reflections with *I* > 2 σ (*I*)
 R _{int} = 0.031
 θ _{max} = 67.2 $^\circ$, θ _{min} = 4.4 $^\circ$
 h = -21 → 22
 k = -4 → 6
 l = -22 → 24

Refinement

Refinement on *F*²
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)]$ = 0.034
 $wR(F^2)$ = 0.089
 S = 1.05
 3658 reflections
 245 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0429P)^2 + 0.3785P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.17 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.30 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of *F*² against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*², conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*². The threshold expression of *F*² > $\sigma(F^2)$ is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*² are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */* <i>U</i> _{eq}
C1	0.85965 (7)	0.6201 (3)	0.54057 (7)	0.0368 (3)
C2	0.82937 (9)	0.6078 (3)	0.60169 (8)	0.0474 (4)
H2	0.7921	0.7121	0.6107	0.057*

C3	0.85494 (9)	0.4403 (4)	0.64878 (8)	0.0527 (4)
H3	0.8347	0.4314	0.6897	0.063*
C4	0.91066 (8)	0.2845 (3)	0.63579 (8)	0.0440 (4)
C5	0.94134 (8)	0.2961 (3)	0.57509 (8)	0.0435 (4)
H5	0.9788	0.1921	0.5662	0.052*
C6	0.91528 (8)	0.4657 (3)	0.52769 (7)	0.0403 (3)
H6	0.9356	0.4754	0.4868	0.048*
C7	0.98789 (12)	-0.0418 (4)	0.67445 (11)	0.0687 (6)
H7A	0.9749	-0.1530	0.6384	0.103*
H7B	0.9983	-0.1396	0.7137	0.103*
H7C	1.0282	0.0537	0.6634	0.103*
C8	0.70575 (8)	0.5843 (3)	0.45894 (8)	0.0420 (3)
H8A	0.6984	0.6863	0.4980	0.050*
H8B	0.7095	0.4093	0.4730	0.050*
C9	0.64333 (8)	0.6121 (3)	0.41093 (8)	0.0447 (4)
H9	0.6499	0.5001	0.3730	0.054*
C10	0.57790 (9)	0.5294 (4)	0.44490 (10)	0.0602 (5)
H10A	0.5853	0.3658	0.4645	0.090*
H10B	0.5397	0.5213	0.4128	0.090*
H10C	0.5675	0.6491	0.4788	0.090*
C11	0.63562 (11)	0.8797 (4)	0.38541 (12)	0.0743 (6)
H11A	0.6288	0.9921	0.4219	0.111*
H11B	0.5963	0.8894	0.3546	0.111*
H11C	0.6769	0.9276	0.3635	0.111*
C12	0.80312 (8)	0.4772 (3)	0.38535 (8)	0.0426 (3)
H12A	0.7712	0.3368	0.3783	0.051*
H12B	0.8454	0.4114	0.4069	0.051*
C13	0.82046 (9)	0.5869 (4)	0.31821 (8)	0.0517 (4)
H13A	0.8475	0.7399	0.3254	0.062*
H13B	0.8493	0.4669	0.2956	0.062*
C14	0.72163 (8)	0.4609 (3)	0.24696 (7)	0.0410 (3)
C15	0.62067 (9)	0.3966 (3)	0.17124 (8)	0.0459 (4)
C16	0.57907 (11)	0.2465 (4)	0.21953 (10)	0.0642 (5)
H16A	0.6083	0.1208	0.2407	0.096*
H16B	0.5408	0.1655	0.1961	0.096*
H16C	0.5617	0.3581	0.2525	0.096*
C17	0.65827 (11)	0.2296 (4)	0.12362 (9)	0.0646 (5)
H17A	0.6895	0.3305	0.0989	0.097*
H17B	0.6250	0.1496	0.0936	0.097*
H17C	0.6842	0.1028	0.1481	0.097*
C18	0.57448 (13)	0.5846 (4)	0.13381 (13)	0.0827 (7)
H18A	0.5534	0.6958	0.1647	0.124*
H18B	0.5388	0.4956	0.1086	0.124*
H18C	0.6019	0.6816	0.1044	0.124*
N1	0.77172 (6)	0.6617 (2)	0.42986 (6)	0.0371 (3)
N2	0.76011 (8)	0.6456 (3)	0.27567 (7)	0.0474 (3)
O1	0.93204 (7)	0.1260 (3)	0.68599 (6)	0.0611 (3)
O2	0.78598 (6)	1.0150 (2)	0.50924 (6)	0.0505 (3)

O3	0.88216 (6)	0.8952 (2)	0.43790 (5)	0.0462 (3)
O4	0.73211 (6)	0.2357 (2)	0.25588 (6)	0.0495 (3)
O5	0.67038 (6)	0.5608 (2)	0.20784 (6)	0.0511 (3)
S1	0.825787 (18)	0.82405 (6)	0.478139 (18)	0.03723 (12)
H2A	0.7503 (11)	0.793 (4)	0.2665 (10)	0.060 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0318 (7)	0.0407 (7)	0.0375 (7)	-0.0033 (6)	-0.0025 (6)	-0.0012 (6)
C2	0.0405 (8)	0.0577 (10)	0.0443 (8)	0.0031 (7)	0.0057 (7)	-0.0026 (7)
C3	0.0501 (10)	0.0697 (11)	0.0390 (8)	-0.0005 (8)	0.0101 (7)	0.0035 (8)
C4	0.0441 (8)	0.0474 (8)	0.0400 (8)	-0.0088 (7)	-0.0052 (6)	0.0062 (7)
C5	0.0372 (8)	0.0486 (9)	0.0442 (8)	0.0021 (7)	-0.0017 (6)	0.0011 (7)
C6	0.0345 (7)	0.0497 (8)	0.0366 (7)	0.0005 (6)	0.0007 (6)	0.0017 (6)
C7	0.0723 (13)	0.0631 (12)	0.0693 (12)	0.0052 (10)	-0.0139 (10)	0.0211 (10)
C8	0.0367 (8)	0.0466 (8)	0.0428 (8)	-0.0027 (7)	0.0007 (6)	0.0046 (7)
C9	0.0363 (8)	0.0500 (9)	0.0476 (8)	0.0010 (7)	-0.0013 (6)	0.0004 (7)
C10	0.0389 (9)	0.0742 (12)	0.0675 (11)	-0.0056 (9)	0.0011 (8)	0.0022 (10)
C11	0.0536 (11)	0.0713 (13)	0.0980 (16)	0.0155 (10)	0.0030 (11)	0.0339 (12)
C12	0.0400 (8)	0.0428 (8)	0.0444 (8)	0.0040 (7)	-0.0050 (6)	-0.0050 (7)
C13	0.0422 (9)	0.0688 (11)	0.0442 (8)	-0.0069 (8)	0.0024 (7)	-0.0105 (8)
C14	0.0461 (8)	0.0429 (8)	0.0342 (7)	-0.0037 (7)	0.0034 (6)	-0.0027 (6)
C15	0.0509 (9)	0.0383 (8)	0.0475 (8)	-0.0074 (7)	-0.0082 (7)	-0.0010 (7)
C16	0.0576 (11)	0.0712 (12)	0.0645 (11)	-0.0108 (10)	0.0085 (9)	0.0014 (10)
C17	0.0709 (13)	0.0721 (12)	0.0509 (10)	-0.0149 (10)	0.0038 (9)	-0.0163 (9)
C18	0.0884 (16)	0.0549 (11)	0.1003 (17)	-0.0064 (11)	-0.0478 (14)	0.0101 (11)
N1	0.0324 (6)	0.0399 (6)	0.0386 (6)	-0.0009 (5)	-0.0021 (5)	-0.0010 (5)
N2	0.0557 (8)	0.0455 (8)	0.0405 (7)	-0.0096 (7)	-0.0031 (6)	-0.0014 (6)
O1	0.0663 (8)	0.0678 (8)	0.0489 (7)	0.0010 (7)	-0.0025 (6)	0.0188 (6)
O2	0.0520 (7)	0.0400 (6)	0.0593 (7)	0.0084 (5)	-0.0007 (5)	-0.0097 (5)
O3	0.0422 (6)	0.0462 (6)	0.0502 (6)	-0.0078 (5)	0.0016 (5)	0.0059 (5)
O4	0.0555 (7)	0.0411 (6)	0.0512 (6)	0.0035 (5)	-0.0035 (5)	-0.0034 (5)
O5	0.0606 (7)	0.0367 (6)	0.0543 (6)	-0.0050 (5)	-0.0165 (5)	-0.0016 (5)
S1	0.03497 (19)	0.03481 (19)	0.0416 (2)	-0.00052 (14)	-0.00156 (14)	-0.00101 (14)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.381 (2)	C11—H11C	0.9600
C1—C2	1.389 (2)	C12—N1	1.474 (2)
C1—S1	1.7621 (15)	C12—C13	1.524 (2)
C2—C3	1.375 (2)	C12—H12A	0.9700
C2—H2	0.9300	C12—H12B	0.9700
C3—C4	1.387 (3)	C13—N2	1.448 (2)
C3—H3	0.9300	C13—H13A	0.9700
C4—O1	1.365 (2)	C13—H13B	0.9700
C4—C5	1.384 (2)	C14—O4	1.2205 (19)
C5—C6	1.389 (2)	C14—N2	1.342 (2)

C5—H5	0.9300	C14—O5	1.345 (2)
C6—H6	0.9300	C15—O5	1.4677 (19)
C7—O1	1.421 (3)	C15—C17	1.513 (3)
C7—H7A	0.9600	C15—C16	1.513 (3)
C7—H7B	0.9600	C15—C18	1.514 (3)
C7—H7C	0.9600	C16—H16A	0.9600
C8—N1	1.4797 (19)	C16—H16B	0.9600
C8—C9	1.518 (2)	C16—H16C	0.9600
C8—H8A	0.9700	C17—H17A	0.9600
C8—H8B	0.9700	C17—H17B	0.9600
C9—C11	1.512 (3)	C17—H17C	0.9600
C9—C10	1.524 (2)	C18—H18A	0.9600
C9—H9	0.9800	C18—H18B	0.9600
C10—H10A	0.9600	C18—H18C	0.9600
C10—H10B	0.9600	N1—S1	1.6378 (12)
C10—H10C	0.9600	N2—H2A	0.82 (2)
C11—H11A	0.9600	O2—S1	1.4294 (12)
C11—H11B	0.9600	O3—S1	1.4338 (12)
C6—C1—C2	119.91 (14)	N1—C12—H12B	108.8
C6—C1—S1	119.56 (11)	C13—C12—H12B	108.8
C2—C1—S1	120.50 (12)	H12A—C12—H12B	107.7
C3—C2—C1	119.52 (15)	N2—C13—C12	114.06 (14)
C3—C2—H2	120.2	N2—C13—H13A	108.7
C1—C2—H2	120.2	C12—C13—H13A	108.7
C2—C3—C4	120.60 (15)	N2—C13—H13B	108.7
C2—C3—H3	119.7	C12—C13—H13B	108.7
C4—C3—H3	119.7	H13A—C13—H13B	107.6
O1—C4—C5	123.89 (16)	O4—C14—N2	124.23 (15)
O1—C4—C3	115.82 (15)	O4—C14—O5	125.63 (14)
C5—C4—C3	120.29 (15)	N2—C14—O5	110.14 (14)
C4—C5—C6	118.87 (15)	O5—C15—C17	110.24 (14)
C4—C5—H5	120.6	O5—C15—C16	109.76 (14)
C6—C5—H5	120.6	C17—C15—C16	112.57 (16)
C1—C6—C5	120.81 (14)	O5—C15—C18	102.60 (13)
C1—C6—H6	119.6	C17—C15—C18	110.70 (17)
C5—C6—H6	119.6	C16—C15—C18	110.52 (18)
O1—C7—H7A	109.5	C15—C16—H16A	109.5
O1—C7—H7B	109.5	C15—C16—H16B	109.5
H7A—C7—H7B	109.5	H16A—C16—H16B	109.5
O1—C7—H7C	109.5	C15—C16—H16C	109.5
H7A—C7—H7C	109.5	H16A—C16—H16C	109.5
H7B—C7—H7C	109.5	H16B—C16—H16C	109.5
N1—C8—C9	112.90 (12)	C15—C17—H17A	109.5
N1—C8—H8A	109.0	C15—C17—H17B	109.5
C9—C8—H8A	109.0	H17A—C17—H17B	109.5
N1—C8—H8B	109.0	C15—C17—H17C	109.5
C9—C8—H8B	109.0	H17A—C17—H17C	109.5

H8A—C8—H8B	107.8	H17B—C17—H17C	109.5
C11—C9—C8	111.82 (15)	C15—C18—H18A	109.5
C11—C9—C10	110.57 (16)	C15—C18—H18B	109.5
C8—C9—C10	109.38 (13)	H18A—C18—H18B	109.5
C11—C9—H9	108.3	C15—C18—H18C	109.5
C8—C9—H9	108.3	H18A—C18—H18C	109.5
C10—C9—H9	108.3	H18B—C18—H18C	109.5
C9—C10—H10A	109.5	C12—N1—C8	116.13 (12)
C9—C10—H10B	109.5	C12—N1—S1	116.31 (10)
H10A—C10—H10B	109.5	C8—N1—S1	116.30 (10)
C9—C10—H10C	109.5	C14—N2—C13	120.87 (16)
H10A—C10—H10C	109.5	C14—N2—H2A	118.5 (15)
H10B—C10—H10C	109.5	C13—N2—H2A	120.5 (15)
C9—C11—H11A	109.5	C4—O1—C7	117.91 (14)
C9—C11—H11B	109.5	C14—O5—C15	120.56 (12)
H11A—C11—H11B	109.5	O2—S1—O3	119.77 (7)
C9—C11—H11C	109.5	O2—S1—N1	107.05 (7)
H11A—C11—H11C	109.5	O3—S1—N1	106.12 (7)
H11B—C11—H11C	109.5	O2—S1—C1	107.89 (7)
N1—C12—C13	113.64 (13)	O3—S1—C1	107.48 (7)
N1—C12—H12A	108.8	N1—S1—C1	108.05 (7)
C13—C12—H12A	108.8		
C6—C1—C2—C3	-0.4 (2)	C5—C4—O1—C7	-1.8 (2)
S1—C1—C2—C3	177.48 (13)	C3—C4—O1—C7	178.47 (16)
C1—C2—C3—C4	0.1 (3)	O4—C14—O5—C15	0.1 (2)
C2—C3—C4—O1	179.80 (16)	N2—C14—O5—C15	-179.50 (14)
C2—C3—C4—C5	0.1 (3)	C17—C15—O5—C14	-61.01 (19)
O1—C4—C5—C6	-179.78 (15)	C16—C15—O5—C14	63.5 (2)
C3—C4—C5—C6	-0.1 (2)	C18—C15—O5—C14	-178.95 (17)
C2—C1—C6—C5	0.4 (2)	C12—N1—S1—O2	171.49 (10)
S1—C1—C6—C5	-177.50 (12)	C8—N1—S1—O2	-46.10 (12)
C4—C5—C6—C1	-0.1 (2)	C12—N1—S1—O3	42.49 (12)
N1—C8—C9—C11	-57.0 (2)	C8—N1—S1—O3	-175.09 (10)
N1—C8—C9—C10	-179.83 (14)	C12—N1—S1—C1	-72.54 (12)
N1—C12—C13—N2	-70.15 (18)	C8—N1—S1—C1	69.88 (12)
C13—C12—N1—C8	127.99 (14)	C6—C1—S1—O2	-161.35 (12)
C13—C12—N1—S1	-89.53 (14)	C2—C1—S1—O2	20.79 (15)
C9—C8—N1—C12	-78.70 (17)	C6—C1—S1—O3	-30.90 (14)
C9—C8—N1—S1	138.82 (12)	C2—C1—S1—O3	151.23 (13)
O4—C14—N2—C13	2.9 (2)	C6—C1—S1—N1	83.23 (13)
O5—C14—N2—C13	-177.48 (14)	C2—C1—S1—N1	-94.64 (13)
C12—C13—N2—C14	-72.1 (2)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C8—H8A···O2	0.97	2.43	2.9106 (19)	110

C13—H13A···O3	0.97	2.48	3.107 (2)	122
C3—H3···O4 ⁱ	0.93	2.59	3.402 (2)	147
N2—H2A···O4 ⁱⁱ	0.82 (2)	2.38 (2)	3.190 (2)	171 (2)

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $x, y+1, z$.