

N-[(1-Benzoylpiperidin-4-yl)methyl]-benzamide

K. Prathebha,^a D. Reuben Jonathan,^b Sathya Shanmugam^a and G. Usha^{a*}

^aPG and Research Department of Physics, Queen Mary's College, Chennai-4, Tamilnadu, India, and ^bPG and Research Department of Chemistry, Presidency College, Chennai-5, Tamil Nadu, India

Correspondence e-mail: guqmc@yahoo.com


Received 13 March 2014; accepted 2 June 2014

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.043; wR factor = 0.125; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2$, the piperidine ring adopts a chair conformation. The phenyl rings are inclined to one another by $80.1(1)^\circ$ and make dihedral angles of $46.1(1)$ and $40.2(1)^\circ$ with the mean plane of the piperidine ring. In the crystal, pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into inversion dimers. $\text{C}-\text{H}\cdots\text{O}$ interactions further link the molecules, forming a three-dimensional supramolecular network.

Related literature

For the synthesis of the title compound, see: Prathebha *et al.* (2013); Venkatraj *et al.* (2008). For the biological activity of piperidine derivatives, see: Ramalingan *et al.* (2004); Sergeant & May (1970). For bond-length data, see: Allen *et al.* (1987). For related structures, see: Al-abbas *et al.* (2010); Ávila *et al.* (2010). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2$

$M_r = 322.40$

Triclinic, $P\bar{1}$

$a = 9.8039(2)\text{ \AA}$

$b = 10.4453(2)\text{ \AA}$

$c = 10.6765(2)\text{ \AA}$

$\alpha = 62.208(1)^\circ$

$\beta = 66.009(1)^\circ$

$\gamma = 68.150(1)^\circ$

$V = 860.80(3)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 0.08\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.22 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2004)

$T_{\min} = 0.982$, $T_{\max} = 0.984$

12912 measured reflections

3562 independent reflections

2929 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.125$

$S = 1.04$

3531 reflections

217 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.56\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.21\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C13—H13A \cdots O1 ⁱ	0.97	2.60	3.5548 (18)	169
C3—H3 \cdots O2 ⁱⁱ	0.93	2.47	3.3803 (17)	167
N2—H2A \cdots O2 ⁱⁱ	0.86	2.11	2.9401 (15)	162
C8—H8 \cdots O1 ⁱⁱⁱ	0.93	2.52	3.4506 (19)	176

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 2, -y + 1, -z$; (iii) $x, y, z + 1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97*.

The authors thank Professor Velmurugan, Centre for Advanced Study in Crystallography and Biophysics, University of Madras, for providing data-collection and computer facilities.

Supporting information for this paper is available from the IUCr electronic archives (Reference: BT6968).

References

- Al-abbas, A. A., Yarmo, M. A. & Kassim, M. B. (2010). *Acta Cryst. E66*, o2896.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst. 26*, 343–350.
- Ávila, R. M. D., Landre, I. M. R., Souza, T. E., Veloso, M. P. & Doriguetto, A. C. (2010). *Acta Cryst. E66*, o1630.
- Bruker (2004). *APEX2, SAINT, XPREP* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc. 97*, 1354–1358.
- Farrugia, L. J. (2012). *J. Appl. Cryst. 45*, 849–854.
- Prathebha, K., Revathi, B. K., Usha, G., Ponnuswamy, S. & Abdul Basheer, S. (2013). *Acta Cryst. E69*, o1424.
- Ramalingan, C., Balasubramanian, S., Kabilan, S. & Vasudevan, M. (2004). *Eur. J. Med. Chem. 39*, 527–533.
- Sergeant, L. J. & May, E. L. (1970). *J. Med. Chem. 13*, 1061–1063.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Venkatraj, M., Ponnuswamy, S. & Jeyaraman, R. (2008). *Indian J. Chem. Sect. B*, **47**, 411–426.

supporting information

Acta Cryst. (2014). E70, o771 [https://doi.org/10.1107/S1600536814012793]

N-[(1-Benzoylpiperidin-4-yl)methyl]benzamide

K. Prathebha, D. Reuben Jonathan, Sathya Shanmugam and G. Usha

S1. Comment

Biologically active alkaloids of substituted piperidines have been targeted for their total or partial synthesis (Ramalingan *et al.*, 2004). Piperidines are known to have CNS depressant action at low dosage levels and stimulant activity with increased doses. In addition, the nucleus also possesses analgesic, angionic blocking and anesthetic properties as well (Sergeant & May, 1970). We report in this communication, the synthesis and crystal structure of a new piperidine derivative.

The phenyl rings form dihedral angles of 46.1 (1) $^{\circ}$ and 40.2 (1) $^{\circ}$, respectively, with the best plane through the piperidine ring atoms. The C—N distances [1.337 (2)- 1.468 (2) Å] are in the normal range and are in good agreement with values of a similar reported structure (Ávila *et al.*, 2010). The piperidine ring adopts a chair conformation with puckering parameters (Cremer & Pople, 1975) of q2 = 0.0351 (1) Å, phi2 = -50.61 (3) $^{\circ}$ q3 = 0.5633 (1) Å, QT = 0.5644 (2) Å and θ2 = 3.67 (2) $^{\circ}$.


The crystal packing shows N-H \cdots O hydrogen bonds linking the molecules to centrosymmetric dimers (Fig. 2).

S2. Experimental


The procedure (Prathebha *et al.*, 2013, Venkatraj *et al.*, 2008) adopted in the synthesis of the typical diamide is as follows: In a 250 mL round-bottomed flask 4-methyl piperidine (0.01 mol) was taken in, to which 100 mL of ethyl methyl ketone was added and stirred at room temperature. After 5 minutes, triethylamine (0.02 mol) was added and the mixture was stirred for 15 minutes. Then, benzoyl chloride (0.02 mol) was added and the reaction mixture was stirred at room temperature for about 2 h. A white precipitate of triethyl ammonium chloride was formed. It was filtered and the filtrate was evaporated to get the crude product. The crude product was recrystallized twice from ethyl methyl ketone. Melting Point: 127 °C, yield: 85%.

S3. Refinement


H atoms were positioned geometrically and treated as riding on their parent atoms with C—H = 0.93 - 0.97 Å and N—H. 87 with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$ (C-methyl) and = 1.2 $U_{\text{eq}}(\text{N,C})$ for other H atoms.

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

The packing of the molecules in the crystal structure. The dashed lines indicate the hydrogen bonds.

**Figure 3**

Experimental procedure

N -[(1-Benzoylpiperidin-4-yl)methyl]benzamide

Crystal data

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2$

$M_r = 322.40$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.8039 (2) \text{ \AA}$

$b = 10.4453 (2) \text{ \AA}$

$c = 10.6765 (2) \text{ \AA}$ $\alpha = 62.208 (1)^\circ$ $\beta = 66.009 (1)^\circ$ $\gamma = 68.150 (1)^\circ$ $V = 860.80 (3) \text{ \AA}^3$ $Z = 2$ $F(000) = 344$ $D_x = 1.244 \text{ Mg m}^{-3}$ $D_m = 1.188 \text{ Mg m}^{-3}$ D_m measured by not measuredMo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3562 reflections

 $\theta = 2.3\text{--}26.5^\circ$ $\mu = 0.08 \text{ mm}^{-1}$ $T = 293 \text{ K}$

Block, colourless

 $0.22 \times 0.20 \times 0.20 \text{ mm}$ *Data collection*Bruker Kappa APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω and φ scanAbsorption correction: multi-scan
(*SADABS*; Bruker, 2004) $T_{\min} = 0.982$, $T_{\max} = 0.984$

12912 measured reflections

3562 independent reflections

2929 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.028$ $\theta_{\max} = 26.5^\circ$, $\theta_{\min} = 2.3^\circ$ $h = -12 \rightarrow 12$ $k = -13 \rightarrow 13$ $l = -13 \rightarrow 13$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.043$ $wR(F^2) = 0.125$ $S = 1.04$

3531 reflections

217 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0625P)^2 + 0.1352P]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.56 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$
C1	0.8259 (2)	0.13083 (19)	-0.42562 (19)	0.0716 (5)
H1	0.8464	0.0451	-0.4428	0.086*
C2	0.9233 (2)	0.14731 (18)	-0.37349 (19)	0.0687 (4)
H2	1.0097	0.0725	-0.3554	0.082*
C3	0.89322 (16)	0.27493 (16)	-0.34793 (16)	0.0557 (3)
H3	0.9605	0.2865	-0.3147	0.067*
C4	0.76373 (14)	0.38484 (14)	-0.37167 (13)	0.0462 (3)

C5	0.6671 (2)	0.3663 (2)	-0.4234 (2)	0.0706 (4)
H5	0.5791	0.4396	-0.4391	0.085*
C6	0.6987 (2)	0.2410 (2)	-0.4522 (2)	0.0805 (5)
H6	0.6338	0.2313	-0.4895	0.097*
C7	0.75109 (14)	0.87255 (14)	0.19275 (14)	0.0459 (3)
C8	0.68184 (17)	0.89913 (16)	0.32258 (16)	0.0555 (3)
H8	0.6521	0.8227	0.4115	0.067*
C9	0.65695 (19)	1.03945 (18)	0.32005 (18)	0.0655 (4)
H9	0.6091	1.0571	0.4075	0.079*
C10	0.7016 (2)	1.15236 (17)	0.1911 (2)	0.0705 (4)
H10	0.6842	1.2465	0.1904	0.085*
C11	0.7722 (2)	1.12603 (19)	0.0627 (2)	0.0797 (5)
H11	0.8044	1.2023	-0.0254	0.096*
C12	0.7961 (2)	0.98681 (18)	0.06295 (17)	0.0687 (4)
H12	0.8428	0.9704	-0.0251	0.082*
C13	0.74162 (16)	0.54026 (13)	0.13846 (15)	0.0491 (3)
H13A	0.6493	0.5026	0.1824	0.059*
H13B	0.8142	0.4717	0.1942	0.059*
C14	0.80986 (15)	0.55186 (13)	-0.02091 (15)	0.0489 (3)
H14A	0.8293	0.4553	-0.0244	0.059*
H14B	0.9072	0.5801	-0.0608	0.059*
C15	0.70474 (15)	0.66528 (13)	-0.11683 (15)	0.0485 (3)
H15	0.6124	0.6289	-0.0839	0.058*
C16	0.65720 (17)	0.81395 (14)	-0.09792 (16)	0.0545 (3)
H16A	0.7453	0.8581	-0.1435	0.065*
H16B	0.5798	0.8810	-0.1483	0.065*
C17	0.59429 (16)	0.79586 (15)	0.06342 (16)	0.0535 (3)
H17A	0.5726	0.8908	0.0712	0.064*
H17B	0.4992	0.7629	0.1067	0.064*
C18	0.78445 (14)	0.71670 (14)	0.20002 (14)	0.0455 (3)
C19	0.78213 (19)	0.68686 (15)	-0.27908 (16)	0.0573 (3)
H19A	0.7154	0.7658	-0.3369	0.069*
H19B	0.8762	0.7184	-0.3109	0.069*
C20	0.71745 (15)	0.52476 (14)	-0.34233 (14)	0.0483 (3)
N1	0.70509 (12)	0.68746 (11)	0.14379 (12)	0.0470 (3)
N2	0.81811 (14)	0.55385 (12)	-0.30954 (13)	0.0534 (3)
H2A	0.9056	0.4929	-0.3064	0.064*
O1	0.59104 (12)	0.60806 (12)	-0.34789 (13)	0.0697 (3)
O2	0.88170 (13)	0.62016 (11)	0.25938 (13)	0.0689 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0865 (11)	0.0708 (10)	0.0764 (10)	-0.0293 (9)	-0.0096 (9)	-0.0448 (9)
C2	0.0719 (10)	0.0632 (9)	0.0812 (11)	-0.0047 (7)	-0.0229 (8)	-0.0417 (8)
C3	0.0563 (8)	0.0607 (8)	0.0644 (8)	-0.0080 (6)	-0.0217 (6)	-0.0346 (7)
C4	0.0523 (7)	0.0514 (7)	0.0393 (6)	-0.0144 (5)	-0.0127 (5)	-0.0186 (5)
C5	0.0704 (10)	0.0781 (11)	0.0874 (11)	-0.0067 (8)	-0.0393 (9)	-0.0430 (9)

C6	0.0848 (12)	0.0992 (13)	0.0971 (13)	-0.0290 (10)	-0.0304 (10)	-0.0573 (11)
C7	0.0473 (7)	0.0477 (7)	0.0529 (7)	-0.0087 (5)	-0.0160 (5)	-0.0267 (6)
C8	0.0653 (8)	0.0561 (8)	0.0526 (7)	-0.0163 (6)	-0.0129 (6)	-0.0274 (6)
C9	0.0734 (10)	0.0675 (9)	0.0698 (9)	-0.0176 (7)	-0.0082 (8)	-0.0455 (8)
C10	0.0833 (11)	0.0542 (8)	0.0871 (11)	-0.0199 (8)	-0.0168 (9)	-0.0393 (8)
C11	0.1130 (14)	0.0594 (9)	0.0692 (10)	-0.0401 (9)	-0.0106 (10)	-0.0232 (8)
C12	0.0919 (11)	0.0662 (9)	0.0542 (8)	-0.0308 (8)	-0.0042 (8)	-0.0314 (7)
C13	0.0590 (7)	0.0381 (6)	0.0579 (8)	-0.0068 (5)	-0.0231 (6)	-0.0217 (5)
C14	0.0568 (7)	0.0382 (6)	0.0582 (8)	-0.0033 (5)	-0.0214 (6)	-0.0244 (5)
C15	0.0561 (7)	0.0432 (7)	0.0580 (8)	-0.0076 (5)	-0.0229 (6)	-0.0251 (6)
C16	0.0682 (8)	0.0403 (7)	0.0666 (8)	0.0011 (6)	-0.0362 (7)	-0.0249 (6)
C17	0.0558 (7)	0.0474 (7)	0.0720 (9)	0.0045 (6)	-0.0318 (7)	-0.0344 (6)
C18	0.0471 (6)	0.0467 (7)	0.0497 (7)	-0.0062 (5)	-0.0158 (5)	-0.0249 (5)
C19	0.0773 (9)	0.0465 (7)	0.0576 (8)	-0.0108 (6)	-0.0268 (7)	-0.0228 (6)
C20	0.0530 (7)	0.0493 (7)	0.0442 (6)	-0.0080 (6)	-0.0163 (5)	-0.0190 (5)
N1	0.0534 (6)	0.0408 (5)	0.0580 (6)	-0.0019 (4)	-0.0242 (5)	-0.0265 (5)
N2	0.0609 (7)	0.0520 (6)	0.0609 (7)	-0.0032 (5)	-0.0262 (5)	-0.0314 (5)
O1	0.0602 (6)	0.0635 (6)	0.0921 (8)	0.0024 (5)	-0.0320 (6)	-0.0379 (6)
O2	0.0756 (7)	0.0571 (6)	0.0973 (8)	0.0054 (5)	-0.0534 (6)	-0.0382 (6)

Geometric parameters (\AA , $^{\circ}$)

C1—C6	1.369 (3)	C13—N1	1.4665 (14)	
C1—C2	1.377 (2)	C13—C14	1.5166 (18)	
C1—H1	0.9300	C13—H13A	0.9700	
C2—C3	1.3854 (19)	C13—H13B	0.9700	
C2—H2	0.9300	C14—C15	1.5272 (18)	
C3—C4	1.3782 (19)	C14—H14A	0.9700	
C3—H3	0.9300	C14—H14B	0.9700	
C4—C5	1.3788 (19)	C15—C19	1.5236 (19)	
C4—C20	1.5026 (17)	C15—C16	1.5319 (16)	
C5—C6	1.378 (2)	C15—H15	0.9800	
C5—H5	0.9300	C16—C17	1.517 (2)	
C6—H6	0.9300	C16—H16A	0.9700	
C7—C12	1.376 (2)	C16—H16B	0.9700	
C7—C8	1.3830 (18)	C17—N1	1.4610 (16)	
C7—C18	1.5065 (16)	C17—H17A	0.9700	
C8—C9	1.3823 (19)	C17—H17B	0.9700	
C8—H8	0.9300	C18—O2	1.2277 (15)	
C9—C10	1.363 (2)	C18—N1	1.3367 (16)	
C9—H9	0.9300	C19—N2	1.4573 (16)	
C10—C11	1.369 (2)	C19—H19A	0.9700	
C10—H10	0.9300	C19—H19B	0.9700	
C11—C12	1.383 (2)	C20—O1	1.2270 (16)	
C11—H11	0.9300	C20—N2	1.3387 (17)	
C12—H12	0.9300	N2—H2A	0.8600	
C6—C1—C2		119.84 (14)	C13—C14—C15	112.38 (10)

C6—C1—H1	120.1	C13—C14—H14A	109.1
C2—C1—H1	120.1	C15—C14—H14A	109.1
C1—C2—C3	120.26 (15)	C13—C14—H14B	109.1
C1—C2—H2	119.9	C15—C14—H14B	109.1
C3—C2—H2	119.9	H14A—C14—H14B	107.9
C4—C3—C2	120.12 (13)	C19—C15—C14	111.50 (11)
C4—C3—H3	119.9	C19—C15—C16	109.95 (11)
C2—C3—H3	119.9	C14—C15—C16	109.78 (10)
C3—C4—C5	118.85 (13)	C19—C15—H15	108.5
C3—C4—C20	124.40 (11)	C14—C15—H15	108.5
C5—C4—C20	116.74 (12)	C16—C15—H15	108.5
C6—C5—C4	121.14 (15)	C17—C16—C15	111.99 (11)
C6—C5—H5	119.4	C17—C16—H16A	109.2
C4—C5—H5	119.4	C15—C16—H16A	109.2
C1—C6—C5	119.76 (14)	C17—C16—H16B	109.2
C1—C6—H6	120.1	C15—C16—H16B	109.2
C5—C6—H6	120.1	H16A—C16—H16B	107.9
C12—C7—C8	118.99 (12)	N1—C17—C16	110.21 (10)
C12—C7—C18	122.18 (12)	N1—C17—H17A	109.6
C8—C7—C18	118.70 (12)	C16—C17—H17A	109.6
C9—C8—C7	119.91 (13)	N1—C17—H17B	109.6
C9—C8—H8	120.0	C16—C17—H17B	109.6
C7—C8—H8	120.0	H17A—C17—H17B	108.1
C10—C9—C8	120.86 (14)	O2—C18—N1	122.09 (11)
C10—C9—H9	119.6	O2—C18—C7	119.08 (11)
C8—C9—H9	119.6	N1—C18—C7	118.82 (11)
C9—C10—C11	119.43 (14)	N2—C19—C15	113.75 (11)
C9—C10—H10	120.3	N2—C19—H19A	108.8
C11—C10—H10	120.3	C15—C19—H19A	108.8
C10—C11—C12	120.43 (15)	N2—C19—H19B	108.8
C10—C11—H11	119.8	C15—C19—H19B	108.8
C12—C11—H11	119.8	H19A—C19—H19B	107.7
C7—C12—C11	120.36 (14)	O1—C20—N2	121.84 (12)
C7—C12—H12	119.8	O1—C20—C4	120.44 (12)
C11—C12—H12	119.8	N2—C20—C4	117.72 (11)
N1—C13—C14	109.34 (10)	C18—N1—C17	126.14 (10)
N1—C13—H13A	109.8	C18—N1—C13	120.63 (10)
C14—C13—H13A	109.8	C17—N1—C13	112.60 (9)
N1—C13—H13B	109.8	C20—N2—C19	121.23 (11)
C14—C13—H13B	109.8	C20—N2—H2A	119.4
H13A—C13—H13B	108.3	C19—N2—H2A	119.4
C6—C1—C2—C3	-0.1 (3)	C12—C7—C18—O2	107.97 (17)
C1—C2—C3—C4	1.3 (2)	C8—C7—C18—O2	-67.72 (17)
C2—C3—C4—C5	-1.1 (2)	C12—C7—C18—N1	-73.19 (18)
C2—C3—C4—C20	177.72 (13)	C8—C7—C18—N1	111.12 (14)
C3—C4—C5—C6	-0.4 (2)	C14—C15—C19—N2	63.59 (15)
C20—C4—C5—C6	-179.32 (15)	C16—C15—C19—N2	-174.42 (11)

C2—C1—C6—C5	−1.4 (3)	C3—C4—C20—O1	−170.49 (13)
C4—C5—C6—C1	1.7 (3)	C5—C4—C20—O1	8.4 (2)
C12—C7—C8—C9	0.9 (2)	C3—C4—C20—N2	9.15 (19)
C18—C7—C8—C9	176.74 (12)	C5—C4—C20—N2	−172.00 (13)
C7—C8—C9—C10	−0.8 (2)	O2—C18—N1—C17	−176.01 (13)
C8—C9—C10—C11	−0.1 (3)	C7—C18—N1—C17	5.18 (19)
C9—C10—C11—C12	1.0 (3)	O2—C18—N1—C13	−5.8 (2)
C8—C7—C12—C11	0.0 (2)	C7—C18—N1—C13	175.39 (11)
C18—C7—C12—C11	−175.70 (15)	C16—C17—N1—C18	110.77 (14)
C10—C11—C12—C7	−1.0 (3)	C16—C17—N1—C13	−60.11 (14)
N1—C13—C14—C15	−56.11 (14)	C14—C13—N1—C18	−111.16 (13)
C13—C14—C15—C19	174.11 (10)	C14—C13—N1—C17	60.29 (14)
C13—C14—C15—C16	52.02 (15)	O1—C20—N2—C19	−0.2 (2)
C19—C15—C16—C17	−174.15 (11)	C4—C20—N2—C19	−179.78 (11)
C14—C15—C16—C17	−51.14 (15)	C15—C19—N2—C20	89.66 (15)
C15—C16—C17—N1	55.05 (15)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C13—H13A···O1 ⁱ	0.97	2.60	3.5548 (18)	169
C3—H3···O2 ⁱⁱ	0.93	2.47	3.3803 (17)	167
N2—H2A···O2 ⁱⁱ	0.86	2.11	2.9401 (15)	162
C8—H8···O1 ⁱⁱⁱ	0.93	2.52	3.4506 (19)	176

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+2, -y+1, -z$; (iii) $x, y, z+1$.