

Received 23 April 2014
Accepted 1 July 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Keywords: dispiro; pyrrolidine-indole; benzothiophene; crystal structure

CCDC references: 1011623; 1011624


Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structures of ethyl (2*S*^{*},2*R*^{*})-1'-methyl-2'',3-dioxo-2,3-dihydrodispiro[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate and ethyl (2*S*^{*},2*R*^{*})-5''-chloro-1'-methyl-2'',3-dioxo-2,3-dihydrodispiro[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate

M. P. Savithri,^a M. Suresh,^b R. Raghunathan,^b G. Vimala,^c R. Raja^c and A. Subbiah Pandi^{c*}

^aDepartment of Physics, Queen Mary's College (Autonomous), Chennai 600 004, India, ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, and ^cDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India. *Correspondence e-mail: aspandian59@gmail.com


In the title compounds, $C_{22}H_{20}N_2O_4S$, (I), and $C_{22}H_{19}ClN_2O_4S$, (II), the pyrrolidine rings have twist conformations on the spiro–spiro C–C bonds. In (I), the five-membered ring of the oxindole moiety has an envelope conformation with the spiro C atom as the flap, while in (II) this ring is flat (r.m.s. deviation = 0.042 Å). The mean planes of the pyrrolidine rings are inclined to the mean planes of the indole units [r.m.s. deviations = 0.073 and 0.069 Å for (I) and (II), respectively] and the benzothiophene ring systems (r.m.s. deviations = 0.019 and 0.034 Å for (I) and (II), respectively) by 79.57 (8) and 88.61 (7)° for (I), and by 81.99 (10) and 88.79 (10)° for (II). In both compounds, the ethoxycarbonyl group occupies an equatorial position with an extended conformation. The overall conformation of the two molecules differs in the angle of inclination of the indole unit with respect to the benzothiophene ring system, with a dihedral angle between the planes of 71.59 (5) in (I) and 82.27 (7)° in (II). In the crystal of (I), molecules are linked via pairs of N–H···O hydrogen bonds, forming inversion dimers enclosing $R_2^2(14)$ loops. The dimers are linked via C–H···O and bifurcated C–H···O(O) hydrogen bonds, forming sheets lying parallel to (100). In the crystal of (II), molecules are again linked via pairs of N–H···O hydrogen bonds, forming inversion dimers but enclosing smaller $R_2^2(8)$ loops. Here, the dimers are linked by C–H···O hydrogen bonds, forming ribbons propagating along [010].



1. Chemical context

The spiro-indole-pyrrolidine ring system is a frequently encountered structural motif in many biologically important and pharmacologically relevant alkaloids, such as vincristine, vinblastine and spirotypostatins (Cordell, 1981). Highly functionalized pyrrolidines have gained much interest in the past few years as they constitute the main structural element of many natural and synthetic pharmacologically active compounds (Waldmann, 1995). Optically active pyrrolidines have been used as intermediates, chiral ligands or auxiliaries in controlled asymmetric synthesis (Suzuki *et al.*, 1994; Huryn *et al.*, 1991). In view of this importance, the title compounds were synthesized and we report herein on their molecular and crystal structures.

OPEN  ACCESS



2. Structural commentary

The molecular structure of molecule (I) is shown in Fig. 1. The pyrrolidine ring (N_2/C_8-C_{11}) exhibits a twist conformation on bond C_8-C_{11} . The five-membered ring ($N_1/C_{11}-C_{14}$) of the oxindole moiety adopts an envelope conformation with C_{11} as the flap atom. The $C_{12}=O_2$ bond length of $1.213(1)$ Å confirms the presence of a keto group in the indoline moiety. The benzothiophene ring system (S_1/C_1-C_8 ; r.m.s. deviation = 0.019 Å) and the mean plane of the indole ring system ($N_1/C_{11}-C_{18}$; r.m.s. deviation = 0.073 Å) are inclined to one another by $71.59(5)$ °, and are both almost normal to the mean plane of the pyrrolidine ring (N_2/C_8-C_{11}) with dihedral angles of $88.61(17)$ and $79.57(8)$ °, respectively.

The molecular structure of the compound (II) is illustrated in Fig. 2. The overall geometry of the molecule is similar to that of (I). The pyrrolidine ring (N_2/C_8-C_{11}) also adopts a twist conformation on the C_8-C_{11} bond, and the five-membered ring ($N_1/C_{11}-C_{14}$) of the oxindole moiety has an r.m.s. deviation = 0.042 Å. The mean plane of the benzothiophene ring system (S_1/C_1-C_8 ; r.m.s. deviation = 0.034 Å) and the mean plane of the indole ring system ($N_1/C_{11}-C_{18}$; r.m.s.




Figure 1

The molecular structure of molecule (I), with the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

Table 1
Hydrogen-bond geometry (Å, °) for (I).

C_g is the centroid of the C_1-C_6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N_1-H_1N\cdots O_1^i$	0.83 (2)	2.09 (2)	2.890 (2)	164 (2)
$C_3-H_3\cdots O_2^{ii}$	0.93	2.56	3.385 (2)	148
$C_{18}-H_{18}\cdots O_3^{iii}$	0.93	2.56	3.299 (2)	136
$C_{21}-H_{21B}\cdots O_2^{iv}$	0.97	2.59	3.560 (2)	174
$C_2-H_2\cdots C_g^y$	0.93	2.81	3.649 (2)	151

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $x, y - 1, z$; (iii) $x, y, z + 1$; (iv) $-x, -y + 1, -z$; (v) $-x, -y, -z + 1$.

Table 2
Hydrogen-bond geometry (Å, °) for (II).

C_g is the centroid of the C_1-C_6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N_1-H_1N\cdots O_2^i$	0.81 (2)	2.03 (2)	2.842 (2)	172 (2)
$C_{18}-H_{18}\cdots O_3^{ii}$	0.93	2.57	3.496 (3)	171
$C_2-H_2\cdots C_g^{iii}$	0.93	2.83	3.649 (2)	155

Symmetry codes: (i) $-x + 1, -y, -z + 2$; (ii) $x, y - 1, z$; (iii) $-x, -y, -z + 1$.

deviation = 0.069 Å) are inclined to one another by $82.27(7)$ °, and are both almost normal to the mean plane of the pyrrolidine ring (N_2/C_8-C_{11}) with dihedral angles of $88.79(10)$ and $81.99(10)$ °, respectively.

Molecules (I) and (II) differ only in the presence of a chloride atom at position 5 in the oxoindole unit in (II). The conformation of the two molecules differ in the angle of inclination of the indole moiety with respect to the benzothiophene ring system, with a dihedral angle of $71.59(5)$ in (I) and $82.27(7)$ ° in (II). This is illustrated in Fig. 3, which shows a view of the superposition of the two molecules (*Mercury*; Macrae *et al.*, 2008). There is also a small difference in the orientation of the ester function, the $C_{20}-O_4-C_{21}-C_{22}$ torsion angle being $173.44(19)$ in (I) and $162.3(3)$ ° in (II).





Figure 2

The molecular structure of molecule (II), with the atom labelling. Displacement ellipsoids are drawn at the 50% probability level.


**Figure 3**

A view of the molecular superposition of molecules (I) and (II) [red (I); blue (II); Cl atom in (II) is shown as a blue ball (*Mercury*; Macrae *et al.*, 2008)].


3. Supramolecular features

In the crystal of (I), molecules are linked *via* pairs of N—H···O hydrogen bonds, forming inversion dimers enclosing $R_2^2(14)$ loops (Table 1 and Fig. 4). The dimers are linked *via* C—H···O and bifurcated C—H···O(O) hydrogen bonds, forming sheets lying parallel to (100).

In the crystal of (II), molecules are again linked *via* pairs of N—H···O hydrogen bonds, forming inversion dimers but enclosing smaller $R_2^2(8)$ loops (Table 2 and Fig. 5). Here the dimers are linked by C—H···O hydrogen bonds, forming double-stranded chains propagating along [010].

**Figure 4**

The crystal packing of compound (I), viewed along the a axis. The hydrogen bonds are shown as dashed lines (see Table 1 for details; H atoms not involved in hydrogen bonding have been omitted for clarity).

**Figure 5**

A partial view along the a axis of the crystal packing of compound (II). The hydrogen bonds are shown as dashed lines (see Table 2 for details; H atoms not involved in hydrogen bonding have been omitted for clarity).

4. Database survey

A search of the Cambridge Structural Database (Version 5.35, last update November 2013; Allen, 2002) revealed that the title compounds are the first examples of dispiro-indole-pyrrolidine derivatives with a benzothiophene substituent on the pyrrolidine ring creating the second spiro C atom. There are a large number of indole-spiro-pyrrolidine compounds but there was only one hit for a dispiro-indole-pyrrolidine-'cyclopentane-type' compound, namely 4'-(*p*-methoxyphenyl)-1'-methyl-1*H*-indole-3-spiro-2'-pyrrolidine-3'-spiro-1''-cyclopentane-2(3*H*),2''-dione (refcode: ILIMUL; Govind *et al.*, 2003). The geometry of the pyrrolidine and oxindole ring systems of the two molecules compare well with those reported for similar structures, for example, ethyl 1''-benzyl-2''-oxo-2',3',5',6',7',7a'-hexahydro-1'*H*-dispiro[1,2-*b*]-quinoxaline-11,2'-pyrrolizine-3',3''-indoline]-1'-carboxylate monohydrate (refcode: IFOVUW; Kannan *et al.*, 2013a) and methyl 5''-chloro-1',1''-dimethyl-2,2''-dioxodispiro[indoline-3,2'-pyrrolidine-3',3''-indoline]-4'-carboxylate (refcode: IFOQUR; Kannan *et al.*, 2013b).

5. Synthesis and crystallization

The two compounds were prepared in a similar manner using isatin (1.1 mmol) for (I) and 5-chloro isatin (1.1 mmol) for (II). A mixture of (*E*)-ethyl 2-(3-oxobenzo[*b*]thiophen-2(3*H*)-ylidene) acetate (1.0 mmol) and the relevant isatin together with sarcosine (1.1 mmol) was refluxed in methanol (20 ml) until completion of the reaction, as evidenced by TLC analysis. After completion of the reaction, the solvent was evaporated under reduced pressure. The crude reaction mixture was dissolved in dichloromethane (2×50 ml) and washed with water followed by brine solution. The organic layer was separated and dried over sodium sulfate. After filtration, the solvent was evaporation under reduced pressure. The product was separated by column chromatography using hexane and ethyl acetate (9:1) as eluent to give a white solid. This was dissolved in chloroform (3 ml) and heated for 2 min. The resulting solutions were allowed to evaporate slowly at room temperature and yielded colourless block-like crystals of compounds (I) and (II).

Table 3
Experimental details.

	(I)	(II)
Crystal data		
Chemical formula	C ₂₂ H ₂₀ N ₂ O ₄ S	C ₂₂ H ₁₉ ClN ₂ O ₄ S
M _r	408.46	442.90
Crystal system, space group	Triclinic, P <bar{1}< bar=""></bar{1}<>	Triclinic, P <bar{1}< bar=""></bar{1}<>
Temperature (K)	293	293
a, b, c (Å)	8.7196 (4), 10.7874 (5), 11.3488 (5)	10.4678 (5), 10.9074 (5), 11.5652 (5)
α, β, γ (°)	82.624 (2), 82.775 (2), 79.214 (2)	85.973 (2), 65.612 (2), 62.089 (2)
V (Å ³)	1034.27 (8)	1050.26 (9)
Z	2	2
Radiation type	Mo K α	Mo K α
μ (mm ⁻¹)	0.19	0.31
Crystal size (mm)	0.35 × 0.30 × 0.30	0.35 × 0.30 × 0.30
Data collection		
Diffractometer	Bruker Kappa APEXII CCD	Bruker AXS kappa APEX2 CCD
Absorption correction	Multi-scan (SADABS; Sheldrick, 1996)	Multi-scan (SADABS; Sheldrick, 1996)
T _{min} , T _{max}	0.938, 0.946	0.931, 0.940
No. of measured, independent and observed [I > 2σ(I)] reflections	18982, 3745, 3389	16876, 3788, 3178
R _{int}	0.024	0.024
(sin θ/λ) _{max} (Å ⁻¹)	0.600	0.600
Refinement		
R[F ² > 2σ(F ²)], wR(F ²), S	0.032, 0.087, 1.05	0.036, 0.103, 1.11
No. of reflections	3745	3788
No. of parameters	268	276
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.25, -0.17	0.29, -0.27

Computer programs: APEX2, SAINT and XPREP (Bruker, 2004), SHELLS97 and SHELLXL2013 (Sheldrick, 2008), Mercury (Macrae *et al.*, 2008), PLATON (Spek, 2009), and publCIF (Westrip, 2010).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. For both molecules (I) and (II), the NH H atoms were located in difference Fourier maps and freely refined. The C-bound H atoms were included in calculated positions and treated as riding atoms: C—H = 0.93–0.98 Å with U_{iso}(H) = 1.5U_{eq}(C-methyl) and = 1.2U_{eq}(C) for other H atoms.

Acknowledgements

The authors thank Dr Babu Vargheese, SAIF, IIT, Madras, India, for his help with the data collection.

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Bruker (2004). APEX2, SAINT and XPREP. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cordell, G. (1981). In *Introduction to Alkaloids: A Biogenic Approach*. New York: Wiley International.
- Govind, M. M., Selvanayagam, S., Velmurugan, D., Ravikumar, K., Sridhar, G. & Raghunathan, R. (2003). *Acta Cryst.* **E59**, o1438–o1440.
- Huryn, D. M., Trost, B. M. & Fleming, I. (1991). *Comp. Org. Synth.* **1**, 64–74.
- Kannan, P. S., Lanka, S., Thennarasu, S., Vimala, G. & SubbiahPandi, A. (2013a). *Acta Cryst.* **E69**, o854–o855.
- Kannan, P. S., Yuvaraj, P. S., Manivannan, K., Reddy, B. S. R. & SubbiahPandi, A. (2013b). *Acta Cryst.* **E69**, o825–o826.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Suzuki, H., Aoyagi, S. & Kibayashi, C. (1994). *Tetrahedron Lett.* **35**, 6119–6122.
- Waldmann, H. (1995). *Synlett*. pp. 133–141.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2014). E70, 94-97 [doi:10.1107/S1600536814015426]

Crystal structures of ethyl (2S*,2'R*)-1'-methyl-2'',3-dioxo-2,3-dihydrodispiro-[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate and ethyl (2S*,2'R*)-5''-chloro-1'-methyl-2'',3-dioxo-2,3-dihydrodispiro[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate

M. P. Savithri, M. Suresh, R. Raghunathan, G. Vimala, R. Raja and A. Subbiah Pandi

Computing details

For both compounds, data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2008), *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

(I) Ethyl (2S*,2'R*)-1'-methyl-2'',3-dioxo-2,3-dihydrodispiro[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate

Crystal data

C ₂₂ H ₂₀ N ₂ O ₄ S	V = 1034.27 (8) Å ³
M _r = 408.46	Z = 2
Triclinic, <i>P</i> ̄1	F(000) = 428
Hall symbol: -P 1	D _x = 1.312 Mg m ⁻³
a = 8.7196 (4) Å	Mo <i>Kα</i> radiation, λ = 0.71073 Å
b = 10.7874 (5) Å	θ = 2.4–25.0°
c = 11.3488 (5) Å	μ = 0.19 mm ⁻¹
α = 82.624 (2)°	T = 293 K
β = 82.775 (2)°	Block, colourless
γ = 79.214 (2)°	0.35 × 0.30 × 0.30 mm

Data collection

Bruker Kappa APEXII CCD diffractometer	18982 measured reflections 3745 independent reflections 3389 reflections with <i>I</i> > 2σ(<i>I</i>)
Radiation source: fine-focus sealed tube	<i>R</i> _{int} = 0.024
Graphite monochromator	θ _{max} = 25.3°, θ _{min} = 2.4°
ω and φ scans	<i>h</i> = -10→10
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	<i>k</i> = -12→12
<i>T</i> _{min} = 0.938, <i>T</i> _{max} = 0.946	<i>l</i> = -13→13

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.087$
 $S = 1.05$
 3745 reflections
 268 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Hydrogen site location: mixed
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/\sigma^2(F_{\text{o}}^2) + (0.0417P)^2 + 0.2979P$
 where $P = (F_{\text{o}}^2 + 2F_{\text{c}}^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.25 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL2013* (Sheldrick, 2008), $F_{\text{c}}^* = kF_{\text{c}}[1 + 0.001x F_{\text{c}}^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0099 (17)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.30889 (4)	0.13663 (3)	0.21159 (3)	0.03693 (12)
O1	-0.05178 (12)	0.36372 (10)	0.34437 (10)	0.0491 (3)
O2	0.15077 (17)	0.58243 (10)	0.29000 (11)	0.0647 (4)
O3	0.2116 (2)	0.32476 (19)	-0.07349 (11)	0.1024 (6)
O4	0.01130 (15)	0.28291 (12)	0.05809 (10)	0.0580 (3)
N1	0.18901 (15)	0.47295 (12)	0.47309 (12)	0.0449 (3)
H1N	0.145 (2)	0.5301 (17)	0.5141 (16)	0.054*
N2	0.42620 (15)	0.38340 (12)	0.21313 (10)	0.0452 (3)
C1	-0.11000 (18)	0.09770 (16)	0.37801 (14)	0.0466 (4)
H1	-0.1955	0.1524	0.4109	0.056*
C2	-0.1138 (2)	-0.02952 (17)	0.38248 (16)	0.0569 (4)
H2	-0.2027	-0.0613	0.4179	0.068*
C3	0.0142 (2)	-0.11030 (16)	0.33441 (16)	0.0556 (4)
H3	0.0100	-0.1962	0.3379	0.067*
C4	0.14759 (19)	-0.06681 (14)	0.28153 (14)	0.0448 (4)
H4	0.2334	-0.1225	0.2503	0.054*
C5	0.15164 (16)	0.06238 (12)	0.27566 (11)	0.0337 (3)
C6	0.02306 (15)	0.14358 (13)	0.32381 (11)	0.0343 (3)
C7	0.04308 (15)	0.27552 (13)	0.30944 (11)	0.0333 (3)
C8	0.20210 (16)	0.29422 (12)	0.24261 (11)	0.0326 (3)
C9	0.1918 (2)	0.38436 (14)	0.12494 (12)	0.0447 (4)
H9	0.1131	0.4595	0.1411	0.054*
C10	0.3518 (2)	0.42508 (18)	0.10187 (14)	0.0581 (4)
H10A	0.4142	0.3851	0.0355	0.070*
H10B	0.3404	0.5165	0.0835	0.070*
C11	0.30303 (16)	0.36257 (12)	0.30895 (12)	0.0339 (3)
C12	0.20227 (18)	0.48789 (13)	0.35267 (13)	0.0425 (3)

C13	0.28113 (15)	0.36078 (13)	0.51820 (12)	0.0354 (3)
C14	0.35773 (15)	0.29409 (12)	0.42447 (11)	0.0314 (3)
C15	0.46804 (16)	0.18729 (13)	0.44844 (13)	0.0382 (3)
H15	0.5232	0.1431	0.3867	0.046*
C16	0.49556 (18)	0.14672 (15)	0.56609 (14)	0.0480 (4)
H16	0.5709	0.0755	0.5832	0.058*
C17	0.4122 (2)	0.21104 (16)	0.65801 (14)	0.0504 (4)
H17	0.4297	0.1805	0.7365	0.060*
C18	0.30339 (18)	0.31964 (15)	0.63592 (13)	0.0448 (4)
H18	0.2475	0.3632	0.6978	0.054*
C19	0.5353 (2)	0.4616 (2)	0.23767 (18)	0.0693 (5)
H19A	0.6191	0.4613	0.1741	0.104*
H19B	0.5777	0.4281	0.3116	0.104*
H19C	0.4810	0.5470	0.2435	0.104*
C20	0.1438 (2)	0.32731 (17)	0.02394 (14)	0.0562 (4)
C21	-0.0480 (3)	0.2184 (2)	-0.02642 (17)	0.0750 (6)
H21A	0.0326	0.1509	-0.0550	0.090*
H21B	-0.0799	0.2776	-0.0944	0.090*
C22	-0.1847 (4)	0.1653 (3)	0.0376 (2)	0.1083 (9)
H22A	-0.2277	0.1220	-0.0158	0.162*
H22B	-0.2633	0.2329	0.0658	0.162*
H22C	-0.1514	0.1066	0.1043	0.162*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0377 (2)	0.0370 (2)	0.0362 (2)	-0.00533 (14)	0.00290 (14)	-0.01255 (14)
O1	0.0431 (6)	0.0461 (6)	0.0584 (7)	0.0027 (5)	-0.0034 (5)	-0.0240 (5)
O2	0.0975 (10)	0.0318 (6)	0.0652 (8)	0.0029 (6)	-0.0309 (7)	-0.0053 (5)
O3	0.1453 (15)	0.1480 (16)	0.0305 (7)	-0.0721 (13)	0.0019 (8)	-0.0137 (8)
O4	0.0749 (8)	0.0634 (7)	0.0427 (6)	-0.0156 (6)	-0.0209 (6)	-0.0118 (5)
N1	0.0507 (7)	0.0394 (7)	0.0452 (7)	0.0031 (6)	-0.0083 (6)	-0.0197 (6)
N2	0.0532 (7)	0.0504 (7)	0.0361 (6)	-0.0256 (6)	0.0020 (5)	-0.0023 (5)
C1	0.0385 (8)	0.0581 (10)	0.0459 (8)	-0.0140 (7)	-0.0024 (6)	-0.0097 (7)
C2	0.0565 (10)	0.0616 (11)	0.0597 (10)	-0.0318 (9)	-0.0057 (8)	-0.0026 (8)
C3	0.0719 (11)	0.0419 (9)	0.0602 (10)	-0.0243 (8)	-0.0173 (9)	-0.0009 (7)
C4	0.0539 (9)	0.0347 (7)	0.0480 (8)	-0.0051 (6)	-0.0135 (7)	-0.0084 (6)
C5	0.0384 (7)	0.0344 (7)	0.0300 (6)	-0.0060 (5)	-0.0083 (5)	-0.0056 (5)
C6	0.0357 (7)	0.0381 (7)	0.0311 (7)	-0.0073 (6)	-0.0063 (5)	-0.0065 (5)
C7	0.0354 (7)	0.0380 (7)	0.0278 (6)	-0.0023 (6)	-0.0077 (5)	-0.0099 (5)
C8	0.0410 (7)	0.0302 (6)	0.0269 (6)	-0.0054 (5)	-0.0041 (5)	-0.0052 (5)
C9	0.0641 (10)	0.0407 (8)	0.0305 (7)	-0.0127 (7)	-0.0096 (6)	0.0015 (6)
C10	0.0809 (12)	0.0603 (10)	0.0365 (8)	-0.0313 (9)	-0.0019 (8)	0.0055 (7)
C11	0.0410 (7)	0.0304 (7)	0.0317 (7)	-0.0092 (5)	-0.0032 (5)	-0.0050 (5)
C12	0.0529 (9)	0.0322 (7)	0.0456 (8)	-0.0063 (6)	-0.0137 (7)	-0.0091 (6)
C13	0.0343 (7)	0.0383 (7)	0.0364 (7)	-0.0090 (6)	-0.0046 (5)	-0.0098 (6)
C14	0.0312 (6)	0.0332 (7)	0.0322 (7)	-0.0106 (5)	-0.0029 (5)	-0.0056 (5)
C15	0.0329 (7)	0.0383 (7)	0.0448 (8)	-0.0060 (6)	-0.0057 (6)	-0.0088 (6)

C16	0.0470 (8)	0.0441 (8)	0.0553 (9)	-0.0072 (7)	-0.0215 (7)	0.0000 (7)
C17	0.0611 (10)	0.0588 (10)	0.0368 (8)	-0.0214 (8)	-0.0183 (7)	0.0030 (7)
C18	0.0485 (8)	0.0576 (9)	0.0334 (7)	-0.0176 (7)	-0.0034 (6)	-0.0126 (6)
C19	0.0765 (13)	0.0797 (13)	0.0629 (11)	-0.0506 (11)	-0.0017 (9)	-0.0009 (10)
C20	0.0842 (13)	0.0554 (10)	0.0313 (8)	-0.0168 (9)	-0.0142 (8)	0.0018 (7)
C21	0.1095 (17)	0.0731 (13)	0.0542 (11)	-0.0218 (12)	-0.0389 (11)	-0.0129 (9)
C22	0.134 (2)	0.125 (2)	0.0921 (18)	-0.0653 (19)	-0.0405 (17)	-0.0196 (16)

Geometric parameters (\AA , $^{\circ}$)

S1—C5	1.7496 (14)	C9—C20	1.505 (2)
S1—C8	1.8330 (13)	C9—C10	1.522 (2)
O1—C7	1.2109 (16)	C9—H9	0.9800
O2—C12	1.2125 (18)	C10—H10A	0.9700
O3—C20	1.187 (2)	C10—H10B	0.9700
O4—C20	1.327 (2)	C11—C14	1.5063 (18)
O4—C21	1.450 (2)	C11—C12	1.5696 (19)
N1—C12	1.348 (2)	C13—C18	1.378 (2)
N1—C13	1.3972 (19)	C13—C14	1.3904 (18)
N1—H1N	0.827 (18)	C14—C15	1.3773 (19)
N2—C11	1.4561 (17)	C15—C16	1.387 (2)
N2—C19	1.457 (2)	C15—H15	0.9300
N2—C10	1.473 (2)	C16—C17	1.380 (2)
C1—C2	1.373 (2)	C16—H16	0.9300
C1—C6	1.388 (2)	C17—C18	1.380 (2)
C1—H1	0.9300	C17—H17	0.9300
C2—C3	1.381 (3)	C18—H18	0.9300
C2—H2	0.9300	C19—H19A	0.9600
C3—C4	1.375 (2)	C19—H19B	0.9600
C3—H3	0.9300	C19—H19C	0.9600
C4—C5	1.394 (2)	C21—C22	1.486 (3)
C4—H4	0.9300	C21—H21A	0.9700
C5—C6	1.3871 (19)	C21—H21B	0.9700
C6—C7	1.4523 (19)	C22—H22A	0.9600
C7—C8	1.5295 (18)	C22—H22B	0.9600
C8—C9	1.5495 (18)	C22—H22C	0.9600
C8—C11	1.5611 (18)		
C5—S1—C8	92.66 (6)	N2—C11—C8	99.56 (10)
C20—O4—C21	117.69 (15)	C14—C11—C8	116.51 (10)
C12—N1—C13	112.19 (12)	N2—C11—C12	114.06 (11)
C12—N1—H1N	122.9 (12)	C14—C11—C12	101.21 (10)
C13—N1—H1N	124.0 (12)	C8—C11—C12	110.24 (11)
C11—N2—C19	115.58 (13)	O2—C12—N1	126.23 (14)
C11—N2—C10	107.91 (12)	O2—C12—C11	126.46 (14)
C19—N2—C10	115.00 (13)	N1—C12—C11	107.26 (12)
C2—C1—C6	119.16 (15)	C18—C13—C14	122.52 (13)
C2—C1—H1	120.4	C18—C13—N1	127.67 (13)

C6—C1—H1	120.4	C14—C13—N1	109.76 (12)
C1—C2—C3	120.05 (15)	C15—C14—C13	119.22 (12)
C1—C2—H2	120.0	C15—C14—C11	131.91 (12)
C3—C2—H2	120.0	C13—C14—C11	108.80 (11)
C4—C3—C2	121.62 (15)	C14—C15—C16	118.95 (14)
C4—C3—H3	119.2	C14—C15—H15	120.5
C2—C3—H3	119.2	C16—C15—H15	120.5
C3—C4—C5	118.61 (15)	C17—C16—C15	120.63 (14)
C3—C4—H4	120.7	C17—C16—H16	119.7
C5—C4—H4	120.7	C15—C16—H16	119.7
C6—C5—C4	119.78 (13)	C18—C17—C16	121.38 (14)
C6—C5—S1	114.47 (10)	C18—C17—H17	119.3
C4—C5—S1	125.75 (11)	C16—C17—H17	119.3
C5—C6—C1	120.77 (13)	C13—C18—C17	117.16 (14)
C5—C6—C7	113.66 (12)	C13—C18—H18	121.4
C1—C6—C7	125.57 (13)	C17—C18—H18	121.4
O1—C7—C6	126.02 (13)	N2—C19—H19A	109.5
O1—C7—C8	121.74 (12)	N2—C19—H19B	109.5
C6—C7—C8	112.24 (11)	H19A—C19—H19B	109.5
C7—C8—C9	114.43 (11)	N2—C19—H19C	109.5
C7—C8—C11	115.22 (10)	H19A—C19—H19C	109.5
C9—C8—C11	99.72 (10)	H19B—C19—H19C	109.5
C7—C8—S1	106.91 (9)	O3—C20—O4	124.34 (17)
C9—C8—S1	110.04 (9)	O3—C20—C9	125.41 (18)
C11—C8—S1	110.43 (9)	O4—C20—C9	110.23 (14)
C20—C9—C10	115.35 (14)	O4—C21—C22	107.01 (17)
C20—C9—C8	114.02 (12)	O4—C21—H21A	110.3
C10—C9—C8	103.71 (12)	C22—C21—H21A	110.3
C20—C9—H9	107.8	O4—C21—H21B	110.3
C10—C9—H9	107.8	C22—C21—H21B	110.3
C8—C9—H9	107.8	H21A—C21—H21B	108.6
N2—C10—C9	105.51 (12)	C21—C22—H22A	109.5
N2—C10—H10A	110.6	C21—C22—H22B	109.5
C9—C10—H10A	110.6	H22A—C22—H22B	109.5
N2—C10—H10B	110.6	C21—C22—H22C	109.5
C9—C10—H10B	110.6	H22A—C22—H22C	109.5
H10A—C10—H10B	108.8	H22B—C22—H22C	109.5
N2—C11—C14	115.81 (11)		
C6—C1—C2—C3	0.6 (2)	C9—C8—C11—N2	-47.31 (12)
C1—C2—C3—C4	0.1 (3)	S1—C8—C11—N2	68.47 (11)
C2—C3—C4—C5	-0.7 (2)	C7—C8—C11—C14	64.43 (15)
C3—C4—C5—C6	0.6 (2)	C9—C8—C11—C14	-172.58 (11)
C3—C4—C5—S1	-179.58 (11)	S1—C8—C11—C14	-56.80 (13)
C8—S1—C5—C6	-2.05 (11)	C7—C8—C11—C12	-50.11 (14)
C8—S1—C5—C4	178.17 (12)	C9—C8—C11—C12	72.88 (13)
C4—C5—C6—C1	0.0 (2)	S1—C8—C11—C12	-171.35 (9)
S1—C5—C6—C1	-179.75 (11)	C13—N1—C12—O2	-170.75 (15)

C4—C5—C6—C7	−179.06 (12)	C13—N1—C12—C11	6.75 (16)
S1—C5—C6—C7	1.15 (15)	N2—C11—C12—O2	43.7 (2)
C2—C1—C6—C5	−0.7 (2)	C14—C11—C12—O2	168.82 (15)
C2—C1—C6—C7	178.32 (14)	C8—C11—C12—O2	−67.27 (19)
C5—C6—C7—O1	−179.54 (13)	N2—C11—C12—N1	−133.76 (13)
C1—C6—C7—O1	1.4 (2)	C14—C11—C12—N1	−8.68 (14)
C5—C6—C7—C8	0.68 (16)	C8—C11—C12—N1	115.23 (13)
C1—C6—C7—C8	−178.36 (13)	C12—N1—C13—C18	175.59 (14)
O1—C7—C8—C9	−59.73 (17)	C12—N1—C13—C14	−1.69 (17)
C6—C7—C8—C9	120.06 (12)	C18—C13—C14—C15	−4.4 (2)
O1—C7—C8—C11	55.04 (17)	N1—C13—C14—C15	173.00 (12)
C6—C7—C8—C11	−125.17 (12)	C18—C13—C14—C11	178.13 (12)
O1—C7—C8—S1	178.16 (11)	N1—C13—C14—C11	−4.43 (15)
C6—C7—C8—S1	−2.05 (12)	N2—C11—C14—C15	−45.34 (19)
C5—S1—C8—C7	2.25 (9)	C8—C11—C14—C15	71.24 (18)
C5—S1—C8—C9	−122.58 (10)	C12—C11—C14—C15	−169.23 (14)
C5—S1—C8—C11	128.30 (9)	N2—C11—C14—C13	131.64 (12)
C7—C8—C9—C20	−73.99 (17)	C8—C11—C14—C13	−111.78 (12)
C11—C8—C9—C20	162.46 (14)	C12—C11—C14—C13	7.76 (13)
S1—C8—C9—C20	46.39 (16)	C13—C14—C15—C16	2.27 (19)
C7—C8—C9—C10	159.77 (12)	C11—C14—C15—C16	178.99 (13)
C11—C8—C9—C10	36.22 (14)	C14—C15—C16—C17	1.0 (2)
S1—C8—C9—C10	−79.85 (13)	C15—C16—C17—C18	−2.3 (2)
C11—N2—C10—C9	−19.43 (17)	C14—C13—C18—C17	3.1 (2)
C19—N2—C10—C9	−150.08 (15)	N1—C13—C18—C17	−173.84 (14)
C20—C9—C10—N2	−137.43 (14)	C16—C17—C18—C13	0.3 (2)
C8—C9—C10—N2	−12.03 (16)	C21—O4—C20—O3	4.9 (3)
C19—N2—C11—C14	−61.82 (18)	C21—O4—C20—C9	−176.82 (15)
C10—N2—C11—C14	167.85 (12)	C10—C9—C20—O3	−8.5 (3)
C19—N2—C11—C8	172.43 (14)	C8—C9—C20—O3	−128.4 (2)
C10—N2—C11—C8	42.11 (14)	C10—C9—C20—O4	173.27 (14)
C19—N2—C11—C12	55.08 (18)	C8—C9—C20—O4	53.38 (19)
C10—N2—C11—C12	−75.25 (15)	C20—O4—C21—C22	173.44 (19)
C7—C8—C11—N2	−170.30 (11)		

Hydrogen-bond geometry (Å, °)

Cg is the centroid of the C1—C6 ring.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O1 ⁱ	0.83 (2)	2.09 (2)	2.890 (2)	164 (2)
C3—H3···O2 ⁱⁱ	0.93	2.56	3.385 (2)	148
C18—H18···O3 ⁱⁱⁱ	0.93	2.56	3.299 (2)	136
C21—H21B···O2 ^{iv}	0.97	2.59	3.560 (2)	174
C2—H2···Cg ^v	0.93	2.81	3.649 (2)	151

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $x, y-1, z$; (iii) $x, y, z+1$; (iv) $-x, -y+1, -z$; (v) $-x, -y, -z+1$.

(II) Ethyl (2*S*,2'*R)-5''-chloro-1'-methyl-2'',3-dioxo-2,3-dihydrodSpiro[1-benzothiophene-2,3'-pyrrolidine-2',3''-indoline]-4'-carboxylate**

Crystal data

C ₂₂ H ₁₉ ClN ₂ O ₄ S	V = 1050.26 (9) Å ³
M _r = 442.90	Z = 2
Triclinic, P $\bar{1}$	F(000) = 460
Hall symbol: -P 1	D _x = 1.401 Mg m ⁻³
a = 10.4678 (5) Å	Mo K α radiation, λ = 0.71073 Å
b = 10.9074 (5) Å	θ = 2.0–25.0°
c = 11.5652 (5) Å	μ = 0.31 mm ⁻¹
α = 85.973 (2)°	T = 293 K
β = 65.612 (2)°	Block, colourless
γ = 62.089 (2)°	0.35 × 0.30 × 0.30 mm

Data collection

Bruker AXS kappa APEX2 CCD diffractometer	3178 reflections with $I > 2\sigma(I)$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$R_{\text{int}} = 0.024$
$T_{\min} = 0.931$, $T_{\max} = 0.940$	$\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 2.1^\circ$
16876 measured reflections	$h = -12 \rightarrow 10$
3788 independent reflections	$k = -13 \rightarrow 11$
	$l = -13 \rightarrow 12$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: mixed
$R[F^2 > 2\sigma(F^2)] = 0.036$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.103$	$w = 1/[\sigma^2(F_o^2) + (0.0458P)^2 + 0.3849P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.11$	$(\Delta/\sigma)_{\max} < 0.001$
3788 reflections	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
276 parameters	$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$
0 restraints	
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^* / U_{\text{eq}}$
Cl1	0.78453 (7)	-0.07384 (7)	0.27294 (5)	0.06115 (18)
S1	0.47022 (6)	0.40379 (5)	0.61542 (4)	0.04426 (15)
O1	0.24464 (18)	0.30234 (17)	0.93183 (14)	0.0622 (4)
O2	0.5511 (2)	0.14387 (16)	0.96174 (13)	0.0596 (4)
O3	0.5065 (3)	0.6187 (2)	0.8095 (3)	0.1135 (8)
O4	0.2715 (3)	0.62165 (17)	0.89290 (19)	0.0794 (5)
N1	0.5579 (2)	-0.00867 (18)	0.83019 (16)	0.0476 (4)

H1N	0.534 (3)	-0.054 (2)	0.886 (2)	0.057*
N2	0.7301 (2)	0.20503 (19)	0.70036 (16)	0.0493 (4)
C1	0.0252 (3)	0.4697 (2)	0.8114 (2)	0.0612 (6)
H1	-0.0287	0.4448	0.8874	0.073*
C2	-0.0533 (3)	0.5469 (3)	0.7396 (3)	0.0782 (8)
H2	-0.1613	0.5752	0.7672	0.094*
C3	0.0283 (3)	0.5822 (3)	0.6265 (3)	0.0793 (8)
H3	-0.0262	0.6340	0.5788	0.095*
C4	0.1878 (3)	0.5431 (3)	0.5823 (2)	0.0629 (6)
H4	0.2411	0.5676	0.5058	0.075*
C5	0.2670 (2)	0.4661 (2)	0.65526 (19)	0.0444 (4)
C6	0.1862 (2)	0.42945 (19)	0.76880 (18)	0.0443 (4)
C7	0.2853 (2)	0.34936 (19)	0.83467 (17)	0.0420 (4)
C8	0.4562 (2)	0.33081 (19)	0.76494 (16)	0.0378 (4)
C9	0.4971 (3)	0.4035 (2)	0.84540 (19)	0.0498 (5)
H9	0.4533	0.3873	0.9344	0.060*
C10	0.6786 (3)	0.3230 (3)	0.7914 (2)	0.0643 (6)
H10	0.7241	0.3823	0.7485	0.077*
H11	0.7118	0.2898	0.8595	0.077*
C11	0.5959 (2)	0.17760 (19)	0.73600 (16)	0.0378 (4)
C12	0.5630 (2)	0.1056 (2)	0.85934 (18)	0.0448 (4)
C13	0.6002 (2)	-0.03474 (19)	0.69931 (17)	0.0396 (4)
C14	0.6242 (2)	0.07240 (18)	0.63836 (16)	0.0358 (4)
C15	0.6818 (2)	0.0617 (2)	0.50663 (17)	0.0394 (4)
H15	0.7009	0.1311	0.4639	0.047*
C16	0.7104 (2)	-0.0557 (2)	0.43996 (18)	0.0439 (4)
C17	0.6825 (2)	-0.1597 (2)	0.5010 (2)	0.0491 (5)
H17	0.7014	-0.2364	0.4533	0.059*
C18	0.6265 (2)	-0.1503 (2)	0.6330 (2)	0.0492 (5)
H18	0.6074	-0.2198	0.6755	0.059*
C19	0.8815 (3)	0.0837 (3)	0.6825 (3)	0.0811 (8)
H22A	0.9067	0.0101	0.6227	0.122*
H22B	0.8732	0.0515	0.7634	0.122*
H19	0.9647	0.1094	0.6496	0.122*
C20	0.4289 (4)	0.5589 (3)	0.8460 (3)	0.0688 (7)
C21	0.1907 (6)	0.7722 (3)	0.8925 (5)	0.1379 (18)
H20A	0.1867	0.8208	0.9620	0.165*
H20B	0.2502	0.7944	0.8122	0.165*
C22	0.0345 (5)	0.8180 (3)	0.9074 (5)	0.1294 (15)
H21A	-0.0217	0.9186	0.9177	0.194*
H21B	-0.0205	0.7869	0.9820	0.194*
H21C	0.0389	0.7797	0.8328	0.194*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0577 (3)	0.0831 (4)	0.0385 (3)	-0.0329 (3)	-0.0163 (2)	-0.0048 (2)
S1	0.0478 (3)	0.0544 (3)	0.0399 (3)	-0.0309 (2)	-0.0221 (2)	0.0220 (2)

O1	0.0585 (9)	0.0682 (10)	0.0430 (8)	-0.0321 (8)	-0.0069 (7)	0.0171 (7)
O2	0.0964 (12)	0.0631 (9)	0.0399 (8)	-0.0478 (9)	-0.0388 (8)	0.0224 (7)
O3	0.158 (2)	0.0845 (14)	0.163 (2)	-0.0951 (16)	-0.0874 (18)	0.0504 (14)
O4	0.1093 (16)	0.0397 (9)	0.0970 (14)	-0.0343 (10)	-0.0535 (12)	0.0116 (8)
N1	0.0641 (11)	0.0461 (9)	0.0387 (9)	-0.0316 (8)	-0.0234 (8)	0.0172 (7)
N2	0.0486 (9)	0.0683 (11)	0.0483 (9)	-0.0363 (9)	-0.0274 (8)	0.0151 (8)
C1	0.0441 (11)	0.0570 (13)	0.0658 (14)	-0.0223 (10)	-0.0099 (10)	-0.0021 (11)
C2	0.0431 (12)	0.0802 (18)	0.101 (2)	-0.0206 (12)	-0.0317 (14)	0.0097 (16)
C3	0.0645 (16)	0.0875 (19)	0.095 (2)	-0.0303 (14)	-0.0514 (16)	0.0261 (16)
C4	0.0618 (14)	0.0712 (15)	0.0665 (14)	-0.0327 (12)	-0.0387 (12)	0.0247 (12)
C5	0.0468 (11)	0.0438 (10)	0.0456 (11)	-0.0231 (9)	-0.0211 (9)	0.0070 (8)
C6	0.0425 (10)	0.0387 (10)	0.0453 (11)	-0.0183 (8)	-0.0138 (9)	-0.0021 (8)
C7	0.0460 (10)	0.0368 (10)	0.0355 (10)	-0.0215 (8)	-0.0084 (8)	0.0001 (8)
C8	0.0491 (10)	0.0410 (10)	0.0314 (9)	-0.0281 (8)	-0.0180 (8)	0.0115 (7)
C9	0.0785 (14)	0.0518 (12)	0.0426 (11)	-0.0435 (11)	-0.0336 (10)	0.0153 (9)
C10	0.0840 (17)	0.0814 (16)	0.0663 (15)	-0.0578 (14)	-0.0467 (13)	0.0207 (12)
C11	0.0434 (10)	0.0463 (10)	0.0322 (9)	-0.0260 (8)	-0.0197 (8)	0.0136 (7)
C12	0.0554 (11)	0.0462 (11)	0.0364 (10)	-0.0252 (9)	-0.0234 (9)	0.0144 (8)
C13	0.0366 (9)	0.0421 (10)	0.0389 (10)	-0.0175 (8)	-0.0174 (8)	0.0087 (8)
C14	0.0322 (8)	0.0405 (9)	0.0348 (9)	-0.0167 (7)	-0.0159 (7)	0.0069 (7)
C15	0.0341 (9)	0.0495 (11)	0.0367 (9)	-0.0211 (8)	-0.0163 (8)	0.0084 (8)
C16	0.0334 (9)	0.0574 (12)	0.0380 (10)	-0.0188 (9)	-0.0153 (8)	0.0004 (8)
C17	0.0448 (11)	0.0481 (11)	0.0536 (12)	-0.0193 (9)	-0.0226 (9)	-0.0035 (9)
C18	0.0494 (11)	0.0431 (11)	0.0589 (13)	-0.0238 (9)	-0.0254 (10)	0.0100 (9)
C19	0.0530 (14)	0.104 (2)	0.091 (2)	-0.0317 (14)	-0.0412 (14)	0.0100 (16)
C20	0.116 (2)	0.0585 (14)	0.0698 (16)	-0.0582 (16)	-0.0565 (16)	0.0242 (12)
C21	0.190 (5)	0.0420 (16)	0.224 (5)	-0.044 (2)	-0.142 (4)	0.033 (2)
C22	0.142 (4)	0.060 (2)	0.162 (4)	-0.028 (2)	-0.071 (3)	0.034 (2)

Geometric parameters (\AA , $^\circ$)

C11—C16	1.7448 (19)	C8—C11	1.559 (3)
S1—C5	1.755 (2)	C9—C20	1.499 (3)
S1—C8	1.8308 (17)	C9—C10	1.519 (3)
O1—C7	1.198 (2)	C9—H9	0.9800
O2—C12	1.221 (2)	C10—H10	0.9700
O3—C20	1.199 (3)	C10—H11	0.9700
O4—C20	1.318 (3)	C11—C14	1.506 (2)
O4—C21	1.454 (3)	C11—C12	1.568 (2)
N1—C12	1.344 (3)	C13—C18	1.374 (3)
N1—C13	1.399 (2)	C13—C14	1.392 (2)
N1—H1N	0.81 (2)	C14—C15	1.378 (2)
N2—C19	1.453 (3)	C15—C16	1.384 (3)
N2—C10	1.459 (3)	C15—H15	0.9300
N2—C11	1.460 (2)	C16—C17	1.377 (3)
C1—C2	1.376 (4)	C17—C18	1.382 (3)
C1—C6	1.388 (3)	C17—H17	0.9300
C1—H1	0.9300	C18—H18	0.9300

C2—C3	1.379 (4)	C19—H22A	0.9600
C2—H2	0.9300	C19—H22B	0.9600
C3—C4	1.374 (3)	C19—H19	0.9600
C3—H3	0.9300	C21—C22	1.402 (6)
C4—C5	1.390 (3)	C21—H20A	0.9700
C4—H4	0.9300	C21—H20B	0.9700
C5—C6	1.388 (3)	C22—H21A	0.9600
C6—C7	1.460 (3)	C22—H21B	0.9600
C7—C8	1.543 (3)	C22—H21C	0.9600
C8—C9	1.558 (3)		
C5—S1—C8	93.13 (9)	N2—C11—C8	100.04 (14)
C20—O4—C21	116.5 (3)	C14—C11—C8	119.28 (14)
C12—N1—C13	111.90 (16)	N2—C11—C12	114.02 (14)
C12—N1—H1N	120.4 (16)	C14—C11—C12	101.08 (14)
C13—N1—H1N	127.7 (16)	C8—C11—C12	109.68 (14)
C19—N2—C10	114.16 (18)	O2—C12—N1	126.35 (17)
C19—N2—C11	115.48 (18)	O2—C12—C11	125.72 (17)
C10—N2—C11	107.82 (16)	N1—C12—C11	107.87 (15)
C2—C1—C6	119.0 (2)	C18—C13—C14	122.24 (17)
C2—C1—H1	120.5	C18—C13—N1	127.88 (17)
C6—C1—H1	120.5	C14—C13—N1	109.75 (16)
C1—C2—C3	119.9 (2)	C15—C14—C13	119.77 (17)
C1—C2—H2	120.0	C15—C14—C11	130.86 (16)
C3—C2—H2	120.0	C13—C14—C11	109.00 (15)
C4—C3—C2	122.0 (2)	C14—C15—C16	117.82 (17)
C4—C3—H3	119.0	C14—C15—H15	121.1
C2—C3—H3	119.0	C16—C15—H15	121.1
C3—C4—C5	118.1 (2)	C17—C16—C15	122.18 (18)
C3—C4—H4	120.9	C17—C16—Cl1	118.56 (15)
C5—C4—H4	120.9	C15—C16—Cl1	119.27 (15)
C6—C5—C4	120.30 (19)	C16—C17—C18	120.25 (19)
C6—C5—S1	114.19 (14)	C16—C17—H17	119.9
C4—C5—S1	125.49 (17)	C18—C17—H17	119.9
C5—C6—C1	120.6 (2)	C13—C18—C17	117.71 (18)
C5—C6—C7	113.80 (17)	C13—C18—H18	121.1
C1—C6—C7	125.61 (19)	C17—C18—H18	121.1
O1—C7—C6	126.40 (18)	N2—C19—H22A	109.5
O1—C7—C8	121.59 (18)	N2—C19—H22B	109.5
C6—C7—C8	112.01 (15)	H22A—C19—H22B	109.5
C7—C8—C9	113.63 (15)	N2—C19—H19	109.5
C7—C8—C11	115.91 (14)	H22A—C19—H19	109.5
C9—C8—C11	100.13 (14)	H22B—C19—H19	109.5
C7—C8—S1	106.50 (12)	O3—C20—O4	124.3 (3)
C9—C8—S1	110.40 (12)	O3—C20—C9	124.8 (3)
C11—C8—S1	110.23 (12)	O4—C20—C9	110.9 (2)
C20—C9—C10	114.2 (2)	C22—C21—O4	110.4 (3)
C20—C9—C8	114.23 (17)	C22—C21—H20A	109.6

C10—C9—C8	104.53 (16)	O4—C21—H20A	109.6
C20—C9—H9	107.9	C22—C21—H20B	109.6
C10—C9—H9	107.9	O4—C21—H20B	109.6
C8—C9—H9	107.9	H20A—C21—H20B	108.1
N2—C10—C9	105.75 (16)	C21—C22—H21A	109.5
N2—C10—H10	110.6	C21—C22—H21B	109.5
C9—C10—H10	110.6	H21A—C22—H21B	109.5
N2—C10—H11	110.6	C21—C22—H21C	109.5
C9—C10—H11	110.6	H21A—C22—H21C	109.5
H10—C10—H11	108.7	H21B—C22—H21C	109.5
N2—C11—C14	113.34 (15)		
C6—C1—C2—C3	0.5 (4)	S1—C8—C11—N2	71.69 (14)
C1—C2—C3—C4	-0.3 (5)	C7—C8—C11—C14	68.6 (2)
C2—C3—C4—C5	-0.2 (4)	C9—C8—C11—C14	-168.71 (15)
C3—C4—C5—C6	0.5 (3)	S1—C8—C11—C14	-52.40 (18)
C3—C4—C5—S1	179.1 (2)	C7—C8—C11—C12	-47.1 (2)
C8—S1—C5—C6	-4.41 (16)	C9—C8—C11—C12	75.54 (16)
C8—S1—C5—C4	177.0 (2)	S1—C8—C11—C12	-168.15 (12)
C4—C5—C6—C1	-0.3 (3)	C13—N1—C12—O2	-170.6 (2)
S1—C5—C6—C1	-179.02 (16)	C13—N1—C12—C11	6.5 (2)
C4—C5—C6—C7	-179.70 (19)	N2—C11—C12—O2	49.3 (3)
S1—C5—C6—C7	1.6 (2)	C14—C11—C12—O2	171.21 (19)
C2—C1—C6—C5	-0.2 (3)	C8—C11—C12—O2	-62.0 (2)
C2—C1—C6—C7	179.1 (2)	N2—C11—C12—N1	-127.93 (18)
C5—C6—C7—O1	-177.65 (19)	C14—C11—C12—N1	-5.98 (19)
C1—C6—C7—O1	3.0 (3)	C8—C11—C12—N1	120.84 (17)
C5—C6—C7—C8	2.9 (2)	C12—N1—C13—C18	171.55 (19)
C1—C6—C7—C8	-176.43 (18)	C12—N1—C13—C14	-4.3 (2)
O1—C7—C8—C9	-63.4 (2)	C18—C13—C14—C15	-2.4 (3)
C6—C7—C8—C9	116.04 (17)	N1—C13—C14—C15	173.75 (15)
O1—C7—C8—C11	51.8 (2)	C18—C13—C14—C11	-176.13 (16)
C6—C7—C8—C11	-128.75 (16)	N1—C13—C14—C11	0.0 (2)
O1—C7—C8—S1	174.81 (16)	N2—C11—C14—C15	-46.9 (2)
C6—C7—C8—S1	-5.73 (18)	C8—C11—C14—C15	70.5 (2)
C5—S1—C8—C7	5.57 (13)	C12—C11—C14—C15	-169.32 (18)
C5—S1—C8—C9	-118.22 (15)	N2—C11—C14—C13	125.89 (16)
C5—S1—C8—C11	132.08 (13)	C8—C11—C14—C13	-116.74 (16)
C7—C8—C9—C20	-78.9 (2)	C12—C11—C14—C13	3.47 (18)
C11—C8—C9—C20	156.84 (19)	C13—C14—C15—C16	1.4 (2)
S1—C8—C9—C20	40.7 (2)	C11—C14—C15—C16	173.56 (17)
C7—C8—C9—C10	155.62 (16)	C14—C15—C16—C17	0.3 (3)
C11—C8—C9—C10	31.38 (17)	C14—C15—C16—Cl1	-179.61 (13)
S1—C8—C9—C10	-84.80 (16)	C15—C16—C17—C18	-1.0 (3)
C19—N2—C10—C9	-153.17 (19)	Cl1—C16—C17—C18	178.82 (14)
C11—N2—C10—C9	-23.4 (2)	C14—C13—C18—C17	1.6 (3)
C20—C9—C10—N2	-132.10 (19)	N1—C13—C18—C17	-173.82 (18)
C8—C9—C10—N2	-6.6 (2)	C16—C17—C18—C13	0.1 (3)

C19—N2—C11—C14	−59.7 (2)	C21—O4—C20—O3	3.5 (4)
C10—N2—C11—C14	171.34 (16)	C21—O4—C20—C9	−176.8 (3)
C19—N2—C11—C8	172.22 (17)	C10—C9—C20—O3	−0.9 (4)
C10—N2—C11—C8	43.22 (18)	C8—C9—C20—O3	−121.1 (3)
C19—N2—C11—C12	55.3 (2)	C10—C9—C20—O4	179.43 (19)
C10—N2—C11—C12	−73.7 (2)	C8—C9—C20—O4	59.2 (3)
C7—C8—C11—N2	−167.27 (14)	C20—O4—C21—C22	162.3 (3)
C9—C8—C11—N2	−44.62 (16)		

Hydrogen-bond geometry (Å, °)

Cg is the centroid of the C1—C6 ring.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···O2 ⁱ	0.81 (2)	2.03 (2)	2.842 (2)	172 (2)
C18—H18···O3 ⁱⁱ	0.93	2.57	3.496 (3)	171
C2—H2···Cg ⁱⁱⁱ	0.93	2.83	3.649 (2)	155

Symmetry codes: (i) $-x+1, -y, -z+2$; (ii) $x, y-1, z$; (iii) $-x, -y, -z+1$.