

# Crystal structure of 4-(prop-2-yn-1-yloxy)benzonitrile

Mayu Kanagawa and Tsunehisa Okuno\*

Department of Material Science and Chemistry, Wakayama University, Sakaedani, Wakayama, 640-8510, Japan. \*Correspondence e-mail: okuno@center.wakayama-u.ac.jp

Received 22 November 2014; accepted 24 December 2014

Edited by W. T. A. Harrison, University of Aberdeen, Scotland


In the title compound, C<sub>10</sub>H<sub>7</sub>NO, the dihedral angle between the aromatic ring and the prop-2-yn-1-yloxy grouping is 9.47 (10)°. The bond lengths indicate electronic conjugation between the cyano group, the benzene ring and the propyn-yloxy oxygen atom. In the crystal, a hydrogen bond between the acetylenic C—H atom and the cyano nitrogen atom link the molecules into wave-like [30 $\bar{1}$ ] C(11) chains. These chains are connected by Csp<sup>2</sup>—H···π<sub>ac</sub> (π<sub>ac</sub> is the acetylenic C—C triple bond) close contacts [2.794 (1) Å], resulting in a rolling sheet structure parallel to the *ac* plane and aromatic π—π stacking interactions between the sheets [centroid—centroid distance = 3.593 (2) Å] generate a three-dimensional network.

**Keywords:** crystal structure; prop-2-yn-1-yloxy; hydrogen bonding; C—H···π interactions; π—π stacking interactions.

**CCDC reference:** 1041123

## 1. Related literature

The title compound is an aryl propargyl ether derivative which attracts interest with regard to Claisen rearrangement (Kenny *et al.* 2006; Wang *et al.* 2012) or cleavage of the O—CH<sub>2</sub> bond by boron reagents (Yao *et al.* 2009). For related structures of 4-(prop-2-yn-1-yloxy)benzenes, see: Lindeman *et al.* (1993); Zhu *et al.* (2006); Zhang *et al.* (2008); Marsh (2009); Ranjith *et al.* (2010); Li *et al.* (2009); Ao *et al.* (2011); Al-Mehana *et al.* (2011); Belay *et al.* (2012); Doi & Okuno (2013).



## 2. Experimental

### 2.1. Crystal data

C <sub>10</sub> H <sub>7</sub> NO	<i>V</i> = 781.7 (9) Å <sup>3</sup>
<i>M<sub>r</sub></i> = 157.17	<i>Z</i> = 4
Monoclinic, <i>P</i> 2 <sub>1</sub> / <i>n</i>	Mo <i>K</i> α radiation
<i>a</i> = 6.033 (4) Å	<i>μ</i> = 0.09 mm <sup>-1</sup>
<i>b</i> = 7.393 (5) Å	<i>T</i> = 93 K
<i>c</i> = 17.527 (11) Å	0.20 × 0.07 × 0.03 mm
<i>β</i> = 90.836 (11)°	

### 2.2. Data collection

Rigaku Saturn724+ diffractometer	6174 measured reflections
Absorption correction: numerical ( <i>NUMABS</i> ; Rigaku, 1999)	1795 independent reflections
<i>T<sub>min</sub></i> = 0.986, <i>T<sub>max</sub></i> = 0.997	1457 reflections with <i>F</i> <sup>2</sup> > 2.0σ( <i>F</i> <sup>2</sup> )
	<i>R<sub>int</sub></i> = 0.046

### 2.3. Refinement

<i>R</i> [ <i>F</i> <sup>2</sup> > 2σ( <i>F</i> <sup>2</sup> )] = 0.050	H atoms treated by a mixture of independent and constrained refinement
<i>wR</i> ( <i>F</i> <sup>2</sup> ) = 0.120	<i>Δρ<sub>max</sub></i> = 0.22 e Å <sup>-3</sup>
<i>S</i> = 1.08	<i>Δρ<sub>min</sub></i> = -0.19 e Å <sup>-3</sup>
1795 reflections	
113 parameters	

**Table 1**

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C10—H1···N1 <sup>i</sup>	0.94 (2)	2.41 (2)	3.300 (3)	158.18 (11)
C6—H6···C10 <sup>ii</sup>	0.95	2.79	3.616 (3)	145

Symmetry codes: (i) *x* -  $\frac{3}{2}$ , -*y* +  $\frac{3}{2}$ , *z* +  $\frac{1}{2}$ ; (ii) *x* +  $\frac{1}{2}$ , -*y* +  $\frac{3}{2}$ , *z* -  $\frac{1}{2}$ .

Data collection: *CrystalClear* (Rigaku, 2008); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXD2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *CrystalStructure* (Rigaku, 2014).

## Acknowledgements

This work was supported by Research for Promoting Technological Seeds from the Japan Science and Technology Agency (JST).

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7326).

## References

- Al-Mehana, W. N. A., Yahya, R. & Lo, K. M. (2011). *Acta Cryst.* **E67**, o2900.  
 Ao, L., Tu, J.-H., Huang, X. & Ding, B.-Y. (2011). *Acta Cryst.* **E67**, o2642.  
 Belay, Y. H., Kinfe, H. H. & Muller, A. (2012). *Acta Cryst.* **E68**, o3072.  
 Doi, I. & Okuno, T. (2013). *Acta Cryst.* **E69**, o125.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 Kenny, R. S., Mashelkar, U. C., Rane, D. M. & Bezawada, D. K. (2006). *Tetrahedron*, **62**, 9280–9288.  
 Li, Z.-X., Ren, C.-M., Yang, S., Yao, G.-J. & Shi, Q.-Z. (2009). *Acta Cryst.* **E65**, o65.  
 Lindeman, S. V., Dvorikova, R. A., Gol'ding, I. R., Struchkov, Y. T. & Teplyakov, M. M. (1993). *Russ. Chem. Bull.* **42**, 1601–1605.

- Marsh, R. E. (2009). *Acta Cryst.* **B65**, 782–783.
- Ranjith, S., Thirunarayanan, A., Raja, S., Rajakumar, P. & SubbiahPandi, A. (2010). *Acta Cryst.* **E66**, o2261–o2262.
- Rigaku (1999). *NUMABS*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2008). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (2014). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, Y., Ji, K., Lan, S. & Zhang, L. (2012). *Angew. Chem. Int. Ed.* **51**, 1915–1918.
- Yao, M. L., Reddy, M. S., Zeng, W., Hall, K., Walfish, I. & Kabalka, G. W. (2009). *J. Org. Chem.* **74**, 1385–1387.
- Zhang, W., Yao, L. & Tao, R.-J. (2008). *Acta Cryst.* **E64**, o307.
- Zhu, N., Lightsey, D., Foroozesh, M., Alworth, W., Chaudhary, A., Willett, K. L. & Stevens, C. L. K. (2006). *J. Chem. Cryst.* **36**, 289–296.

## supporting information

*Acta Cryst.* (2015). E71, o97–o98 [doi:10.1107/S2056989014028035]

## Crystal structure of 4-(prop-2-yn-1-yloxy)benzotrile

Mayu Kanagawa and Tsunehisa Okuno

### S1. Comment

The title compound, C<sub>10</sub>H<sub>7</sub>N<sub>1</sub>O<sub>1</sub>, is an aryl propargyl ether derivative which attracts interest from viewpoints of Claisen rearrangement (Kenny *et al.* 2006; Wang *et al.* 2012) or cleavage of O–CH<sub>2</sub> bond by boron reagents (Yao *et al.* 2009). In these reactions, a direction of the lone pair of the oxygen has large influence upon reactivity.

The molecule has an almost planar structure (atoms C1—C10/N1/O1 are essentially co-planar with r.m.s. deviation = 0.0862 Å), indicating an effective conjugation of the cyano group, the C1—C6 benzene ring and the lone pair of the O1 (Fig. 1). This is presumably because push-pull effect between an electron donating alkyloxy group and an electron withdrawing cyano group (Zhu *et al.* 2006; Marsh 2009; Ranjith *et al.* 2010; Ao *et al.* 2011; Al-Mehana *et al.* 2011; Belay *et al.* 2012; Doi & Okuno 2013).

In the crystal, C10–H1<sup>i</sup>⋯N1<sup>i</sup> hydrogen bonds [Symmetry code: (i)  $x - 3/2, -y + 3/2, z + 1/2$ ] connect the molecules to make a one-dimensional wavy chain. Intermolecular C6–H6<sup>ii</sup>⋯C10<sup>ii</sup> interaction [Symmetry code: (ii)  $x + 1/2, -y + 3/2, z - 1/2$ ], whose distance is 2.794 (1) Å, binds the chains to form a rolling sheet structure as shown in Fig. 2.


Fig. 3 shows  $\pi$ ⋯ $\pi$  stacking interactions between the sheets, where the centroid to centroid distance is 3.593 (2) Å and the C3<sup>v</sup>⋯C5<sup>v</sup> is 3.387 (3) Å [Symmetry code: (v)  $-x + 2, -y + 1, -z$ ]. The molecules also form weak intersheet C5–H5<sup>vi</sup>⋯O1<sup>vi</sup> bonds whose distance is 2.690 (1) Å [Symmetry code: (vi)  $-x + 1, -y + 1, -z$ ]. In this crystal, the intermolecular hydrogen bonds, the C–H⋯ $\pi$  interactions and the  $\pi$ ⋯ $\pi$  stacking interactions are found to make a three-dimensional molecular network.

### S2. Experimental


The title compound is commercially available. Colourless platelets of sufficient quality for diffraction measurements were prepared by sublimation at room temperature.

### S3. Refinement


The C-bound H atoms except for C<sub>sp</sub>–H were placed at ideal positions and were refined as riding on their parent C atoms.  $U_{\text{iso}}(\text{H})$  values of the H atoms were set at  $1.2U_{\text{eq}}(\text{parent atom})$ . The C<sub>sp</sub>-bound H atom was obtained from a difference Fourier map and was refined isotropically without any restrictions.

**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level and H atoms are shown as small spheres.

**Figure 2**

Part of the crystal structure showing the rolling sheet structure formed by the C–H $\cdots$ N and C–H $\cdots$  $\pi$  hydrogen bonds [Symmetry codes: (i)  $x - 3/2, -y + 3/2, z + 1/2$ ; (ii)  $x + 1/2, -y + 3/2, z - 1/2$ ; (iii)  $x + 3/2, -y + 3/2, z - 1/2$ ; (iv)  $x - 1/2, -y + 3/2, z + 1/2$ ].

**Figure 3**

Part of the crystal structure showing the intersheet  $\pi\cdots\pi$  stacking interactions and the weak C–H $\cdots$ O hydrogen bonds [Symmetry codes: (v)  $-x + 2, -y + 1, -z$ ; (vi)  $-x + 1, -y + 1, -z$ ].

#### 4-(Prop-2-yn-1-yloxy)benzonitrile

##### Crystal data

$C_{10}H_7NO$

$M_r = 157.17$

Monoclinic,  $P2_1/n$

$a = 6.033$  (4) Å

$b = 7.393$  (5) Å

$c = 17.527$  (11) Å

$\beta = 90.836$  (11)°

$V = 781.7$  (9) Å<sup>3</sup>

$Z = 4$

$F(000) = 328.00$

$D_x = 1.335$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71075$  Å

Cell parameters from 2559 reflections

$\theta = 2.3$ – $31.1$ °

$\mu = 0.09$  mm<sup>-1</sup>

$T = 93$  K

Platelet, colorless

$0.20 \times 0.07 \times 0.03$  mm

##### Data collection

Rigaku Saturn724+  
diffractometer

Detector resolution: 28.445 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: numerical  
(*NUMABS*; Rigaku, 1999)

$T_{\min} = 0.986$ ,  $T_{\max} = 0.997$

6174 measured reflections

1795 independent reflections

1457 reflections with  $F^2 > 2\sigma(F^2)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 27.5$ °,  $\theta_{\min} = 2.3$ °

$h = -7 \rightarrow 7$   
 $k = -9 \rightarrow 9$

$l = -21 \rightarrow 22$

### Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.050$   
 $wR(F^2) = 0.120$   
 $S = 1.08$   
 1795 reflections  
 113 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0584P)^2 + 0.1471P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** ENTER SPECIAL DETAILS OF THE MOLECULAR GEOMETRY

**Refinement.** Refinement was performed using all reflections. The weighted  $R$ -factor ( $wR$ ) and goodness of fit ( $S$ ) are based on  $F^2$ .  $R$ -factor (gt) are based on  $F$ . The threshold expression of  $F^2 > 2.0 \sigma(F^2)$  is used only for calculating  $R$ -factor (gt).

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.66646 (16)	0.60053 (14)	0.09599 (6)	0.0181 (3)
N1	1.4021 (2)	0.95596 (18)	-0.14609 (7)	0.0234 (3)
C1	1.1274 (2)	0.81486 (19)	-0.05087 (8)	0.0166 (3)
C2	1.1843 (2)	0.79920 (19)	0.02615 (8)	0.0175 (3)
C3	1.0358 (2)	0.72770 (19)	0.07719 (8)	0.0168 (3)
C4	0.8276 (2)	0.67109 (18)	0.05072 (8)	0.0152 (3)
C5	0.7705 (2)	0.68441 (19)	-0.02634 (8)	0.0162 (3)
C6	0.9187 (2)	0.75633 (19)	-0.07715 (8)	0.0168 (3)
C7	1.2805 (2)	0.8928 (2)	-0.10375 (8)	0.0176 (3)
C8	0.7121 (2)	0.5908 (2)	0.17643 (8)	0.0197 (3)
C9	0.5068 (2)	0.5386 (2)	0.21392 (8)	0.0190 (3)
C10	0.3416 (3)	0.4999 (2)	0.24560 (9)	0.0235 (4)
H1	0.211 (3)	0.479 (3)	0.2728 (11)	0.033 (5)*
H2	1.32622	0.838	0.04368	0.0210*
H3	1.07475	0.71712	0.12971	0.0201*
H5	0.62929	0.64389	-0.0439	0.0195*
H6	0.8798	0.76623	-0.1297	0.0202*
H8A	0.76355	0.70976	0.19566	0.0236*
H8B	0.82942	0.50026	0.18705	0.0236*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0144 (5)	0.0253 (6)	0.0147 (5)	-0.0035 (4)	0.0030 (4)	0.0018 (4)
N1	0.0173 (6)	0.0298 (7)	0.0233 (7)	0.0007 (6)	0.0032 (5)	0.0022 (6)
C1	0.0143 (7)	0.0156 (7)	0.0199 (8)	0.0023 (5)	0.0042 (6)	-0.0003 (6)

C2	0.0117 (7)	0.0177 (7)	0.0231 (8)	0.0021 (5)	0.0005 (6)	-0.0021 (6)
C3	0.0146 (7)	0.0199 (7)	0.0158 (7)	0.0018 (5)	-0.0005 (6)	0.0003 (6)
C4	0.0131 (6)	0.0141 (7)	0.0186 (7)	0.0011 (5)	0.0046 (5)	-0.0006 (6)
C5	0.0119 (7)	0.0174 (7)	0.0193 (8)	0.0003 (5)	-0.0005 (5)	-0.0007 (6)
C6	0.0160 (7)	0.0189 (7)	0.0155 (7)	0.0015 (6)	0.0004 (6)	-0.0003 (6)
C7	0.0141 (7)	0.0195 (7)	0.0193 (8)	0.0012 (6)	-0.0001 (6)	-0.0011 (6)
C8	0.0155 (7)	0.0276 (8)	0.0161 (8)	-0.0004 (6)	0.0024 (6)	0.0006 (6)
C9	0.0179 (7)	0.0239 (8)	0.0151 (7)	0.0018 (6)	-0.0002 (6)	0.0005 (6)
C10	0.0182 (7)	0.0339 (9)	0.0185 (8)	-0.0014 (7)	0.0027 (6)	0.0000 (7)

*Geometric parameters (Å, °)*

O1—C4	1.3673 (18)	C8—C9	1.463 (2)
O1—C8	1.434 (2)	C9—C10	1.183 (2)
N1—C7	1.150 (2)	C2—H2	0.950
C1—C2	1.393 (2)	C3—H3	0.950
C1—C6	1.403 (2)	C5—H5	0.950
C1—C7	1.438 (2)	C6—H6	0.950
C2—C3	1.380 (2)	C8—H8A	0.990
C3—C4	1.397 (2)	C8—H8B	0.990
C4—C5	1.392 (2)	C10—H1	0.94 (2)
C5—C6	1.378 (2)		
C4—O1—C8	117.49 (11)	C1—C2—H2	119.768
C2—C1—C6	119.99 (13)	C3—C2—H2	119.770
C2—C1—C7	120.43 (13)	C2—C3—H3	120.380
C6—C1—C7	119.58 (13)	C4—C3—H3	120.387
C1—C2—C3	120.46 (13)	C4—C5—H5	119.993
C2—C3—C4	119.23 (13)	C6—C5—H5	119.976
O1—C4—C3	124.41 (13)	C1—C6—H6	120.174
O1—C4—C5	114.96 (12)	C5—C6—H6	120.181
C3—C4—C5	120.63 (13)	O1—C8—H8A	110.181
C4—C5—C6	120.03 (13)	O1—C8—H8B	110.187
C1—C6—C5	119.65 (13)	C9—C8—H8A	110.179
N1—C7—C1	179.63 (15)	C9—C8—H8B	110.175
O1—C8—C9	107.64 (12)	H8A—C8—H8B	108.478
C8—C9—C10	178.27 (16)	C9—C10—H1	175.1 (12)
C4—O1—C8—C9	171.29 (10)	C1—C2—C3—C4	0.0 (2)
C8—O1—C4—C3	2.52 (18)	C2—C3—C4—O1	-179.13 (12)
C8—O1—C4—C5	-177.34 (10)	C2—C3—C4—C5	0.7 (2)
C2—C1—C6—C5	0.4 (2)	O1—C4—C5—C6	178.94 (10)
C6—C1—C2—C3	-0.6 (2)	C3—C4—C5—C6	-0.9 (2)
C7—C1—C2—C3	178.70 (12)	C4—C5—C6—C1	0.4 (2)
C7—C1—C6—C5	-178.91 (11)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C10—H1···N1 <sup>i</sup>	0.94 (2)	2.41 (2)	3.300 (3)	158.18 (11)
C6—H6···C10 <sup>ii</sup>	0.95	2.79	3.616 (3)	145

Symmetry codes: (i)  $x-3/2, -y+3/2, z+1/2$ ; (ii)  $x+1/2, -y+3/2, z-1/2$ .