

Crystal structure of (+)-methyl (*E*)-3-[(2*S*,4*S*,5*R*)-2-amino-5-hydroxymethyl-2-trichloromethyl-1,3-dioxolan-4-yl]-2-methylprop-2-enoate

Takeshi Oishi,^{a*} Daichi Yasushima,^b Kihiro Yuasa,^b Takaaki Sato^b and Noritaka Chida^b

Received 22 January 2016
 Accepted 10 February 2016

^aSchool of Medicine, Keio University, Hiyoshi 4-1-1, Kohoku-ku, Yokohama 223-8521, Japan, and ^bDepartment of Applied Chemistry, Faculty of Science and Technology, Keio University, Hiyoshi 3-14-1, Kohoku-ku, Yokohama 223-8522, Japan. *Correspondence e-mail: oec@keio.jp

Edited by H. Ishida, Okayama University, Japan

Keywords: crystal structure; orthoamide; dioxolane; hydrogen bonding; hydroxy group; amino group; disorder.



CCDC reference: 1452471

Supporting information: this article has supporting information at journals.iucr.org/e

In the title compound, C₁₀H₁₄Cl₃NO₅, the five-membered dioxolane ring adopts an envelope conformation. The C atom at the flap, which is bonded to the hydroxymethyl substituent, deviates from the mean plane of other ring atoms by 0.357 (5) Å. There are two intramolecular hydrogen bonds (O—H···N and N—H···O) between the hydroxy and amino groups, so that O- and N-bound H atoms involved in these hydrogen bonds are each disordered with equal occupancies of 0.50. The methyl 2-methylprop-2-enoate substituent also shows a disordered structure over two sets of sites with refined occupancies of 0.482 (5) and 0.518 (5). In the crystal, molecules are connected into a dimer by an O—H···O hydrogen bond. The dimers are further linked by N—H···O, C—H···N and C—H···O interactions, extending a sheet structure parallel to (101).


1. Chemical context

The 3,3-sigmatropic rearrangement of an allylic trichloroacetimidate (Overman rearrangement; Overman, 1974, 1976) is one of the most important reactions in organic chemistry. It has been utilized as a quite powerful tool to introduce the nitrogen functional group because this imidate is easily available from an allylic alcohol with trichloroacetonitrile (Cl₃CC≡N). In the case of a diol, a cyclic orthoamide (2-amino-2-trichloromethyl-1,3-dioxolane) may be afforded by controlling the reaction conditions though bis-imidates are usually produced. We have explored the rearrangement of the cyclic orthoamide prepared from a contiguous diol or triol, and have developed a novel strategy for the total synthesis of certain natural products (Nakayama *et al.*, 2013). As part of our ongoing studies in this area, we now describe the synthesis and structure of the title compound.




2. Structural commentary

The molecular structure of the title compound is shown in Fig. 1. The dioxolane ring (C5/O7/C8/C9/O10) adopts an


Figure 1

The molecular structure of the title compound, showing the atom labelling. Displacement ellipsoids are drawn at the 50% probability levels. Only H atoms connected to O, N and chiral C atoms are shown for clarity. Other possible positions of disordered atoms have been omitted.

envelope form with puckering parameters of $Q(2) = 0.223(3) \text{ \AA}$ and $\varphi(2) = 111.7(8)^\circ$. Atom C9 deviates from the mean plane of the other four atoms by $0.357(5) \text{ \AA}$. The hy-


Figure 2

Three possible combinations of the hydroxy and amino H atoms, the probabilities being (a) 25%, (a') 25% and (b) 50%. Purple dotted lines indicate the intramolecular N—H...O and O—H...N hydrogen bonds. Other H atoms have been omitted for clarity.

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).


$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O12—H12B...N6	0.82 (3)	2.22 (4)	2.986 (4)	155 (7)
N6—H6A...O12	0.87 (3)	2.31 (5)	2.986 (4)	135 (6)
O12—H12A...O12 ⁱ	0.84 (3)	1.98 (4)	2.750 (4)	151 (5)
N6—H6C...O17B ⁱⁱ	0.81 (2)	2.57 (2)	3.369 (7)	169 (3)
C19A—H19C...N6 ⁱⁱⁱ	0.98	2.59	3.555 (15)	167
N6—H6B...O17A ^{iv}	0.85 (2)	2.61 (5)	3.286 (7)	137 (5)
C19B—H19E...O12 ^v	0.98	2.62	3.573 (11)	164

Symmetry codes: (i) $-x + 1, y, -z + 1$; (ii) $x - \frac{1}{2}, y - \frac{1}{2}, z - \frac{1}{2}$; (iii) $x + \frac{1}{2}, y + \frac{1}{2}, z + \frac{1}{2}$; (iv) $x - \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$; (v) $-x + 1, y + 1, -z + 1$.


droxy H atom has two possible positions and the amino H atoms have three possible positions, generating two types of intramolecular hydrogen bonds with an $S(7)$ graph-set motif between the hydroxy and amino groups, N6—H6A...O12 and O12—H12B...N6 (Table 1 and Fig. 2). The occupation factor of atoms H12A, H12B and H6A is 0.5, while that of atoms H6B and H6C is 0.75. The unsaturated ester substituent (C15/C14/C16/O17/O18/C19) is disordered over two orientations with refined occupancies of 0.482 (5) and 0.518 (5).

3. Supramolecular features

The crystal packing is stabilized by O—H...O hydrogen bonding (O12—H12A...O12ⁱ; Table 1), connecting molecules related by a twofold rotation axis into a dimer. As the result of this intermolecular linkage, the intramolecular hydrogen-bonding pattern is restricted, as shown in Fig. 3. The dimers are further linked by weak N—H...O, C—H...N and C—H...O interactions (N6—H6C...O17Bⁱⁱ and N6—H6B...O17A^{iv}, C19A—H19C...N6ⁱⁱⁱ and C19B—


Figure 3

A pair of molecules showing a correlation between the intra- and intermolecular hydrogen bonds. A yellow dashed line indicates the intermolecular O—H...O hydrogen bond. Purple dotted lines indicate the intramolecular N—H...O and O—H...N hydrogen bonds. Only the H atoms of the hydroxy and amino groups are shown for clarity. The other possible position of N-bound H atoms due to the disorder are omitted. [Symmetry code: (i) $-x + 1, y, -z + 1$.]



Figure 4

A packing diagram viewed down to the b axis. Yellow lines indicate the intermolecular O—H...O hydrogen bonds, generating the dimers. Black dashed lines indicate the intermolecular N—H...O, C—H...N and C—H...O interactions. Only H atoms involved in hydrogen bonds are shown for clarity. [Symmetry codes: (i) $-x + 1, y, -z + 1$; (ii) $x - \frac{1}{2}, y - \frac{1}{2}, z - \frac{1}{2}$; (iii) $x + \frac{1}{2}, y + \frac{1}{2}, z + \frac{1}{2}$; (vi) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (vii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$]


H19E...O12^v; Table 1, Figs. 4 and 5) to form a sheet structure parallel to $(\bar{1}01)$.

4. Database survey

In the Cambridge Structural Database (CSD, Version 5.36, November 2014; Groom & Allen, 2014), eight structures possessing a 1,3-dioxolane core with 4-(prop-2-enoate-3-yl) and 5-hydroxymethyl substituents, (a), are registered (Fig. 6). These include its 2,2-dimethyl, (b), 2-oxo, (c) and 2-alkoxy-2-


Figure 5

A partial packing diagram viewed along $[\bar{1}01]$, showing hydrogen bonding in the sheet structure. Overlapped molecules indicate the dimer. Black dashed lines indicate the intermolecular N—H...O, C—H...N and C—H...O interactions. Only H atoms involved in hydrogen bonds are shown for clarity.


Figure 6

The core structures for database survey; (a) 5-hydroxymethyl-4-(prop-2-enoate-3-yl) substituted 1,3-dioxolane, and its (b) 2,2-dimethyl, (c) 2-oxo, (d) 2-alkoxy-2-alkyl and (e) 2-amino-2-trichloromethyl derivatives, (f) 2,2,2-trichloroethan-1-amine, and its derivatives (g) 2-amino-2-trichloromethyl-1,3-dioxolane ($n = 1$) or -1,3-dioxane ($n = 2$); and (h) the structure of the title compound.

alkyl (orthoester) derivative, (d), but its 2-amino-2-trichloromethyl derivative, (e), which is related to the title compound, (h), has not been reported.

On the other hand, a search in CSD for a 2,2,2-trichloroethan-1-amine skeleton, (f), gave 12 entries. These include two structures (LIBHIO: Rondot *et al.*, 2007; WEKWOY: Haeckel *et al.*, 1994) with a 2-amino-2-trichloromethyl-1,3-dioxolane or -1,3-dioxane core, (g). N-bound hydrogen atoms in the structure of LIBHIO were refined as having an sp^3 configuration and tilted towards chlorine atoms, whereas those in other 11 structures were refined assuming an sp^2 configuration of the N atom.

5. Synthesis and crystallization

The title compound was derived from D-erythrose, which was prepared according to the reported procedure (Storz *et al.*, 1999) from D-glucose (Yasushima *et al.*, 2016). Purification was carried out by silica gel column chromatography, and colourless crystals were obtained from a toluene solution by slow evaporation at ambient temperature. M.p. 365–366 K. $[\alpha]_D^{20} + 33.8$ (c 0.32, CHCl_3).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. The unsaturated ester group is disordered; the atoms C15/C16/O17/O18/C19 were split into two sets of positions *A* and *B* with their geometries restrained, and the refined occupancies being 0.482 (5) and 0.518 (5), respectively.

C-bound H atoms were positioned geometrically with C—H = 0.95–1.00 Å, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$. The O-bound hydrogen atom has two possible positions. They were refined

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₁₀ H ₁₄ Cl ₃ NO ₅
<i>M_r</i>	334.57
Crystal system, space group	Monoclinic, <i>I</i> 2
Temperature (K)	90
<i>a</i> , <i>b</i> , <i>c</i> (Å)	14.8821 (9), 5.5847 (3), 17.2971 (14)
β (°)	105.847 (2)
<i>V</i> (Å ³)	1382.96 (16)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.68
Crystal size (mm)	0.26 × 0.22 × 0.16
Data collection	
Diffraction	Broker D8 Venture
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2014)
<i>T_{min}</i> , <i>T_{max}</i>	0.84, 0.90
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	11488, 2420, 2272
<i>R_{int}</i>	0.030
(sin θ/λ) _{max} (Å ⁻¹)	0.595
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.023, 0.052, 1.03
No. of reflections	2420
No. of parameters	207
No. of restraints	36
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.29, -0.21
Absolute structure	Flack <i>x</i> determined using 959 quotients [(<i>I</i> ⁺) - (<i>I</i> ⁻)] / [(<i>I</i> ⁺) + (<i>I</i> ⁻)] (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	-0.02 (2)

Computer programs: *APEX2* (Bruker, 2014), *SAINT* (Bruker, 2014), *SHELXS2013* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015), *Mercury* (Macrae *et al.*, 2006), *publCIF* (Westrip, 2010) and *PLATON* (Spek, 2009).

isotropically with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ and the O—H distances restrained. The N-bound hydrogen atoms have three possible

positions. They were refined isotropically with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ and the N—H and H···H distances restrained. The site-occupation factors of the disordered H atoms of the hydroxy group (H12*A* and H12*B*) and one of the amino group (H6*A*) were uniquely assigned to 0.5 each based on the two possible patterns of the hydrogen-bonding linkages related by the twofold axis. The occupation factors of the other N-bound H6*B* and H6*C* atoms were assumed to be 0.75 each from a difference map.

Acknowledgements

This research was partially supported by the Keio Gijuku Fukuzawa Memorial Fund for the Advancement of Education and Research. We also thank Professor S. Ohba (Keio University, Japan) for his valuable advice.

References

- Bruker (2014). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Haeckel, R., Troll, C., Fischer, H. & Schmidt, R. R. (1994). *Synlett*, pp. 84–86.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nakayama, Y., Sekiya, R., Oishi, H., Hama, N., Yamazaki, M., Sato, T. & Chida, N. (2013). *Chem. Eur. J.* **19**, 12052–12058.
- Overman, L. E. (1974). *J. Am. Chem. Soc.* **96**, 597–599.
- Overman, L. E. (1976). *J. Am. Chem. Soc.* **98**, 2901–2910.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Rondot, C., Retailleau, P. & Zhu, J. (2007). *Org. Lett.* **9**, 247–250.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Storz, T., Bernet, B. & Vasella, A. (1999). *Helv. Chim. Acta*, **82**, 2380–2412.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yasushima, D., Sato, T. & Chida, N. (2016). In preparation.

supporting information

Acta Cryst. (2016). E72, 343-346 [doi:10.1107/S2056989016002474]

Crystal structure of (+)-methyl (*E*)-3-[(2*S*,4*S*,5*R*)-2-amino-5-hydroxymethyl-2-trichloromethyl-1,3-dioxolan-4-yl]-2-methylprop-2-enoate

Takeshi Oishi, Daichi Yasushima, Kihiro Yuasa, Takaaki Sato and Noritaka Chida

Computing details

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINTE* (Bruker, 2014); data reduction: *SAINTE* (Bruker, 2014); program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *pubCIF* (Westrip, 2010) and *PLATON* (Spek, 2009).

(+)-Methyl (*E*)-3-[(2*S*,4*S*,5*R*)-2-amino-5-(hydroxymethyl)-2-trichloromethyl-1,3-dioxolan-4-yl]-2-methylprop-2-enoate

Crystal data

C₁₀H₁₄Cl₃NO₅

M_r = 334.57

Monoclinic, *I*2

a = 14.8821 (9) Å

b = 5.5847 (3) Å

c = 17.2971 (14) Å

β = 105.847 (2)°

V = 1382.96 (16) Å³

Z = 4

F(000) = 688

D_x = 1.607 Mg m⁻³

Melting point = 366–365 K

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 6489 reflections

θ = 2.5–24.9°

μ = 0.68 mm⁻¹

T = 90 K

Prism, colourless

0.26 × 0.22 × 0.16 mm

Data collection

Bruker D8 Venture
diffractometer

Radiation source: fine-focus sealed tube

Multilayered confocal mirror monochromator

Detector resolution: 10.4167 pixels mm⁻¹

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2014)

T_{min} = 0.84, *T_{max}* = 0.90

11488 measured reflections

2420 independent reflections

2272 reflections with *I* > 2σ(*I*)

R_{int} = 0.030

θ_{max} = 25.0°, θ_{min} = 2.5°

h = -17→17

k = -6→6

l = -20→20

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.023

wR(*F*²) = 0.052

S = 1.03

2420 reflections

207 parameters

36 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: mixed

H atoms treated by a mixture of independent

and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0137P)^2 + 1.951P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack x determined using
959 quotients $[(F^+)-(F^-)]/[(F^+)+(F^-)]$ (Parsons *et al.*, 2013)

Absolute structure parameter: -0.02 (2)

Special details

Experimental. M.p. 365–366 K (not corrected); $[\alpha]_{\text{D}}^{20} + 33.8$ (c 0.32, CHCl₃); IR (KBr): 3450, 3393, 3371, 3295, 3020, 2947, 2921, 2873, 1715, 1656, 1613, 1579, 1438, 1376, 1336, 1313, 1251, 1220, 1121, 1098, 1080, 1037, 1003, 977, 933, 909, 826, 803, 740, 585 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ (p.p.m.) 7.12 (dq, $J = 7.5, 1.4$ Hz, 1H; H13), 5.49 (t, $J = 7.5$ Hz, 1H; H8), 4.74 (dt, $J = 7.5, 2.3$ Hz, 1H; H9), 4.63 (bs, 1H; H12), 3.90 (dd, $J = 12.9, 2.3$ Hz, 1H; H11A), 3.77 (s, 3H; H19A–F), 3.57 (bd, $J = 12.9$ Hz, 1H; H11B), 3.06 (bs, 2H; H6), 1.90 (d, $J = 1.4$ Hz, 3H; H15A–F); ¹³C NMR (125 MHz, CDCl₃): δ (p.p.m.) 167.6 (C), 135.6 (CH), 131.7 (C), 115.3 (C), 103.8 (C), 83.6 (CH), 79.8 (CH), 62.1 (CH₂), 52.4 (CH₃), 13.5 (CH₃);

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. Problematic three reflections (1 –5 10, –13 0 15 and –12 0 16) with $|I_{\text{obs}} - I_{\text{calc}}|/\sigma W(I)$ greater than 10 have been omitted in the final refinement.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.49607 (5)	0.41064 (14)	0.10141 (4)	0.02169 (18)	
C12	0.69020 (5)	0.44496 (13)	0.18886 (5)	0.02292 (19)	
C13	0.57725 (5)	0.87077 (13)	0.15202 (5)	0.0279 (2)	
C4	0.5774 (2)	0.5655 (5)	0.17849 (19)	0.0165 (7)	
C5	0.5517 (2)	0.5446 (6)	0.25983 (19)	0.0192 (7)	
N6	0.4607 (2)	0.6381 (8)	0.25371 (19)	0.0432 (10)	
H6A	0.453 (6)	0.657 (12)	0.301 (2)	0.052*	0.5
H6B	0.453 (4)	0.781 (5)	0.236 (3)	0.052*	0.75
H6C	0.433 (2)	0.523 (6)	0.230 (3)	0.052*	0.75
O7	0.61956 (15)	0.6675 (4)	0.31855 (13)	0.0202 (5)	
C8	0.6707 (2)	0.5008 (6)	0.37781 (19)	0.0190 (7)	
H8	0.7322	0.4656	0.3673	0.023*	
C9	0.6103 (2)	0.2743 (6)	0.36247 (19)	0.0206 (7)	
H9	0.6519	0.1318	0.3659	0.025*	
O10	0.55577 (18)	0.3032 (4)	0.28050 (13)	0.0267 (6)	
C11	0.5478 (2)	0.2341 (6)	0.4164 (2)	0.0227 (7)	
H11A	0.5078	0.0928	0.3971	0.027*	
H11B	0.5867	0.1992	0.4715	0.027*	
O12	0.49006 (15)	0.4361 (5)	0.41859 (12)	0.0263 (5)	
H12A	0.496 (4)	0.488 (9)	0.4653 (18)	0.039*	0.5
H12B	0.465 (5)	0.491 (14)	0.374 (2)	0.039*	0.5
C13	0.6872 (2)	0.6143 (6)	0.45876 (19)	0.0230 (7)	

H13	0.647	0.7423	0.4636	0.028*	
C14	0.7528 (2)	0.5524 (7)	0.5242 (2)	0.0294 (9)	
C15A	0.818 (3)	0.343 (6)	0.5238 (16)	0.044 (3)	0.482 (5)
H15A	0.8202	0.311	0.4686	0.066*	0.482 (5)
H15B	0.7958	0.2007	0.5458	0.066*	0.482 (5)
H15C	0.8813	0.3833	0.5568	0.066*	0.482 (5)
C16A	0.767 (4)	0.651 (8)	0.6034 (12)	0.033 (4)	0.482 (5)
O17A	0.8138 (4)	0.5539 (12)	0.6617 (3)	0.0250 (10)	0.482 (5)
O18A	0.7178 (8)	0.851 (4)	0.5940 (10)	0.028 (2)	0.482 (5)
C19A	0.7246 (9)	0.960 (3)	0.6710 (10)	0.032 (2)	0.482 (5)
H19A	0.6983	0.8512	0.7037	0.047*	0.482 (5)
H19B	0.6899	1.1108	0.6631	0.047*	0.482 (5)
H19C	0.7904	0.9908	0.6985	0.047*	0.482 (5)
C15B	0.825 (2)	0.365 (6)	0.5320 (16)	0.044 (3)	0.518 (5)
H15D	0.8045	0.2507	0.4876	0.066*	0.518 (5)
H15E	0.8331	0.2813	0.5832	0.066*	0.518 (5)
H15F	0.8837	0.439	0.5302	0.066*	0.518 (5)
C16B	0.757 (3)	0.706 (7)	0.5989 (12)	0.033 (4)	0.518 (5)
O17B	0.8182 (4)	0.6902 (11)	0.6645 (3)	0.0250 (10)	0.518 (5)
O18B	0.6889 (7)	0.875 (3)	0.5956 (10)	0.028 (2)	0.518 (5)
C19B	0.6959 (9)	1.028 (2)	0.6652 (9)	0.032 (2)	0.518 (5)
H19D	0.6942	0.9286	0.7115	0.047*	0.518 (5)
H19E	0.6433	1.1402	0.6535	0.047*	0.518 (5)
H19F	0.7547	1.1171	0.6773	0.047*	0.518 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0302 (4)	0.0189 (4)	0.0146 (3)	-0.0018 (4)	0.0037 (3)	-0.0028 (3)
C12	0.0207 (4)	0.0251 (5)	0.0273 (4)	0.0083 (3)	0.0141 (3)	0.0056 (4)
C13	0.0237 (4)	0.0137 (4)	0.0420 (5)	0.0004 (3)	0.0018 (4)	0.0047 (4)
C4	0.0155 (16)	0.0148 (16)	0.0214 (16)	0.0008 (12)	0.0086 (13)	-0.0014 (13)
C5	0.0131 (17)	0.0287 (18)	0.0165 (17)	-0.0020 (13)	0.0051 (13)	-0.0073 (14)
N6	0.0131 (15)	0.089 (3)	0.0263 (18)	0.0115 (18)	0.0024 (13)	-0.0247 (19)
O7	0.0208 (12)	0.0169 (12)	0.0172 (11)	0.0036 (9)	-0.0047 (9)	-0.0012 (9)
C8	0.0137 (15)	0.0210 (18)	0.0223 (17)	0.0045 (13)	0.0048 (13)	0.0037 (13)
C9	0.0240 (18)	0.0197 (17)	0.0206 (16)	0.0036 (14)	0.0104 (14)	0.0002 (14)
O10	0.0369 (15)	0.0285 (14)	0.0170 (12)	-0.0178 (11)	0.0112 (11)	-0.0035 (10)
C11	0.0295 (19)	0.0227 (19)	0.0208 (17)	-0.0012 (15)	0.0150 (15)	-0.0007 (14)
O12	0.0254 (12)	0.0362 (15)	0.0209 (11)	0.0051 (12)	0.0123 (10)	-0.0023 (12)
C13	0.0171 (16)	0.0258 (19)	0.0244 (18)	-0.0025 (14)	0.0027 (14)	0.0030 (15)
C14	0.0161 (18)	0.042 (2)	0.0273 (19)	-0.0112 (15)	0.0012 (15)	0.0131 (17)
C15A	0.021 (4)	0.066 (5)	0.039 (4)	0.007 (4)	0.000 (4)	0.030 (4)
C16A	0.014 (8)	0.065 (16)	0.018 (3)	-0.015 (6)	0.000 (2)	0.014 (5)
O17A	0.0230 (16)	0.033 (3)	0.0179 (15)	-0.003 (3)	0.0029 (13)	0.002 (3)
O18A	0.022 (6)	0.039 (4)	0.0231 (14)	-0.011 (5)	0.006 (4)	-0.0089 (19)
C19A	0.031 (7)	0.034 (7)	0.031 (3)	-0.005 (4)	0.012 (5)	-0.017 (4)
C15B	0.021 (4)	0.066 (5)	0.039 (4)	0.007 (4)	0.000 (4)	0.030 (4)

C16B	0.014 (8)	0.065 (16)	0.018 (3)	-0.015 (6)	0.000 (2)	0.014 (5)
O17B	0.0230 (16)	0.033 (3)	0.0179 (15)	-0.003 (3)	0.0029 (13)	0.002 (3)
O18B	0.022 (6)	0.039 (4)	0.0231 (14)	-0.011 (5)	0.006 (4)	-0.0089 (19)
C19B	0.031 (7)	0.034 (7)	0.031 (3)	-0.005 (4)	0.012 (5)	-0.017 (4)

Geometric parameters (Å, °)

C11—C4	1.763 (3)	C13—H13	0.95
C12—C4	1.772 (3)	C14—C16A	1.439 (17)
C13—C4	1.765 (3)	C14—C15B	1.475 (17)
C4—C5	1.560 (4)	C14—C15A	1.524 (18)
C5—O10	1.392 (4)	C14—C16B	1.538 (16)
C5—O7	1.401 (4)	C15A—H15A	0.98
C5—N6	1.428 (5)	C15A—H15B	0.98
N6—H6A	0.87 (3)	C15A—H15C	0.98
N6—H6B	0.85 (2)	C16A—O17A	1.19 (2)
N6—H6C	0.81 (2)	C16A—O18A	1.318 (19)
O7—C8	1.439 (4)	O18A—C19A	1.442 (17)
C8—C13	1.495 (4)	C19A—H19A	0.98
C8—C9	1.532 (4)	C19A—H19B	0.98
C8—H8	1.0	C19A—H19C	0.98
C9—O10	1.437 (4)	C15B—H15D	0.98
C9—C11	1.502 (5)	C15B—H15E	0.98
C9—H9	1.0	C15B—H15F	0.98
C11—O12	1.425 (4)	C16B—O17B	1.252 (19)
C11—H11A	0.99	C16B—O18B	1.370 (18)
C11—H11B	0.99	O18B—C19B	1.455 (16)
O12—H12A	0.84 (3)	C19B—H19D	0.98
O12—H12B	0.82 (3)	C19B—H19E	0.98
C13—C14	1.323 (5)	C19B—H19F	0.98
C5—C4—C11	111.0 (2)	C14—C13—C8	125.8 (3)
C5—C4—C13	108.9 (2)	C14—C13—H13	117.1
C11—C4—C13	109.02 (17)	C8—C13—H13	117.1
C5—C4—C12	109.8 (2)	C13—C14—C16A	126.6 (9)
C11—C4—C12	109.01 (16)	C13—C14—C15B	127.6 (11)
C13—C4—C12	109.09 (16)	C13—C14—C15A	121.4 (11)
O10—C5—O7	108.4 (3)	C16A—C14—C15A	111.8 (12)
O10—C5—N6	110.3 (3)	C13—C14—C16B	115.0 (7)
O7—C5—N6	110.8 (3)	C15B—C14—C16B	117.3 (13)
O10—C5—C4	107.5 (3)	C14—C15A—H15A	109.5
O7—C5—C4	108.2 (3)	C14—C15A—H15B	109.5
N6—C5—C4	111.5 (3)	H15A—C15A—H15B	109.5
C5—N6—H6A	110 (5)	C14—C15A—H15C	109.5
C5—N6—H6B	113 (4)	H15A—C15A—H15C	109.5
H6A—N6—H6B	101 (5)	H15B—C15A—H15C	109.5
C5—N6—H6C	95 (2)	O17A—C16A—O18A	132.0 (16)
H6A—N6—H6C	113 (5)	O17A—C16A—C14	122.1 (15)

H6B—N6—H6C	124 (5)	O18A—C16A—C14	106.0 (14)
C5—O7—C8	109.6 (2)	C16A—O18A—C19A	110.2 (15)
O7—C8—C13	108.2 (2)	O18A—C19A—H19A	109.5
O7—C8—C9	103.9 (2)	O18A—C19A—H19B	109.5
C13—C8—C9	116.8 (3)	H19A—C19A—H19B	109.5
O7—C8—H8	109.2	O18A—C19A—H19C	109.5
C13—C8—H8	109.2	H19A—C19A—H19C	109.5
C9—C8—H8	109.2	H19B—C19A—H19C	109.5
O10—C9—C11	110.5 (3)	C14—C15B—H15D	109.5
O10—C9—C8	103.0 (2)	C14—C15B—H15E	109.5
C11—C9—C8	116.7 (3)	H15D—C15B—H15E	109.5
O10—C9—H9	108.8	C14—C15B—H15F	109.5
C11—C9—H9	108.8	H15D—C15B—H15F	109.5
C8—C9—H9	108.8	H15E—C15B—H15F	109.5
C5—O10—C9	109.6 (2)	O17B—C16B—O18B	115.7 (13)
O12—C11—C9	112.2 (3)	O17B—C16B—C14	124.9 (15)
O12—C11—H11A	109.2	O18B—C16B—C14	119.4 (15)
C9—C11—H11A	109.2	C16B—O18B—C19B	119.0 (13)
O12—C11—H11B	109.2	O18B—C19B—H19D	109.5
C9—C11—H11B	109.2	O18B—C19B—H19E	109.5
H11A—C11—H11B	107.9	H19D—C19B—H19E	109.5
C11—O12—H12A	113 (2)	O18B—C19B—H19F	109.5
C11—O12—H12B	113 (6)	H19D—C19B—H19F	109.5
H12A—O12—H12B	133 (6)	H19E—C19B—H19F	109.5
C11—C4—C5—O10	62.2 (3)	C8—C9—O10—C5	-22.5 (3)
C13—C4—C5—O10	-177.8 (2)	O10—C9—C11—O12	-64.2 (3)
C12—C4—C5—O10	-58.5 (3)	C8—C9—C11—O12	52.9 (4)
C11—C4—C5—O7	179.0 (2)	O7—C8—C13—C14	-158.9 (3)
C13—C4—C5—O7	-61.0 (3)	C9—C8—C13—C14	84.4 (4)
C12—C4—C5—O7	58.4 (3)	C8—C13—C14—C16A	-177 (4)
C11—C4—C5—N6	-58.9 (4)	C8—C13—C14—C15B	2 (2)
C13—C4—C5—N6	61.2 (3)	C8—C13—C14—C15A	-2 (2)
C12—C4—C5—N6	-179.5 (3)	C8—C13—C14—C16B	178 (3)
O10—C5—O7—C8	1.5 (4)	C13—C14—C16A—O17A	164 (4)
N6—C5—O7—C8	122.7 (3)	C15A—C14—C16A—O17A	-11 (7)
C4—C5—O7—C8	-114.7 (3)	C13—C14—C16A—O18A	-15 (6)
C5—O7—C8—C13	-139.8 (3)	C15A—C14—C16A—O18A	170 (4)
C5—O7—C8—C9	-15.0 (3)	O17A—C16A—O18A—C19A	-1 (8)
O7—C8—C9—O10	22.3 (3)	C14—C16A—O18A—C19A	178 (3)
C13—C8—C9—O10	141.3 (3)	C13—C14—C16B—O17B	-174 (4)
O7—C8—C9—C11	-98.9 (3)	C15B—C14—C16B—O17B	3 (7)
C13—C8—C9—C11	20.1 (4)	C13—C14—C16B—O18B	6 (6)
O7—C5—O10—C9	14.0 (4)	C15B—C14—C16B—O18B	-177 (4)
N6—C5—O10—C9	-107.5 (3)	O17B—C16B—O18B—C19B	3 (6)
C4—C5—O10—C9	130.7 (2)	C14—C16B—O18B—C19B	-177 (3)
C11—C9—O10—C5	102.8 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O12—H12 <i>B</i> \cdots N6	0.82 (3)	2.22 (4)	2.986 (4)	155 (7)
N6—H6 <i>A</i> \cdots O12	0.87 (3)	2.31 (5)	2.986 (4)	135 (6)
O12—H12 <i>A</i> \cdots O12 ⁱ	0.84 (3)	1.98 (4)	2.750 (4)	151 (5)
N6—H6 <i>C</i> \cdots O17 <i>B</i> ⁱⁱ	0.81 (2)	2.57 (2)	3.369 (7)	169 (3)
C19 <i>A</i> —H19 <i>C</i> \cdots N6 ⁱⁱⁱ	0.98	2.59	3.555 (15)	167
N6—H6 <i>B</i> \cdots O17 <i>A</i> ^{iv}	0.85 (2)	2.61 (5)	3.286 (7)	137 (5)
C19 <i>B</i> —H19 <i>E</i> \cdots O12 ^v	0.98	2.62	3.573 (11)	164

Symmetry codes: (i) $-x+1, y, -z+1$; (ii) $x-1/2, y-1/2, z-1/2$; (iii) $x+1/2, y+1/2, z+1/2$; (iv) $x-1/2, y+1/2, z-1/2$; (v) $-x+1, y+1, -z+1$.