

## **Crystallization and preliminary X-ray analysis of Na-ASP-1, a multi-domain pathogenesis-related-1 protein from the human hookworm parasite *Necator americanus*. Corrigendum.**

**Oluwatoyin A. Asojo,<sup>a\*</sup> Alex Loukas,<sup>b,c</sup> Mehmet Inan,<sup>d</sup> Rick Barent,<sup>d</sup> Jicai Huang,<sup>d</sup> Brad Plantz,<sup>d</sup> Amber Swanson,<sup>d</sup> Mark Gouthro,<sup>d</sup> Michael M. Meagher<sup>d</sup> and Peter J. Hotez<sup>b</sup>**

<sup>a</sup>Eppley Institute for Research in Cancer and Allied Diseases, 987696 Nebraska Medical Center, Omaha, NE 68198-7696, USA, <sup>b</sup>Department of Microbiology and Tropical Medicine, The George Washington University Medical Center, Washington DC 20037, USA, <sup>c</sup>Division of Infectious Diseases and Immunology, Queensland Institute of Medical Research, Brisbane, QLD 4006, Australia, and <sup>d</sup>Department of Chemical Engineering, The University of Nebraska–Lincoln, Lincoln, NE 68588-0643, USA

Correspondence e-mail: oasojo@unmc.edu

Data collection and instrument information reported in the article by Asojo *et al.* [(2005), *Acta Cryst.* **F61**, 391–394] are corrected.

---

In the article by Asojo *et al.* (2005), the X-ray detector and data collection were reported incorrectly for the 2.2 Å data set. The X-ray system used was actually an FRE-superbright rotating anode generator (Rigaku/MSO) operating at 45 kV and 45 mA, with Osmic Micromax optics and an R-Axis IV++ image-plate detector (Rigaku/MSO). A complete data set was collected from a single crystal using a crystal-to-detector distance of 150 mm, using the *CrystalClear (d\*TREK)* package (Pflugrath, 1999). Raw images are available on request from the authors.

### **References**

- Asojo, O. A., Loukas, A., Inan, M., Barent, R., Huang, J., Plantz, B., Swanson, A., Gouthro, M., Meagher, M. M. & Hotez, P. J. (2005). *Acta Cryst.* **F61**, 391–394.
- Pflugrath, J. W. (1999). *Acta Cryst.* **D55**, 1718–1725.