IUCrData

ISSN 2414-3146

Received 23 December 2017 Accepted 23 December 2017

Edited by W. T. A. Harrison, University of Aberdeen, Scotland

‡ Additional corresponding author, e-mail: kariukib@cardiff.ac.uk.

Keywords: crystal structure; triazole.

CCDC reference: 1813278

**Structural data**: full structural data are available from iucrdata.iucr.org

## (*E*)-3-(4-Fluorophenyl)-1-[1-(4-fluorophenyl)-5methyl-1*H*-1,2,3-triazol-4-yl]prop-2-en-1-one

Gamal A. El-Hiti,<sup>a</sup>\* Bakr F. Abdel-Wahab,<sup>b,c</sup> Mohammad Hayal Alotaibi,<sup>d</sup> Amany S. Hegazy<sup>e</sup> and Benson M. Kariuki<sup>e</sup>‡

<sup>a</sup>Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, PO Box 10219, Riyadh 11433, Saudi Arabia, <sup>b</sup>Department of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia, <sup>c</sup>Applied Organic Chemistry Department, National Research Centre, Dokki, Giza, Egypt, <sup>d</sup>National Center for Petrochemicals Technology, King Abdulaziz City for Science and Technology, PO Box 6086, Riyadh 11442, Saudi Arabia, and <sup>e</sup>School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK. \*Correspondence e-mail: gelhiti@ksu.edu.sa

The asymmetric unit of the title compound,  $C_{18}H_{13}F_2N_3O$ , comprises two molecules with similar conformations. In the crystal, weak  $C-H\cdots F$  interactions form chains of molecules and the chains are stacked to form layers parallel to (101).



Structure description

The asymmetric unit comprises two molecules of  $C_{18}H_{13}F_2N_3O$  (Fig. 1) with similar conformations (r.m.s. overlay fit = 0.180 Å). In the C1 molecule, the dihedral angles between the triazole ring and the adjacent and remote fluorobenzene rings are 39.63 (11) and 17.88 (11)°, respectively. Equivalent values for the C19 molecule are 34.67 (10) and 16.55 (11)°, respectively.

In the crystal, very weak C–H···F interactions link the molecules into chains (Table 1, Fig. 2) and the chains are stacked to form layers parallel to (101) (Fig. 3). Weak aromatic  $\pi$ - $\pi$  stacking is also observed [shortest centroid–centroid separation = 3.7332 (12) Å].

Synthesis and crystallization

1-[1-(4-Fluorophenyl)-5-methyl-1*H*-1,2,3-triazol-4-yl]ethanone, 4-fluorobenzaldehyde and sodium hydroxide (10%) were mixed in ethanol at  $20-25^{\circ}$ C for 4 h. Yellow plates (m.p. 167–168°C) were obtained following recrystallization from dimethylformamide solution of the solid obtained after work-up.





Figure 1

An ORTEP representation of the two unique molecules showing 50% probability ellipsoids.



Figure 2 Intermolecular C-H···F interactions forming chains.



Figure 3 Crystal packing showing layers parallel to (101).

#### Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2.

Table 1			
Hydrogen-bond geometry	' (Å,	°).	

$D - H \cdot \cdot \cdot A$	D-H	$H \cdot \cdot \cdot A$	$D \cdots A$	$D - H \cdots A$
$C24-H24\cdots F1^i$	0.93	2.73	3.418 (2)	132
$C35-H35\cdots F2^{ii}$	0.93	2.70	3.608 (2)	164
$C2-H2\cdots F3^{iii}$	0.93	2.65	3.395 (2)	137
$C17-H17\cdots F4^{iv}$	0.93	2.62	3.546 (2)	178

Symmetry codes: (i) x + 1, y - 1, z; (ii) x, y + 1, z + 1; (iii) x - 1, y + 1, z; (iv) x, y - 1, z - 1.

#### Table 2 Experimental details.

Crystal data Chemical formula  $M_{\rm r}$ Crystal system, space group Temperature (K) a, b, c (Å)  $\alpha, \beta, \gamma$  (°)  $V(Å^3)$ Ζ Radiation type  $\mu \text{ (mm}^{-1})$ Crystal size (mm) Data collection

Diffractometer

Absorption correction  $T_{\min}, T_{\max}$ No. of measured, independent and observed  $[I > 2\sigma(I)]$  reflections  $R_{\rm int}$  $(\sin \theta)$ Refin  $R[F^2$ 

Triclinic,  $P\overline{1}$ 296 8.1891 (4), 14.1804 (6), 14.5050 (6) 68.075 (4), 84.220 (4), 74.627 (4) 1506.65 (13) 4 Μο Κα 0.11  $0.33 \times 0.21 \times 0.10$ Rigaku Oxford Diffraction Super-Nova, Dual, Cu at zero, Atlas Gaussian (CrysAlis PRO; Rigaku OD, 2015) 0.503, 1.000 26379, 7530, 4890

 $C_{18}H_{13}F_2N_3O$ 325.31

0.029
0.700
0.052 0.141 1.05
0.053, 0.141, 1.05
7530
435
H-atom parameters constrained
0.14, -0.21

Computer programs: CrysAlis PRO (Rigaku OD, 2015), SHELXS2013 (Sheldrick, 2008), SHELXL2013 (Sheldrick, 2015), ORTEP-3 for Windows and WinGX (Farrugia, 2012) and CHEMDRAW Ultra (Cambridge Soft, 2001).

#### **Funding information**

The project was supported by King Saud University, Deanship of Scientific Research, Research Chairs.

#### References

- Cambridge Soft (2001). CHEMDRAW Ultra. Cambridge Soft Corporation, Cambridge, Massachusetts, USA.
- Farrugia, L. J. (2012). J. Appl. Cryst. 45, 849-854.
- Rigaku OD (2015). CrysAlis PRO. Rigaku Oxford Diffraction, Yarnton, England.
- Sheldrick, G. M. (2008). Acta Cryst. A64, 112-122.
- Sheldrick, G. M. (2015). Acta Cryst. C71, 3-8.

# full crystallographic data

*IUCrData* (2018). **3**, x171841 [https://doi.org/10.1107/S2414314617018417]

(*E*)-3-(4-Fluorophenyl)-1-[1-(4-fluorophenyl)-5-methyl-1*H*-1,2,3-triazol-4-yl]prop-2-en-1-one

Gamal A. El-Hiti, Bakr F. Abdel-Wahab, Mohammad Hayal Alotaibi, Amany S. Hegazy and Benson M. Kariuki

(E)-3-(4-Fluorophenyl)-1-[1-(4-fluorophenyl)-5-methyl-1H-1,2,3-triazol-4-yl]prop-2-en-1-one

#### Crystal data

 $C_{18}H_{13}F_{2}N_{3}O$   $M_{r} = 325.31$ Triclinic, *P*1 *a* = 8.1891 (4) Å *b* = 14.1804 (6) Å *c* = 14.5050 (6) Å *a* = 68.075 (4)° *β* = 84.220 (4)° *γ* = 74.627 (4)° *V* = 1506.65 (13) Å<sup>3</sup>

### Data collection

Rigaku Oxford Diffraction SuperNova, Dual, Cu at zero, Atlas diffractometer  $\omega$  scans Absorption correction: gaussian (CrysAlisPro; Rigaku OD, 2015)  $T_{\min} = 0.503, T_{\max} = 1.000$ 26379 measured reflections

### Refinement

Refinement on  $F^2$ Least-squares matrix: full  $R[F^2 > 2\sigma(F^2)] = 0.053$  $wR(F^2) = 0.141$ S = 1.057530 reflections 435 parameters 0 restraints Z = 4 F(000) = 672  $D_x = 1.434 \text{ Mg m}^{-3}$ Mo K $\alpha$  radiation,  $\lambda = 0.71073 \text{ Å}$ Cell parameters from 6895 reflections  $\theta = 4.2-27.7^{\circ}$   $\mu = 0.11 \text{ mm}^{-1}$ T = 296 K Plate, yellow  $0.33 \times 0.21 \times 0.10 \text{ mm}$ 

7530 independent reflections 4890 reflections with  $I > 2\sigma(I)$  $R_{int} = 0.029$  $\theta_{max} = 29.8^{\circ}, \ \theta_{min} = 3.0^{\circ}$  $h = -11 \rightarrow 11$  $k = -19 \rightarrow 19$  $l = -19 \rightarrow 19$ 

Primary atom site location: structure-invariant direct methods Hydrogen site location: inferred from neighbouring sites H-atom parameters constrained  $w = 1/[\sigma^2(F_o^2) + (0.0477P)^2 + 0.4741P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{max} < 0.001$  $\Delta\rho_{max} = 0.14 \text{ e } \text{Å}^{-3}$  $\Delta\rho_{min} = -0.21 \text{ e } \text{Å}^{-3}$ 

#### Special details

**Experimental**. Numerical absorption correction based on gaussian integration over a multifaceted crystal model Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm. **Geometry**. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement**. All hydrogen atoms were placed in calculated positions and refined using a riding model. Methyl C—H bonds were fixed at 0.96 Å, with displacement parameters 1.5 times  $U_{eq}(C)$ , and were allowed to spin about the C—C bond. Aromatic C—H distances were set to 0.93 Å and their U(iso) set to 1.2 times the  $U_{eq}$  for the atoms to which they are bonded.

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$
C1	0.2075 (3)	0.01266 (16)	0.03379 (16)	0.0604 (5)
C2	0.2384 (3)	0.05994 (16)	-0.06481 (16)	0.0667 (6)
H2	0.2281	0.1319	-0.0926	0.080*
C3	0.2854 (3)	-0.00171 (14)	-0.12258 (14)	0.0574 (5)
Н3	0.3067	0.0286	-0.1901	0.069*
C4	0.3003 (2)	-0.10814 (13)	-0.07944 (13)	0.0451 (4)
C5	0.2665 (3)	-0.15357 (15)	0.02006 (14)	0.0589 (5)
Н5	0.2756	-0.2254	0.0484	0.071*
C6	0.2191 (3)	-0.09221 (17)	0.07755 (15)	0.0645 (6)
H6	0.1956	-0.1218	0.1448	0.077*
C7	0.5744 (2)	-0.10855 (15)	-0.24935 (15)	0.0554 (5)
H7A	0.5199	-0.0457	-0.3017	0.083*
H7B	0.6744	-0.1441	-0.2748	0.083*
H7C	0.6049	-0.0907	-0.1968	0.083*
C8	0.4564 (2)	-0.17840 (12)	-0.21012 (12)	0.0407 (4)
С9	0.4406 (2)	-0.26215 (13)	-0.23169 (13)	0.0434 (4)
C10	0.5329 (2)	-0.30726 (13)	-0.30344 (13)	0.0459 (4)
C11	0.5127 (2)	-0.41066 (14)	-0.29353 (14)	0.0484 (4)
H11	0.4446	-0.4428	-0.2426	0.058*
C12	0.5872 (2)	-0.45986 (13)	-0.35404 (13)	0.0466 (4)
H12	0.6513	-0.4247	-0.4056	0.056*
C13	0.5791 (2)	-0.56363 (13)	-0.34807 (13)	0.0452 (4)
C14	0.4970 (3)	-0.62760 (15)	-0.27110 (14)	0.0562 (5)
H14	0.4438	-0.6037	-0.2214	0.067*
C15	0.4930 (3)	-0.72559 (16)	-0.26704 (16)	0.0636 (5)
H15	0.4386	-0.7682	-0.2152	0.076*
C16	0.5715 (3)	-0.75854 (15)	-0.34159 (17)	0.0614 (5)
C17	0.6531 (3)	-0.69911 (16)	-0.41866 (16)	0.0638 (5)
H17	0.7049	-0.7236	-0.4682	0.077*
C18	0.6566 (3)	-0.60140 (15)	-0.42123 (14)	0.0549 (5)
H18	0.7123	-0.5599	-0.4733	0.066*
C19	1.2425 (2)	-0.66173 (15)	0.01546 (14)	0.0517 (4)
C20	1.3283 (3)	-0.59030 (17)	0.01026 (15)	0.0616 (5)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters  $(Å^2)$ 

<b>U2</b> 0	1 4220	_0.5934	-0.0206	0.074*
C21	1.4229	-0.52812(16)	-0.0300 0.06637 (14)	$0.074^{\circ}$
U21	1.2730 (5)	-0.4702	0.06037 (14)	0.0388 (3)
C22	1.3317	-0.53858(13)	0.0041 0.12588 (12)	0.070
C22	1.1323(2) 1.0451(2)	-0.61035(13)	0.12388(12) 0.12862(13)	0.0414(4)
U23	0.0485	-0.6164	0.12802 (13)	0.0470 (4)
C24	1,1016(2)	-0.67331(15)	0.1078 0.07208 (14)	$0.050^{\circ}$
U24	1.1010 (2)	-0.7226	0.07238 (14)	0.0554 (5)
C25	0.9663 (3)	-0.58167(13)	0.0747 0.34081 (14)	$0.004^{\circ}$
U25	0.9003 (5)	-0.5847	0.34001 (14)	0.0347(3)
1125A 1125D	0.0730	0.3847	0.3230	0.082*
П256	1.0483	-0.5840	0.4070	0.082*
П23C	1.0403	-0.0400	0.3310 0.26076(12)	$0.082^{\circ}$
C20	1.0010(2)	-0.48221(13) -0.38655(13)	0.20970(12) 0.27027(12)	0.0413(4)
C27	0.9802(2)	-0.38033(13)	0.27927(13) 0.26008(12)	0.0403(4)
C28	0.9078(2)	-0.35110(13)	0.30098(13) 0.24065(14)	0.0478(4)
C29	0.9169 (3)	-0.24485 (14)	0.34965 (14)	0.0526 (4)
H29	0.9774	-0.2092	0.2902	0.003*
C30	0.8435 (2)	-0.19/92 (14)	0.411/0 (13)	0.0494 (4)
H30	0.7826	-0.2348	0.4641	0.059*
C31	0.8486 (2)	-0.09345 (13)	0.40616 (13)	0.0468 (4)
C32	0.9495 (3)	-0.03399 (15)	0.33795 (15)	0.0558 (5)
H32	1.0171	-0.0611	0.2938	0.067*
C33	0.9505 (3)	0.06391 (16)	0.33501 (16)	0.0609 (5)
H33	1.0179	0.1031	0.2895	0.073*
C34	0.8501 (3)	0.10242 (14)	0.40055 (16)	0.0572 (5)
C35	0.7489 (3)	0.04812 (15)	0.46811 (16)	0.0614 (5)
H35	0.6812	0.0765	0.5114	0.074*
C36	0.7492 (3)	-0.05020 (14)	0.47070 (14)	0.0556 (5)
H36	0.6811	-0.0884	0.5168	0.067*
N1	0.34173 (18)	-0.17422 (11)	-0.13693 (11)	0.0452 (3)
N2	0.2574 (2)	-0.25153 (13)	-0.11516 (13)	0.0586 (4)
N3	0.3186 (2)	-0.30441 (12)	-0.17205 (12)	0.0549 (4)
N4	1.07488 (19)	-0.46919 (11)	0.17951 (10)	0.0453 (3)
N5	1.0991 (2)	-0.36974 (12)	0.13550 (11)	0.0628 (5)
N6	1.0419 (2)	-0.32099 (12)	0.19610 (12)	0.0606 (4)
01	0.62385 (19)	-0.26083 (11)	-0.36626 (10)	0.0658 (4)
02	0.84447 (19)	-0.40717 (10)	0.43345 (10)	0.0636 (4)
F1	0.1592 (2)	0.07284 (11)	0.09065 (10)	0.0896 (4)
F2	0.5683 (2)	-0.85579 (10)	-0.33684 (12)	0.0940 (5)
F3	1.29847 (17)	-0.72282 (11)	-0.04006 (10)	0.0808 (4)
F4	0.85150 (19)	0.19891 (9)	0.39781 (11)	0.0828 (4)

Atomic displacement parameters  $(Å^2)$ 

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0705 (13)	0.0620 (12)	0.0615 (12)	-0.0169 (10)	0.0053 (10)	-0.0377 (11)
C2	0.0948 (16)	0.0473 (11)	0.0667 (13)	-0.0239 (11)	0.0071 (12)	-0.0275 (10)
C3	0.0817 (14)	0.0459 (10)	0.0473 (10)	-0.0225 (10)	0.0058 (10)	-0.0167 (9)

C4	0.0502 (10)	0.0429 (9)	0.0470 (10)	-0.0160 (7)	0.0068 (8)	-0.0202 (8)
C5	0.0767 (14)	0.0467 (10)	0.0513 (11)	-0.0178 (9)	0.0158 (10)	-0.0177 (9)
C6	0.0807 (15)	0.0627 (13)	0.0494 (11)	-0.0168 (11)	0.0130 (10)	-0.0232 (10)
C7	0.0626 (12)	0.0536 (11)	0.0584 (11)	-0.0315 (9)	0.0137 (9)	-0.0214 (9)
C8	0.0446 (9)	0.0378 (8)	0.0396 (8)	-0.0150 (7)	0.0058 (7)	-0.0119 (7)
C9	0.0486 (10)	0.0383 (9)	0.0441 (9)	-0.0165 (7)	0.0061 (7)	-0.0135 (7)
C10	0.0508 (10)	0.0415 (9)	0.0459 (9)	-0.0147 (8)	0.0056 (8)	-0.0154 (8)
C11	0.0548 (10)	0.0448 (10)	0.0503 (10)	-0.0186 (8)	0.0080 (8)	-0.0201 (8)
C12	0.0534 (10)	0.0427 (9)	0.0441 (9)	-0.0136 (8)	0.0024 (8)	-0.0154 (8)
C13	0.0494 (10)	0.0434 (9)	0.0434 (9)	-0.0112 (7)	0.0010 (8)	-0.0167 (8)
C14	0.0694 (13)	0.0512 (11)	0.0520 (11)	-0.0205 (9)	0.0122 (9)	-0.0222 (9)
C15	0.0754 (14)	0.0521 (11)	0.0656 (13)	-0.0279 (10)	0.0132 (11)	-0.0191 (10)
C16	0.0702 (13)	0.0431 (10)	0.0762 (14)	-0.0150 (9)	-0.0031 (11)	-0.0260 (10)
C17	0.0777 (14)	0.0532 (11)	0.0676 (13)	-0.0141 (10)	0.0098 (11)	-0.0339 (11)
C18	0.0654 (12)	0.0490 (10)	0.0509 (10)	-0.0154 (9)	0.0099 (9)	-0.0205 (9)
C19	0.0542 (11)	0.0545 (11)	0.0509 (10)	-0.0130 (9)	0.0063 (9)	-0.0260 (9)
C20	0.0556 (11)	0.0835 (15)	0.0582 (12)	-0.0335 (11)	0.0210 (9)	-0.0338 (11)
C21	0.0649 (12)	0.0717 (13)	0.0562 (11)	-0.0410 (10)	0.0155 (10)	-0.0293 (10)
C22	0.0491 (9)	0.0405 (9)	0.0368 (8)	-0.0168 (7)	0.0013 (7)	-0.0127 (7)
C23	0.0494 (10)	0.0464 (10)	0.0497 (10)	-0.0193 (8)	0.0099 (8)	-0.0197 (8)
C24	0.0593 (11)	0.0508 (10)	0.0608 (11)	-0.0248 (9)	0.0106 (9)	-0.0272 (9)
C25	0.0724 (13)	0.0399 (9)	0.0494 (10)	-0.0208 (9)	0.0156 (9)	-0.0126 (8)
C26	0.0470 (9)	0.0380 (9)	0.0384 (8)	-0.0143 (7)	0.0024 (7)	-0.0112 (7)
C27	0.0602 (11)	0.0378 (9)	0.0424 (9)	-0.0185 (8)	0.0013 (8)	-0.0121 (7)
C28	0.0579 (11)	0.0399 (9)	0.0460 (10)	-0.0137 (8)	0.0012 (8)	-0.0151 (8)
C29	0.0681 (12)	0.0442 (10)	0.0484 (10)	-0.0203 (9)	0.0048 (9)	-0.0167 (8)
C30	0.0587 (11)	0.0440 (10)	0.0459 (10)	-0.0146 (8)	0.0020 (8)	-0.0159 (8)
C31	0.0558 (10)	0.0409 (9)	0.0444 (9)	-0.0111 (8)	-0.0017 (8)	-0.0163 (8)
C32	0.0642 (12)	0.0547 (11)	0.0561 (11)	-0.0203 (9)	0.0095 (9)	-0.0271 (9)
C33	0.0696 (13)	0.0536 (11)	0.0655 (13)	-0.0276 (10)	0.0049 (10)	-0.0211 (10)
C34	0.0723 (13)	0.0405 (10)	0.0627 (12)	-0.0126 (9)	-0.0131 (10)	-0.0205 (9)
C35	0.0789 (14)	0.0496 (11)	0.0587 (12)	-0.0101 (10)	0.0024 (10)	-0.0275 (10)
C36	0.0706 (13)	0.0474 (10)	0.0488 (10)	-0.0163 (9)	0.0077 (9)	-0.0184 (9)
N1	0.0529 (8)	0.0399 (7)	0.0481 (8)	-0.0206 (6)	0.0114 (7)	-0.0185 (7)
N2	0.0677 (10)	0.0558 (9)	0.0683 (10)	-0.0358 (8)	0.0276 (8)	-0.0333 (9)
N3	0.0636 (10)	0.0495 (9)	0.0645 (10)	-0.0296 (7)	0.0232 (8)	-0.0301 (8)
N4	0.0620 (9)	0.0388 (7)	0.0398 (7)	-0.0238 (7)	0.0061 (7)	-0.0132 (6)
N5	0.1073 (14)	0.0463 (9)	0.0442 (9)	-0.0409 (9)	0.0165 (9)	-0.0161 (7)
N6	0.0997 (13)	0.0444 (9)	0.0451 (9)	-0.0340 (9)	0.0108 (9)	-0.0162 (7)
01	0.0836 (10)	0.0560 (8)	0.0648 (9)	-0.0332 (7)	0.0338 (8)	-0.0273 (7)
O2	0.0879 (10)	0.0489 (8)	0.0548 (8)	-0.0250 (7)	0.0209 (7)	-0.0193 (7)
F1	0.1226 (12)	0.0847 (9)	0.0852 (9)	-0.0274 (8)	0.0181 (8)	-0.0605 (8)
F2	0.1197 (12)	0.0538 (7)	0.1244 (12)	-0.0332 (8)	0.0155 (9)	-0.0458 (8)
F3	0.0878 (9)	0.0880 (9)	0.0915 (9)	-0.0308 (7)	0.0328 (7)	-0.0623 (8)
F4	0.1129 (11)	0.0499 (7)	0.0981 (10)	-0.0264 (7)	-0.0075 (8)	-0.0345 (7)

Geometric parameters (Å, °)

C1—F1	1.359 (2)	C19—C24	1.363 (3)
C1—C6	1.362 (3)	C20—C21	1.376 (3)
C1—C2	1.364 (3)	C20—H20	0.9300
C2—C3	1.385 (3)	C21—C22	1.378 (2)
C2—H2	0.9300	C21—H21	0.9300
C3—C4	1.377 (2)	C22—C23	1.377 (2)
С3—Н3	0.9300	C22—N4	1.431 (2)
C4—C5	1.376 (3)	C23—C24	1.381 (2)
C4—N1	1.430 (2)	C23—H23	0.9300
C5—C6	1.379 (3)	C24—H24	0.9300
С5—Н5	0.9300	C25—C26	1.485 (2)
С6—Н6	0.9300	C25—H25A	0.9600
C7—C8	1.488 (2)	C25—H25B	0.9600
С7—Н7А	0.9600	C25—H25C	0.9600
С7—Н7В	0.9600	C26—N4	1.352 (2)
С7—Н7С	0.9600	C26—C27	1.378 (2)
C8—N1	1.353 (2)	C27—N6	1.365 (2)
C8—C9	1.375 (2)	C27—C28	1.467 (2)
C9—N3	1.365 (2)	C28—O2	1.220 (2)
C9—C10	1.465 (2)	C28—C29	1.476 (2)
C10—O1	1.222 (2)	C29—C30	1.319 (2)
C10—C11	1.471 (2)	С29—Н29	0.9300
C11—C12	1.324 (2)	C30—C31	1.465 (2)
C11—H11	0.9300	С30—Н30	0.9300
C12—C13	1.460 (2)	C31—C36	1.388 (2)
C12—H12	0.9300	C31—C32	1.398 (3)
C13—C18	1.387 (2)	C32—C33	1.375 (3)
C13—C14	1.392 (2)	С32—Н32	0.9300
C14—C15	1.378 (3)	C33—C34	1.367 (3)
C14—H14	0.9300	С33—Н33	0.9300
C15—C16	1.371 (3)	C34—F4	1.357 (2)
C15—H15	0.9300	C34—C35	1.361 (3)
C16—C17	1.360 (3)	C35—C36	1.380 (3)
C16—F2	1.362 (2)	С35—Н35	0.9300
C17—C18	1.380 (3)	С36—Н36	0.9300
С17—Н17	0.9300	N1—N2	1.3701 (19)
C18—H18	0.9300	N2—N3	1.294 (2)
C19—C20	1.355 (3)	N4—N5	1.3735 (19)
C19—F3	1.358 (2)	N5—N6	1.292 (2)
F1—C1—C6	118.35 (19)	C21—C20—H20	120.5
F1—C1—C2	118.77 (19)	C20—C21—C22	119.83 (17)
C6—C1—C2	122.86 (18)	C20—C21—H21	120.1
C1—C2—C3	118.45 (19)	C22—C21—H21	120.1
С1—С2—Н2	120.8	C23—C22—C21	120.20 (16)
С3—С2—Н2	120.8	C23—C22—N4	121.59 (15)

C4—C3—C2	119.59 (18)	C21—C22—N4	118.11 (15)
С4—С3—Н3	120.2	C22—C23—C24	119.75 (16)
С2—С3—Н3	120.2	С22—С23—Н23	120.1
C5—C4—C3	120.70 (17)	С24—С23—Н23	120.1
C5—C4—N1	117.69 (15)	C19—C24—C23	118.68 (17)
C3—C4—N1	121.52 (16)	C19—C24—H24	120.7
C4—C5—C6	119.75 (18)	C23—C24—H24	120.7
C4—C5—H5	120.1	C26—C25—H25A	109.5
C6—C5—H5	120.1	C26—C25—H25B	109.5
C1—C6—C5	118.65 (19)	H25A—C25—H25B	109.5
C1—C6—H6	120.7	C26—C25—H25C	109.5
C5—C6—H6	120.7	$H_{25A} - C_{25} - H_{25C}$	109.5
C8—C7—H7A	109.5	$H_{25B}$ $C_{25}$ $H_{25C}$	109.5
C8-C7-H7B	109.5	N4—C26—C27	103.98 (14)
H7A - C7 - H7B	109.5	N4—C26—C25	125.31(15)
C8 - C7 - H7C	109.5	$C_{27}$ $C_{26}$ $C_{25}$	120.01(10) 130.60(16)
H7A - C7 - H7C	109.5	N6-C27-C26	108.92(15)
H7B C7 H7C	109.5	N6 C27 C28	100.92(15) 120.75(15)
$\frac{11}{D} - \frac{1}{C} - \frac{11}{C}$	109.5 103 00 (14)	$C_{26} C_{27} C_{28}$	120.75(15) 130.32(15)
N1_C8_C7	103.99(14) 124.74(15)	$C_{20} = C_{27} = C_{28}$	130.32(15)
$\frac{1}{10000000000000000000000000000000000$	124.74(15) 131.10(15)	02 - 028 - 027	121.33(13) 122.77(17)
$C_{2} = C_{3} = C_{1}$	131.19(15) 100.02(15)	$C_{2} = C_{2} = C_{2}$	122.77(17)
$N_3 = C_9 = C_8$	109.02(13) 120.88(15)	$C_{2} = C_{2} = C_{2}$	113.00(13) 123.18(17)
$N_3 = C_9 = C_{10}$	120.88(13) 120.00(15)	$C_{20}$ $C_{29}$ $C_{28}$ $C_{20}$ $C$	123.18 (17)
	130.09 (15)	C30—C29—H29	118.4
01 - 010 - 011	120.54 (15)	C28—C29—H29	118.4
	122.96 (16)	$C_{29} = C_{30} = C_{31}$	126.38 (17)
C9—C10—C11	116.48 (15)	C29—C30—H30	116.8
C12—C11—C10	122.81 (16)	С31—С30—Н30	116.8
C12—C11—H11	118.6	C36—C31—C32	117.71 (16)
C10—C11—H11	118.6	C36—C31—C30	119.36 (16)
C11—C12—C13	126.70 (17)	C32—C31—C30	122.93 (16)
C11—C12—H12	116.7	C33—C32—C31	121.17 (18)
C13—C12—H12	116.7	С33—С32—Н32	119.4
C18—C13—C14	117.89 (16)	С31—С32—Н32	119.4
C18—C13—C12	119.41 (16)	C34—C33—C32	118.52 (19)
C14—C13—C12	122.69 (16)	С34—С33—Н33	120.7
C15—C14—C13	121.32 (18)	С32—С33—Н33	120.7
C15—C14—H14	119.3	F4—C34—C35	118.76 (19)
C13—C14—H14	119.3	F4—C34—C33	118.50 (19)
C16—C15—C14	118.13 (19)	C35—C34—C33	122.74 (17)
C16—C15—H15	120.9	C34—C35—C36	118.30 (19)
C14—C15—H15	120.9	С34—С35—Н35	120.8
C17—C16—F2	119.13 (19)	С36—С35—Н35	120.8
C17—C16—C15	122.97 (18)	C35—C36—C31	121.56 (18)
F2-C16-C15	117.90 (19)	С35—С36—Н36	119.2
C16—C17—C18	118.12 (19)	С31—С36—Н36	119.2
С16—С17—Н17	120.9	C8—N1—N2	110.51 (13)
C18—C17—H17	120.9	C8—N1—C4	132.05 (14)

# data reports

C17—C18—C13	121.57 (18)	N2—N1—C4	117.40 (13)
C17—C18—H18	119.2	N3—N2—N1	107.40 (13)
C13—C18—H18	119.2	N2—N3—C9	109.07 (14)
C20—C19—F3	118.33 (16)	C26—N4—N5	110.53 (14)
C20—C19—C24	122.52 (17)	C26—N4—C22	132.38 (14)
F3—C19—C24	119.15 (17)	N5—N4—C22	117.07 (13)
C19—C20—C21	119.01 (17)	N6—N5—N4	107.30 (13)
C19—C20—C21	119.01 (17)	N6—N5—N4	107.30 (13)
C19—C20—H20	120.5	N5—N6—C27	109.27 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D····A	D—H···A
$C24$ — $H24$ ···· $F1^i$	0.93	2.73	3.418 (2)	132
C35—H35…F2 <sup>ii</sup>	0.93	2.70	3.608 (2)	164
C2—H2···F3 <sup>iii</sup>	0.93	2.65	3.395 (2)	137
C17—H17…F4 <sup>iv</sup>	0.93	2.62	3.546 (2)	178

Symmetry codes: (i) *x*+1, *y*-1, *z*; (ii) *x*, *y*+1, *z*+1; (iii) *x*-1, *y*+1, *z*; (iv) *x*, *y*-1, *z*-1.