

lytic-like protease. In contrast, the high-resolution crystal structures of another prokaryote serpin, tengpin, reveals that the serpin domain of this molecule folds spontaneously and rapidly to most stable (i.e. relaxed) conformation. This is an exciting result, since tengpin represents the first serpin identified to date that obeys Anfinsen's conjecture. Furthermore, the X-ray crystal structures of tengpin reveals the structural basis for a novel mechanism for loop-C-sheet serpin-polymerisation. Analysis of the structural data provides striking insight into the mechanism of serpin metastability and the structural basis for serpin polymerisation.

[1] a) Irving J.A., et al., *Structure* 2003; b) Fulton K.F., et al., *J Biol Chem*, 2005.

**Keywords:** serpin, folding, polymerisation

### MS03 CHIRAL AND NON-CENTROSYMMETRIC STRUCTURES

**Chairpersons:** Shiv P. Halasyamani, Reiko Kuroda


#### MS03.24.1

*Acta Cryst.* (2005). A61, C11

#### Spontaneous Resolutions in Halogen Bonded Fluorinated Networks

Pierangelo Metrangolo<sup>a</sup>, Hannes Neukirch<sup>a</sup>, Tullio Pilati<sup>b</sup>, Giuseppe Resnati<sup>a</sup>, <sup>a</sup>*Department of Chemistry, Materials, and Chemical Engineering "G. Natta"; Polytechnic of Milan.* <sup>b</sup>*ISTM-C.N.R., University of Milan; Italy.* E-mail: hannes.neukirch@chem.polimi.it

Halogen bonding is an efficient tool for self-assembling halo-perfluorocarbons (PFC) and hydrocarbons (HC) [1]. Its particular ability to control spontaneous resolution in hybrid PFC-HC systems has been discovered only recently [2]. Up to now we observed spontaneous resolutions in four cases affording chiral cocrystals, space group  $P2_12_12_1$ . Three of them involved long-chain iodo-PFC's ( $C_8-C_{10}$ ) with either QUATS or  $N,N,N',N'$ -tetramethyl-*p*-phenyldiamine as bases. Their different features with regard to the segregation behaviour and the conformation of the PFC chains will be outlined. The X-ray structure of a chiral alkali halide complex **1** (Figure) involving a tripodand will also be presented.



[1] Metrangolo P., Neukirch H., Pilati T., Resnati G., *Acc. Chem. Res.*, 2005, *in press*. [2] Neukirch H., Guido E., Liantonio R., Metrangolo P., Pilati T., Resnati G., *Chem. Commun.*, 2005, 1534.

**Keywords:** halogen bonding, chiral resolution, molecular cocrystals

#### MS03.24.2

*Acta Cryst.* (2005). A61, C11

#### Invariom Modeling for Improving Absolute Structure Determination

Marianna Strumpel<sup>a</sup>, Martina Schaefer<sup>b</sup>, Tibor Koritsanszky<sup>c</sup>, Birger Dittrich<sup>d</sup>, <sup>a</sup>*Institut für Chemie/Kristallographie, FU Berlin, 14195 Berlin, Germany.* <sup>b</sup>*Schering Research Laboratory, Structural Biology, Müllerstrasse 178, 13342 Berlin, Germany.* <sup>c</sup>*Department of Chemistry, MTSU Box 0395, Murfreesboro, TN, USA* <sup>d</sup>*Chemistry, M313, University of Western Australia, WA 6009 Nedlands, Australia.* E-mail: strumpel@chemie.fu-berlin.de

A reliable determination of the Flack parameter [1] for structures of organic molecules, containing only the elements H, C, N, and O usually fails. The reason for this is the very weak anomalous signal obtained from the light atoms [2]. Recently we have introduced invarioms [3] and here we try to improve the absolute structure determination by replacing the independent atom model with the aspherical invariom scattering model. The determination of the Flack parameter was included in the program XDLSM [4]. Alternatively, its calculation has been attempted via a hole-in-one procedure. A precise

data set on a steroid compound was collected using copper radiation and CCD detection, and first results are reported.

[1] Flack H. D., *Acta Cryst.*, 1983, A39, 876. [2] Flack H. D., Bernardinelli G., *J. Appl. Cryst.*, 2000, 33, 1143. [3] Dittrich B., Koritsanszky T., Luger P., *Angew. Chem. Int. Ed.*, 2004, 43, 2718. [4] Koritsanszky T., Richter T., Macci P., Gatti C., Howard S., Mallinson P.R., Farrugia L., Su Z.W., Hansen N.K., *XD*, Freie Universität Berlin, Berlin, 2003.

**Keywords:** Invarioms, Flack parameter refinement, chiral structures

#### MS03.24.3

*Acta Cryst.* (2005). A61, C11

#### Optical Topographies of Chiral Structures

Werner Kaminsky, Bart Kahr, *University of Washington, Department of Chemistry Seattle, USA.* E-mail: wernerka@u.washington.edu

Can optical rotatory power, a phenomenon typically associated with chirality or handedness, be used as a contrast mechanism in microscopy? Chiroptical imaging techniques have not heretofore been implemented. This neglect has created a hole in the science of molecular chirality, particularly with respect to complex, heterogeneous, organized media. We built a circular extinction imaging microscope to examine chromophores in anisotropic hosts.




Figure 1. Circular Dichroism in 1-8-Dihydroxyanthraquinone.

With this instrument, images of crystals were made via two mechanisms, intrinsic circular dichroism (CD) and a new effect that was discovered and called anomalous circular extinction (ACE). Through these new chirality "spectacles" we have observed left and right handed twinning in crystals of a dye that was masked by all previous methods of analysis, Figure 1 [1]. However, when turned onto unusual dyed crystals, we observed optical effects that mimic those due to chirality.

[1] Claborn K., Puklin-Faucher E., Kurimoto M., Kaminsky W., Kahr B., *J. Am. Chem. Soc.*, 2003, 125, 14825-14831.

**Keywords:** chiroptical properties, circular dichroism measurement, dyes

#### MS03.24.4

*Acta Cryst.* (2005). A61, C11

#### Nonlinear Optical Properties of Chiral Polymers and Systems

André Persoons, *Department of Chemistry, University of Leuven, Belgium and Optical Sciences Center, University of Arizona, Tucson, AZ 85721 USA.* E-mail: andre.persoons@fys.kuleuven.ac.be

We present the nonlinear optical properties of different thin film films of chiral (conjugated) polymers. These systems exhibit large magnetic dipole nonlinearities, in some cases larger than the effects linked to electric dipole interactions. The nonlinear optical effects observed indicate the links between magnetic hypersusceptibilities and chirality. We also investigated supramolecular assemblies of helicenes where the nonlinear optical effects are exclusively described by electric dipole interactions. In the crystalline liquid state the chirality, as expressed by nonlinear CD effects, of these helicene assemblies could be switched by the application of an electric field.

**Keywords:** polymers, chirality, nonlinear optics

#### MS03.24.5

*Acta Cryst.* (2005). A61, C11-C12

#### Some Reminiscences of Non-centrosymmetric Structures

Joel Bernstein, *Department of Chemistry, Ben-Gurion University Beer Sheva, Israel 84105.* E-mail: joel@bgumail.bgu.ac.il

In the "neanderthal" age of crystallography, a light atom non-centrosymmetric crystal was usually relegated to the skeleton collection of unsolvable structures. The development of MULTAN