

jebe@crc.dk

Fatty acid breakdown is an essential process in all organisms, and the complex pathway require several sets of enzymes working in different cellular compartments. β -oxidation cycles are the center of the fatty acid breakdown, in each round the lipid chain shortening by a C₂ unit. The acyl-CoA oxidase family catalyzes the first and rate limiting step in the peroxisomal β -oxidation cycle, where acyl-CoA is converted to *trans*-2-enoyl-CoA. *Arabidopsis thaliana* have 4 different acyl-CoA oxidases, each with different chain length specificities. The structure of acyl-CoA oxidase 1 (ACX1), specific for long chain lipids, has previously been determined in our group.

We here present ACX1 in complex with acetoacetyl-CoA; the first structure of an acyl-CoA oxidase in complex with a substrate analogue. The fatty acyl moiety could be modeled between the isoalloxazine ring of FAD and the putative catalytic residue Glu424, forming hydrogen bonds to the backbone nitrogen of Glu424 and to N5 of FAD. Glu424 has moved to better accommodate the inhibitor, and is also more ordered in the complex structure. This confirms Glu424 to be the catalytic residue. The beginning of helix G and the end of helix H has moved slightly, and an interesting rotamer change of His374 and Tyr278 can be observed, bringing both residues closer to the CoA tail of the modeled ligand.

Keywords: acylCoA oxidase, beta oxidation, peroxisome

P.04.02.99

Acta Cryst. (2005). A61, C203

Structural Studies on Sulfurtransferase/Phosphatase Enzymes

Domenico Bordo^a, Andrea Spallarossa^{a,b}, Silvia Pagani^c, Timoty Larson^d, Martino Bolognesi^b, ^aNational Cancer Research Institute, Genova. ^bUniversity of Genova. ^cUniversity of Milano. ^dVirginia University. E-mail: bordo@fisica.unige.it

Sulfurtransferases are widespread enzymes that *in vitro* catalyze the transfer of a sulfur atom from a donor molecule to cyanide [1]. In order to elucidate the molecular basis of sulfur transfer reaction and identify the structural determinants for enzyme selectivity, crystallographic analyses were carried out on three different sulfurtransferases: *Azotobacter vinelandii* rhodanese (RhDA) and *Escherichia coli* GlpE and *Escherichia coli* SseA, a MST enzyme.

The crystal structure of the RhDA has been determined with the method of Multiple Isomorphous Replacement and refined at 1.8 Å resolution in the sulfur-free and persulfide-containing forms [2]. GlpE crystal structure has been determined at 1.06 Å and displays a three-dimensional fold similar to that of either RhDA domains [3]. Notably, GlpE is also structurally similar to the catalytic domain of the human cell cycle-control Cdc25 phosphatase. The distinct substrate specificity, sulphur for rhodanese enzymes and phosphate for Cdc25 phosphatases, appears to be primarily consequence of the different active site loop length in the two enzymes. These structural findings provide guidelines for the identification of the as yet unknown biological role of this protein. Also the crystal structure of SseA, solved at 2.8 Å resolution by molecular replacement method [4].

[1] Bordo D., Bork P., *EMBO Reports*, 2002, **3**, 741. [2] Bordo D. *et al.*, *J. Mol. Biol.*, 2000, **398**, 691. [3] Spallarossa A., *et al.*, *Structure*, 2001, **9**, 1117. [4] Spallarossa A., *et al.*, *J. Mol. Biol.*, 2004, **335**, 583.

Keywords: sulfurtransferases, phosphatases, rhodanese

P.04.02.100

Acta Cryst. (2005). A61, C203

Regulation of *Sulfolobus solfataricus* Uracil Phosphoribosyltransferase

Anders Kadziola^a, Stig Christoffersen^b, Eva Johansson^a, Susan Arent^a, Sine Larsen^a, Kaj Frank Jensen^b, ^aDepartment of Chemistry and ^bInstitute of Molecular Biology and Physiology, University of Copenhagen, Denmark. E-mail: anders@ccs.ki.ku.dk

UPRTase is a salvage enzyme that catalyzes the formation of UMP from PRPP (5-phosphoribosyl-1- α -diphosphate) and uracil.

CTP and UMP can independently bind to SsUPRTase and simultaneous binding of CTP and UMP strongly inhibits the enzyme. A structure to 2.8 Å resolution of SsUPRTase-CTP-UMP has already

been determined [1].

GTP causes a 20-fold increase in the turnover number k_{cat} and raises K_M for PRPP and uracil by 2- and >10-fold, respectively [2]. In order to make an SsUPRTase-GTP-PRPP complex, the enzyme must be depleted for the product UMP. Co-purified UMP binds so strongly to SsUPRTase that an unfolding and refolding procedure was necessary to remove it. UMP-depleted SsUPRTase (5 mg/mL) with 5 mM GTP and 5 mM PRPP was crystallized by vapour diffusion with PEG8000 at pH 6.5. Synchrotron data to 2.8 Å resolution has been recorded (P6₄22, a=b=122.2 Å, c=62.2 Å) and the structure determined by molecular replacement.

SsUPRTase in solution as well as in crystals is tetrameric with 222 symmetry. The allosteric binding sites for CTP/GTP are situated in the middle of the tetramer ca. 24 Å from the active sites. Transformation from inhibited to activated structure involves structural changes in the quaternary structure along with major active site movements.

[1] Arent S., Harris P., Jensen K.F., Larsen S., *Biochemistry*, 2005, **44**, 883. [2] Jensen K.F., Arent S., Larsen S., Schack L., *FEBS Journal*, 2005, **272**, 1440.

Keywords: nucleotide metabolism, protein regulation, enzyme catalysis

P.04.02.101

Acta Cryst. (2005). A61, C203

Crystal Structure of Glutathione-dependent Dehydroascorbate Reductase from Spinach Chloroplast

Tomonori Yadani^a, Shinichi Kurata^a, Hiroyoshi Matsumura^a, Tsuyoshi Inoue^a, Eiichi Mizohata^a, Taise Shimaoka^b, Chikahiro Miyake^b, Akiho Yokota^b, Yasushi Kai^a, ^aDepartment of Materials Chemistry, Graduate School of Engineering, Osaka University, Japan. ^bDepartment of Molecular biology, Graduate School of Biological Science, Nara Institute of Science and Technology (NAIST), Japan. E-mail: tomo@chem.eng.osaka-u.ac.jp

Glutathione-dependent dehydroascorbate reductase (GSH-DHAR) catalyzes the reduction of dehydroascorbate (DHA) to ascorbate using reduced glutathione as the electron donor. GSH-DHAR existing in chloroplast plays a pivotal role in the regeneration of ascorbate, which is oxidized to scavenge active oxygen species in the process of photosynthesis. The catalytic mechanism of the GSH-DHAR from spinach chloroplast is intriguing, because the specific constants for DHA and GSH are much higher than those of the other characterized DHARs.

Here, we report the three-dimensional structure of GSH-DHAR from spinach chloroplasts at 1.65 Å resolution, which has been determined by the multiwavelength anomalous dispersion (MAD) phasing method. The crystal structure reveals a monomeric form, which corresponds to the results observed in the analyses of gel filtration and dynamic light scattering. The structure is mainly composed of the similar domain to that of previously solved thioltransferase [1] and an extra alpha-helical domain. The catalytically essential cysteine was completely reduced, because it was crystallized in the solution including high concentration of DTT. The model study using the coordinates of glutathione transferases suggested that the putative glutathione binding site was formed by the amino residues corresponding to those of the other glutathione transferases. These observations support that glutathione could be bound near the catalytically essential cysteine. We will show the detailed reaction mechanism to describe how it establishes its high specificities.

[1] Katti S. K., *et al.*, *Protein Sci.*, 1995, **4**, 1998.

Keywords: crystal structure, dehydroascorbate, reductase

P.04.02.102

Acta Cryst. (2005). A61, C203-C204

Snapshots Along the PEPCK Catalytic Pathway

Todd Holyoak, Department of Biochemistry and Molecular Biology, The University of Kansas Medical Center, Kansas City, Kansas. E-mail: tholyoak@KUMC.edu

Phosphoenolpyruvate carboxykinase (PEPCK) catalyzes the reversible decarboxylation of oxaloacetic acid with the concomitant