

spectra we will discuss the electric structure, distortion of polyhedra related to ferroelectricity and thermal vibration.

[1] B. Ravel, E.A. Stern *Physica B* 208&209 **1995** 316-318 [2] T. Miyanaga, D. Diop, S.Ikeda, and H. Kon, *Ferroelectrics*, **2002** 274, 41-53

Keywords: ferroelectric, XANES, phase transition

MS56.P09

Acta Cryst. (2011) **A67**, C585

Temperature dependence of pre-edge feature in Ti K-edge XANES spectra for ATiO_3 , A_2TiO_4 (A=Mg, Ca, Fe, Sr and Ba) and TiO_2 compounds

Tatsuya Hiratoko,^a Tomotaka Nakatani,^a Hiroki Okudera,^b Akihiko Nakatsuka^c and Akira Yoshias,^a ^a*Graduate School of Sciences and Technology, Kumamoto University, Kumamoto 860-8555 (Japan).* ^b*Graduate School of Natural Science, Kanazawa University, Kanazawa, 920-1192 (Japan)* ^c*Department of Advanced Materials Science and Engineering, Yamaguchi University, Ube 755-8611 (Japan).* E-mail: 118d8052@st.kumamoto-u.ac.jp

X-ray absorption near edge structure (XANES) provides important information on the electronic structure and local symmetry around X-ray absorption atom. There are three distinct pre-edge peaks correspond to electronic transitions in Ti K-edge XANES spectra. The composition, structure and temperature dependence of XANES spectra on three peaks was investigated. We measured Ti K-edge XANES spectra of various titanates, MgTiO_3 , CaTiO_3 , SrTiO_3 , BaTiO_3 , Mg_2TiO_4 , Fe_2TiO_4 , TiO_2 rutile and anatase, in the temperature range from 20K to 800K. Ti atoms are placed in TiO_6 octahedral and TiO_4 tetrahedral sites in crystal structures. The measurements of Ti k-edge XANES spectra were carried out in transmission mode at beam line BL-7C and BL-9A of the Photon Factory in KEK, Tsukuba. High temperature X-ray absorption measurements were made under a helium atmosphere. XANES spectra of all sample on each peaks is increasing as the temperature increases except for tetragonal BaTiO_3 and tetragonal SrTiO_3 phases. TiO_2 rutile and anatase have largely different rate of rising pre-edge absorption to the temperature. The XANES spectra in the high temperature region were strongly affected by the harmonic thermal vibration of the atoms. There is an interesting relation between electronic transition and local distortion of coordination environment coming from the harmonic thermal vibration. We will discuss harmonic and anharmonic thermal vibration models by XAFS method.

Keywords: XANES, pre-edge peak, titanate


MS56.P10

Acta Cryst. (2011) **A67**, C585

A gold(III) complex of the neuroepileptic drug gabapentin

Demetrius C. Levendis, Ahmed Shaikjee, Helder M. Marques and Richard Mampa, *Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, PO WITS, 2050, Johannesburg, (South Africa).* Email: demetrius.levendis@wits.ac.za

Gabapentin, a neuro-epileptic drug, has been the subject of interest lately as new polymorphs, [1] salts and hydrates have been reported, as well as the high pressure crystallisation of a novel heptahydrate [2]. The first complexes with transition metal complexes, Cu(II) and Zn(II) were reported recently by Braga and co-workers [3]. Since gold is known to have pharmaceutical applications [4] we were interested to see if we could prepare a Au(III)-Gp complex.



Auric acid was reacted with gabapentin in an aqueous solution. Two distinct types of crystals precipitated from this solution: pale yellow needle shaped crystals (**I**) and dark yellow blocks (**II**, not shown here). Their structures were determined by single crystal structure analysis. The MeOD solution ^1H and ^{15}N -NMR were then recorded to confirm that the gold remains coordinated to the nitrogen of the amino group in solution (**I**).

The crystal structure of (**I**) clearly shows the Au-N coordination (Au-N bond 2.043(2) Å) in the hydrogen bonded dimer. The cyclohexane is disordered over two positions, both having a chair conformation. The ^1H -NMR shows peaks at $\delta = 2.45, 2.88$ ppm for the free gabapentin $-\text{CH}_2-$ groups ($-\text{CH}_2-\text{COOH}$ and $-\text{CH}_2-\text{NH}_3^+$ respectively). On dissolving (**I**) in MeOD four peaks are apparent in the ^1H -NMR; at 2.45, 2.51, 3.05 and 3.12 ppm. Our current interpretation of this is that (**I**) exists as two different isomers in solution and that on crystallisation these persist in the solid state.

In conclusion, auric acid reacts with gabapentin in aqueous solution to form a Au(III)-gabapentin complex in which the Au(III) is coordinated to the amino nitrogen [5]. It appears that in this reaction an intermediate complex has been trapped given that Au(III) is known to oxidise amino acids.

[1] H.A. Reece, D.C. Levendis, *Acta Cryst.* **2008**, C64, o105 [2] F.P.A. Fabbiani, D.C. Levendis, G. Buth, W.F. Kuhs, N. Shankland, H. Sowa, *CrystEngComm*, **2010**, 12, 2354. [3] D. Braga, F. Grepioni, L. Maini, R. Brescello, L. Cotarca, *CrystEngComm*. **2008**, 10, 469-471. [4] S.P. Fricker, *Transition Met. Chem.* **1996**, 21, 377. [5] A. Shaikjee, D.C. Levendis, H.M. Marques, R. Mampa, *Inorg. Chem. Comm.* **2011**, 14, 534-538.

Acknowledgments: University of the Witwatersrand and the National Research Foundation for funding; Manuel Fernandes for X-ray data collection and advice on the disorder refinement

Keywords: neuro-epileptic drug, gold(III) complex, gabapentin

MS56.P11

Acta Cryst. (2011) **A67**, C585-C586

Crystal structure of Zn complex with chelidamic acid and acridine

Zohreh Derikvand,^a Andya Nemati,^b Masomeh Mansouriki,^a ^a*Department of Chemistry, Faculty of Science, Islamic Azad University, Khorramabad Branch, Khorramabad, (Iran).* ^b*Iran Compiling Encyclopedia Foundation, Tajrish, Tehran, (Iran).* E-mail: zderik@yahoo.com

4-Hydroxypyridine-2,6-dicarboxylic acid as carboxylate derivative has drawn extensive attention in coordination chemistry. This ligand could potentially provide various coordination motifs to form both discrete and consecutive metal complexes under appropriate synthesis condition [1,2]. We prepared new mononuclear complex of Zn(II) containing 4-hydroxypyridine-2,6-dicarboxylic