

## Microsymposium

**MS20.O04**

### *Feasibility study of visibility-based X-ray photon correlation spectroscopy*

I. Inoue<sup>1,2</sup>, Y. Shinohara<sup>1,3</sup>, A. Watanabe<sup>1</sup>, Y. Amemiya<sup>1,3</sup>

<sup>1</sup>The University of Tokyo, Graduate School of Frontier Sciences, Kashiwa, Japan, <sup>2</sup>RIKEN SPring-8 Center, XFEL Research and Development Division, Sayo, Japan, <sup>3</sup>JST-CREST, Kawaguchi, Japan

When coherent X-rays impinge upon a disordered system, a grainy scattering pattern called speckle pattern is observed. If the system evolves with time, the corresponding speckle pattern also changes. Temporal changes in the speckle patterns therefore provide information on the dynamics of the system. This technique, which is called X-ray photon correlation spectroscopy (XPCS) [1], has shown the potential to access dynamic properties of various materials, such as colloidal suspensions, block copolymer, supercooled liquids, alloys, and antiferromagnetic materials. Although XPCS is a powerful technique for material science as recent studies show, it has a limitation of time resolution: dynamics faster than the frame rate of detector cannot be measured. When a two-dimensional (2D) detector is used in XPCS, the time resolution is limited to the order of milliseconds. For improving the time resolution of XPCS, we have extended speckle visibility spectroscopy (SVS) in the region of visible light [2] to the region of X-rays (X-ray SVS; XSVS) [3]. Since the minimum exposure time of the scattering patterns determines the time resolutions of XSVS and SVS, micro- or nano- second dynamics can be measured even with a 2D detector. Thus, XSVS has potential to bridge the time gap between XPCS and inelastic neutron/X-ray scattering techniques, and will be one of the promising tools for various science with the next generation synchrotron X-ray sources, such as diffraction limited storage rings and energy recovery linac based X-ray sources. In this presentation, we will describe the principle of XSVS and show the result of the application of XSVS to Brownian colloidal suspensions. This study was performed under the approval of JASRI (2011A1112, 2011B1131). We acknowledge Drs. N. Yagi and N. Ohta for their kind support in performing experiments.

[1] M. Sutton, *C. R. Phys.*, 2008, 9, 657-667, [2] P. K. Dixon and D. J. Durian, *Phys. Rev. Lett.*, 2003, 90, 184302, [3] I. Inoue et al., *Opt. Express*, 2012, 20, 26878-26887

**Keywords:** Coherent X-rays, X-ray speckle visibility spectroscopy, X-ray photon correlation spectroscopy