## **Poster Presentation**

## MS74.P01

## Crystallochemical aspects of two microlite-group new mineral species

M. Andrade<sup>1</sup>, J. Ellena<sup>1</sup>, D. Atencio<sup>2</sup>

<sup>1</sup>São Carlos Institute of Physics (IFSC-USP), Department of Physics and Informatics, Sao Carlos, Brazil, <sup>2</sup>Geosciences Institute (IGc-USP), Department of Mineralogy and Geotectonics, Sao Paulo, Brazil

Fluorcalciomicrolite,  $Ca_{1.5}Ta_2O_6F$ , and hydroxycalciomicrolite,  $Ca_{1.5}Ta_2O_6(OH)$ , are new microlite-group [1] minerals found in the Volta Grande pegmatite, Nazareno, Minas Gerais, Brazil. Both occur as octahedral and rhombododecahedral crystals. The crystals are colourless, yellow and translucent, with vitreous to resinous luster. The densities calculated for fluorcalciomicrolite [2] and hydroxycalciomicrolite are 6.160 and 6.176 g/cm³, respectively. The empirical formulae obtained from electron microprobe analysis are  $(Ca_{1.07}Na_{0.81} \bigcirc_{0.12})\Sigma_2$   $(Ta_{1.84}Nb_{0.14}Sn_{0.02})\Sigma_2[O_{5.93}(OH)_{0.07}]\Sigma_{6.00}[F_{0.79}(OH)_{0.21}]$  for fluorcalciomicrolite and  $(Ca_{1.48}Na_{0.06}Mn_{0.01})\Sigma_{1.55}(Ta_{1.88}Nb_{0.11}Sn_{0.01})\Sigma_2O_6[(OH)_{0.76}F_{0.20}O_{0.04}]$  for hydroxycalmicrolite. Fluorcalciomicrolite is cubic, space group Fd-3m, a = 10.4191(6) Å, V = 1131.07(11) ų, and Z = 8. Hydroxycalciomicrolite is also cubic; however, the presence of P-lattice is confirmed by the large number of weak reflections observed by X-ray diffraction. As a result, the space group is  $P4_332$  and unit-cell parameters are a = 10.4211(8) Å, and V = 1131.72(15) ų.

[1] D. Atencio, M. B. Andrade, A. G. Christy, R. Gieré, P. M. Kartashov, Canadian Mineralogist, 2010, 48, 673-698, [2] M. B. ANDRADE, D. Atencio, A. I. C. Persiano, J. Ellena, Mineralogical Magazine, 2013, 77, 2989-2996

Keywords: fluorcalciomicrolite, hydroxycalciomicrolite, microlite-group