

**References**

- CROMER, D. T. & LIBERMAN, D. (1970). *J. Chem. Phys.* **53**, 1891–1898.  
 CROMER, D. T. & WABER, J. T. (1965). *Acta Cryst.* **18**, 104–109.
- GOSSAUER, A. & HINZE, J.-P. (1981). *Hoppe-Seyler's Z. Physiol. Chem.* In the press.  
 SHELDICK, G. M. (1976). *SHELX*. A program for crystal structure determination. Univ. of Cambridge, England.  
 SHELDICK, W. S. (1981). *Acta Cryst.* **B37**, 299–300.

**SHORT COMMUNICATION**

*Contributions intended for publication under this heading should be expressly so marked; they should not exceed about 1000 words; they should be forwarded in the usual way to the appropriate Co-editor; they will be published as speedily as possible.*

*Acta Cryst.* (1981). **B37**, 302

**Racemic 3-[benzyl(phenyl)phosphinyl]-2-butenoic acid: errata.** By M. L. GŁÓWKA, *Institute of General Chemistry, Technical University of Łódź, 36 Zwirki, 90-924 Łódź, Poland*

(Received 20 August 1980; accepted 1 September 1980)

**Abstract**

E.s.d.'s for bond lengths and angles, the  $\gamma$  coordinate for C(3) [new value 0.7156 (9)], and the bond length C(16)–C(11) [new value 1.415 (10) Å] in the paper by Główka [*Acta Cryst.* (1978), **B34**, 3089–3091] are revised and a missing hydrogen-bond distance is added [O(1)…O(2) 2.587 (7) Å].

The  $\gamma$  coordinate of C(3) is 0.7156 (9). The remaining corrections are given in Table 1.

The author thanks Professor G. Ferguson for pointing out the errors.

Table 1. Bond lengths (Å) and angles (°)

P–O(1)	1.493 (4)	C(3)–C(4)	1.415 (14)	C(8)–C(9)	1.510 (9)	C(13)–C(14)	1.379 (13)
P–C(1)	1.796 (6)	C(4)–C(5)	1.402 (14)	C(9)–O(2)	1.310 (8)	C(14)–C(15)	1.402 (13)
P–C(7)	1.827 (7)	C(5)–C(6)	1.405 (10)	C(9)–O(3)	1.194 (8)	C(15)–C(16)	1.409 (10)
P–C(10)	1.827 (8)	C(6)–C(1)	1.402 (10)	C(10)–C(11)	1.520 (8)	C(16)–C(11)	1.415 (10)
C(1)–C(2)	1.410 (11)	C(7)–C(8)	1.333 (8)	C(11)–C(12)	1.404 (10)		
C(2)–C(3)	1.431 (10)	C(7)–C(17)	1.525 (8)	C(12)–C(13)	1.434 (10)	O(1)…O(2)	2.587 (7)
P–C(7)–C(8)	113.6 (5)	O(1)–P–C(7)	113.2 (3)	C(6)–C(1)–C(2)	121.2 (6)	C(12)–C(13)–C(14)	120.0 (7)
P–C(7)–C(17)	117.1 (5)	O(1)–P–C(10)	112.9 (3)	C(7)–C(8)–C(9)	127.5 (6)	C(13)–C(14)–C(15)	121.7 (7)
P–C(10)–C(11)	109.9 (5)	C(7)–P–C(10)	106.6 (3)	C(8)–C(7)–C(17)	129.3 (6)	C(14)–C(15)–C(16)	118.9 (8)
P–C(1)–C(2)	116.3 (5)	C(1)–C(2)–C(3)	119.1 (7)	C(8)–C(9)–O(2)	117.5 (5)	C(15)–C(16)–C(11)	120.4 (7)
P–C(1)–C(6)	122.5 (5)	C(2)–C(3)–C(4)	119.5 (8)	C(8)–C(9)–O(3)	119.5 (6)	C(16)–C(11)–C(12)	120.1 (6)
C(1)–P–O(1)	112.5 (3)	C(3)–C(4)–C(5)	119.9 (7)	O(2)–C(9)–O(3)	122.9 (7)	C(16)–C(11)–C(10)	120.7 (6)
C(1)–P–C(7)	105.0 (3)	C(4)–C(5)–C(6)	121.1 (7)	C(10)–C(11)–C(12)	119.2 (6)	C(9)–O(2)–H(1)	107 (3)
C(1)–P–C(10)	106.1 (3)	C(5)–C(6)–C(1)	119.2 (7)	C(11)–C(12)–C(13)	118.9 (7)	P–O(1)…H(1)	140 (2)

0567-7408/81/010302-01\$01.00

© 1981 International Union of Crystallography

**International Union of Crystallography**

*Acta Cryst.* (1981). **B37**, 302

**Submission of papers for publication in  
*Acta Crystallographica***

The unhindered selection by authors of the most convenient Co-editor for the submission of papers is one of the policies of *Acta Crystallographica* stated on the inside front cover of all issues. In recent years, a disproportionate number of

papers have been submitted to Dr P. J. Wheatley and Professor M. M. Woolfson. Many of these papers originated in countries outside the United Kingdom. With the resignations of Professor Woolfson and Dr Wheatley it would be helpful if the authors of such papers, particularly those resident in other European countries, would select a Co-editor not resident in the United Kingdom. Manuscripts receive comparable treatment from all Co-editors, and possible delays in handling or by manuscript transfer will be avoided by a more random manuscript distribution.