



# The oxygenating constituent of 3,6-diketocamphane monooxygenase from the CAM plasmid of *Pseudomonas putida*: the first crystal structure of a type II Baeyer–Villiger monooxygenase. Corrigendum

**Keywords:** FMN-dependent monooxygenase; protein structure; industrial biocatalysis; corrigendum

Michail N. Isupov,<sup>a</sup> Ewald Schröder,<sup>a</sup> Robert P. Gibson,<sup>a</sup> Jean Beecher,<sup>a</sup> Giuliana Donadio,<sup>a</sup> Vahid Saneei,<sup>a</sup> Stephlina A. Dcunha,<sup>a</sup> Emma J. McGhie,<sup>a</sup> Christopher Sayer,<sup>a</sup> Colin F. Davenport,<sup>a</sup> Peter C. K. Lau,<sup>b</sup> Yoshie Hasegawa,<sup>c</sup> Hiroaki Iwaki,<sup>c</sup> Maria Kadow,<sup>d</sup> Kathleen Balke,<sup>d</sup> Uwe T. Bornscheuer,<sup>d</sup> Gleb Bourenkov<sup>e</sup> and Jennifer A. Littlechild<sup>a\*</sup>

<sup>a</sup>The Henry Wellcome Building for Biocatalysis, Biosciences, College of Life and Environmental Sciences, University of Exeter, Stocker Road, Exeter EX4 4QD, England, <sup>b</sup>Biotechnology Research Institute, National Research Council Canada, 6100 Royalmount Avenue, Montreal, QC H4P 2R2, Canada, <sup>c</sup>Department of Biotechnology, Faculty of Engineering, Kansai University, Japan, <sup>d</sup>Department of Biotechnology and Enzyme Catalysis, Institute of Biochemistry, Greifswald University, Felix-Hausdorff-Strasse 4, 17487 Greifswald, Germany, and <sup>e</sup>European Molecular Biology Laboratory (EMBL), Hamburg Outstation, Notkestrasse 85, 22607 Hamburg, Germany. \*Correspondence e-mail: j.a.littlechild@exeter.ac.uk

A statement is amended in the article by Isupov *et al.* [(2015). *Acta Cryst.* **D71**, 2344–2353].

In the article by Isupov *et al.* (2015), the statement in §3.10 ‘Partial sequencing of the large CAM plasmid has now identified a flavin reductase adjacent to the 3,6-DKMO gene on the CAM plasmid (Littlechild & Isupov, unpublished data)’ is withdrawn. The following statement is added: ‘A cognate flavin reductase (Fred) from the genome of *Pseudomonas putida* (strain PpCam), which provides the required FMNH<sub>2</sub> for the 3,6-DKMO activity, has been described by Iwaki *et al.* (2013). Other non-cognate reductases have been reported which can partially support the DKMO activity but not as efficiently as the cognate Fred’.

## References

- Isupov, M. N., Schröder, E., Gibson, R. P., Beecher, J., Donadio, G., Saneei, V., Dcunha, S. A., McGhie, E. J., Sayer, C., Davenport, C. F., Lau, P. C., Hasegawa, Y., Iwaki, H., Kadow, M., Balke, K., Bornscheuer, U. T., Bourenkov, G. & Littlechild, J. A. (2015). *Acta Cryst.* **D71**, 2344–2353.
- Iwaki, H., Grosse, S., Bergeron, H., Leisch, H., Morley, K., Hasegawa, Y. & Lau, P. C. K. (2013). *Appl. Environ. Microbiol.* **79**, 3282–3293.

