

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Reinvestigation of the crystal structure of lautite, CuAsS

Luca Bindi,^a Tiziano Catelani,^b Laura Chelazzi^b and Paola Bonazzi^{b*}

^aMuseo di Storia Naturale, Sezione di Mineralogia, Università degli Studi di Firenze, Via La Pira 4, I-50121 Firenze, Italy, and ^bDipartimento di Scienze della Terra, Università degli Studi di Firenze, Via La Pira 4, I-50121 Firenze, Italy
Correspondence e-mail: pbcry@geo.unifi.it

Received 25 January 2008; accepted 14 February 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{As}-\text{S}) = 0.002$ Å; R factor = 0.046; wR factor = 0.109; data-to-parameter ratio = 30.2.

The crystal structure of the mineral lautite (copper arsenic sulfide), CuAsS, previously described as either centrosymmetric [*Pnma*; Marumo & Nowacki (1964). *Schweiz. Miner. Petro. Mitt.* **44**, 439–454] or noncentrosymmetric [*Pna2*₁; Craig & Stephenson (1965). *Acta Cryst.* **19**, 543–547], was reinvestigated by means of single-crystal X-ray diffraction. The centrosymmetric structural model reported previously was confirmed, although with improved precision for the atomic coordinates and interatomic distances. Lautite shows a sphalerite-derivative structure with a linking of Cu[AsS₃], As[CuAs₂S] and S[Cu₃As] tetrahedra. All atoms lie on special positions (Wyckoff position 4*c*, site symmetry *m*).

Related literature

For related literature, see: Craig & Stephenson (1965); Marumo & Nowacki (1964); Wyckoff (1963).

Experimental

Crystal data

AsCuS	$V = 232.24(10)$ Å ³
$M_r = 170.54$	$Z = 4$
Orthorhombic, <i>Pnma</i>	Mo $K\alpha$ radiation
$a = 11.347(4)$ Å	$\mu = 24.00$ mm ⁻¹
$b = 3.7533(7)$ Å	$T = 298(2)$ K
$c = 5.453(1)$ Å	$0.12 \times 0.10 \times 0.08$ mm

Data collection

Bruker <i>P4</i> diffractometer	483 reflections with $I > 2\sigma(I)$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$R_{\text{int}} = 0.077$
$T_{\text{min}} = 0.070$, $T_{\text{max}} = 0.150$	3 standard reflections every 150 reflections
3824 measured reflections	intensity decay: none
574 independent reflections	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	19 parameters
$wR(F^2) = 0.108$	$\Delta\rho_{\text{max}} = 1.28$ e Å ⁻³
$S = 1.09$	$\Delta\rho_{\text{min}} = -1.07$ e Å ⁻³
574 reflections	

Data collection: *XSCANS* (Bruker, 1997); cell refinement: *XSCANS*; data reduction: *XSCANS*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XtalDraw* (Downs & Hall-Wallace, 2003); software used to prepare material for publication: *SHELXL97*.

This work was funded by CNR, Istituto di Geoscienze e Georisorse, Sezione di Firenze.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FI2059).

References

- Bruker (1997). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Craig, D. C. & Stephenson, N. C. (1965). *Acta Cryst.* **19**, 543–547.
 Downs, R. T. & Hall-Wallace, M. (2003). *Am. Mineral.* **88**, 247–250.
 Marumo, F. & Nowacki, W. (1964). *Schweiz. Miner. Petro. Mitt.* **44**, 439–454.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wyckoff, R. W. G. (1963). *Crystal Structures*, 2nd ed. New York: Interscience Publishers.

supporting information

Acta Cryst. (2008). E64, i22 [doi:10.1107/S1600536808004492]

Reinvestigation of the crystal structure of lautite, CuAsS

Luca Bindi, Tiziano Catelani, Laura Chelazzi and Paola Bonazzi

S1. Comment

The crystal structure of lautite was solved by Marumo & Nowacki (1964) in the space group $Pnma$ ($R = 9.0\%$) and by Craig & Stephenson (1965) in the space group $Pna2_1$ ($R = 13.7\%$) by means of photographic data and three-dimensional Patterson-function. The low quality of the structural data given by these authors, however, did not allow to obtain an anisotropic model of the structure. Nevertheless, the topologies and interatomic distances of both centrosymmetric and non-centrosymmetric models are very similar.

Although the structural results obtained by Craig & Stephenson (1965) indicate the acentricity of the structure of CuAsS, no clear crystal-chemical reason for the choice of a noncentrosymmetric arrangement was given. To help resolve the concerns relating to the structure of natural lautite, we present new crystal structure data for lautite from its type locality (*i.e.*, Marienberg, Saxony, Germany).


The centrosymmetric structural model previously reported by Marumo & Nowacki (1964) was confirmed, although a higher precision of refinement was achieved (e.s.d. improved by a factor of two) and refinement with anisotropic displacement parameters could be performed (Fig. 1). All atoms lie on special positions (Wyckoff position 4c, site symmetry m). Lautite shows a sphalerite-derivative structure with a linking of $\text{Cu}[\text{AsS}_3]$, $\text{As}[\text{CuAs}_2\text{S}]$ and $\text{S}[\text{Cu}_3\text{As}]$ tetrahedra (Fig. 2). Within the framework, the As atoms form zigzag As—As chains along [010] exhibiting As—As bond distances [2.4965 (8) Å] and angles [97.48 (4)°] resembling the covalent As—As linkage observed within the sheets of the crystal structure of arsenic (Wyckoff, 1963).

S2. Experimental


A crystal was selected from a natural specimen belonging to the Mineralogical Collection of the Natural History Museum of Florence (catalogue number 44202/G).

S3. Refinement

The crystal structure refinement was performed starting from the atomic coordinates reported by Marumo & Nowacki (1964). Convergence was rapidly obtained for an anisotropic model of the structure.


Figure 1

The crystal structure of lautite down [010]. Displacement parameters are drawn at the 70% probability level. The unit-cell is outlined. Symmetry codes are: (i) $-x + 1/2; -y; z + 1/2$; (ii) $x + 1/2; -y + 1/2; -z + 1/2$; (iii) $-x; y + 1/2; -z$.


Figure 2

The crystal structure of lautite showing the linking of $\text{Cu}[\text{AsS}_3]$, $\text{As}[\text{CuAs}_2\text{S}]$ and $\text{S}[\text{Cu}_3\text{As}]$ tetrahedra. Blu, red and yellow circles indicate Cu, As and S atoms, respectively. The unit-cell is outlined.

copper arsenic sulfide

Crystal data

AsCuS

$M_r = 170.54$

Orthorhombic, $Pnma$

Hall symbol: $-P\ 2ac\ 2n$

$a = 11.347\ (4)\ \text{\AA}$

$b = 3.7533\ (7)\ \text{\AA}$

$c = 5.453\ (1)\ \text{\AA}$

$V = 232.24\ (10)\ \text{\AA}^3$

$Z = 4$

$F(000) = 312$

$D_x = 4.878\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 38 reflections

$\theta = 12.5\text{--}24.3^\circ$

$\mu = 24.00\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, black

$0.12 \times 0.10 \times 0.08\ \text{mm}$

Data collection

Bruker P4 diffractometer	574 independent reflections
Radiation source: fine-focus sealed tube	483 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.077$
ω scans	$\theta_{\text{max}} = 35.0^\circ$, $\theta_{\text{min}} = 3.6^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = -18 \rightarrow 18$
$T_{\text{min}} = 0.070$, $T_{\text{max}} = 0.150$	$k = -6 \rightarrow 6$
3824 measured reflections	$l = -8 \rightarrow 8$
	3 standard reflections every 150 reflections
	intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0856P)^2 + 1.9844P]$
$R[F^2 > 2\sigma(F^2)] = 0.046$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.108$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.09$	$\Delta\rho_{\text{max}} = 1.28$ at 0.0736 0.2500 0.3457 (0.68 Å from As) $e \text{ \AA}^{-3}$
574 reflections	$\Delta\rho_{\text{min}} = -1.07$ at 0.0052 0.0883 0.4402 (0.78 Å from As) $e \text{ \AA}^{-3}$
19 parameters	
0 restraints	
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	0.17454 (7)	0.2500	0.06264 (18)	0.0165 (2)
As	0.01373 (5)	0.2500	0.35177 (11)	0.00894 (18)
S	0.16576 (12)	0.7500	0.8196 (3)	0.0100 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu	0.0171 (3)	0.0168 (4)	0.0157 (4)	0.000	-0.0015 (3)	0.000
As	0.0105 (3)	0.0084 (3)	0.0080 (3)	0.000	0.00083 (17)	0.000
S	0.0108 (5)	0.0114 (5)	0.0079 (5)	0.000	-0.0005 (4)	0.000

Geometric parameters (\AA , $^\circ$)

Cu—S ⁱ	2.2908 (16)	As—As ^v	2.4965 (8)
Cu—S ⁱⁱ	2.2996 (10)	S—As ^{iv}	2.2408 (16)
Cu—S ⁱⁱⁱ	2.2996 (10)	S—Cu ^{vi}	2.2908 (16)

Cu—As	2.4114 (11)	S—Cu ^{vii}	2.2996 (10)
As—S ^{iv}	2.2408 (16)	S—Cu ^{viii}	2.2996 (10)
As—As ^{iv}	2.4965 (8)		
S ⁱ —Cu—S ⁱⁱ	112.76 (3)	S ^{iv} —As—As ^v	99.01 (4)
S ⁱ —Cu—S ⁱⁱⁱ	112.76 (3)	Cu—As—As ^v	121.19 (3)
S ⁱⁱ —Cu—S ⁱⁱⁱ	109.39 (7)	As ^{iv} —As—As ^v	97.48 (4)
S ⁱ —Cu—As	101.46 (5)	As ^{iv} —S—Cu ^{vi}	117.64 (7)
S ⁱⁱ —Cu—As	110.12 (4)	As ^{iv} —S—Cu ^{vii}	106.24 (5)
S ⁱⁱⁱ —Cu—As	110.12 (4)	Cu ^{vi} —S—Cu ^{vii}	108.56 (4)
S ^{iv} —As—Cu	114.52 (5)	As ^{iv} —S—Cu ^{viii}	106.24 (5)
S ^{iv} —As—As ^{iv}	99.01 (4)	Cu ^{vi} —S—Cu ^{viii}	108.56 (4)
Cu—As—As ^{iv}	121.19 (3)	Cu ^{vii} —S—Cu ^{viii}	109.39 (7)

Symmetry codes: (i) $-x+1/2, -y+1, z-1/2$; (ii) $x, y-1, z-1$; (iii) $x, y, z-1$; (iv) $-x, -y+1, -z+1$; (v) $-x, -y, -z+1$; (vi) $-x+1/2, -y+1, z+1/2$; (vii) $x, y+1, z+1$; (viii) $x, y, z+1$.