

Bis(2-bromopyridinium) hexachlorido-stannate(IV)

Basem Fares Ali,^a Rawhi Al-Far^{b*} and Salim F. Haddad^c

^aDepartment of Chemistry, Al al-Bayt University, Mafraq, Jordan, ^bFaculty of Science and IT, Al-Balqa'a Applied University, Salt, Jordan, and ^cDepartment of Chemistry, The University of Jordan, Amman, Jordan
Correspondence e-mail: rohi@bau.edu.jo


Received 6 March 2008; accepted 3 April 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.040; wR factor = 0.098; data-to-parameter ratio = 18.3.

The asymmetric unit of the title compound, $(\text{C}_5\text{H}_5\text{BrN})_2[\text{SnCl}_6]$, contains one cation and one half-anion. The $[\text{SnCl}_6]^{2-}$ anion is located on an inversion center and forms a quasi-regular octahedral arrangement. Hydrogen-bonding interactions of two kinds, *viz.* N—H···Cl—Sn and C—H···Cl—Sn, along with Cl···Br interactions [3.4393 (15) Å], connect the ions in the crystal structure into two-dimensional supramolecular arrays. These supramolecular arrays are arranged in layers approximately parallel to (110) built up from anions interacting with six symmetry-related surrounding cations.

Related literature

The title salt is isomorphous with the Te-analogue, see: Fernandes *et al.* (2004). For related literature, see: Al-Far & Ali (2007); Ali, Al-Far & Al-Sou'od (2007); Ali & Al-Far (2007); Ali, Al-Far & Ng (2007); Allen *et al.* (1987); Aruta *et al.* (2005); Awwadi *et al.* (2007); Bouacida *et al.* (2007); Ellis & Macdonald (2006); Hill (1998); Kagan *et al.* (1999); Knutson *et al.* (2005); Li *et al.* (2005); Mitzi *et al.* (2001); Raptopoulou *et al.* (2002); Willett & Haddad (2000).



Experimental

Crystal data

$(\text{C}_5\text{H}_5\text{BrN})_2[\text{SnCl}_6]$
 $M_r = 649.41$

Monoclinic, $P2_1/n$
 $a = 9.0843$ (14) Å

$b = 10.6827$ (9) Å
 $c = 10.6345$ (17) Å
 $\beta = 109.843$ (11)°
 $V = 970.8$ (2) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 6.25$ mm⁻¹
 $T = 296$ (2) K
 $0.20 \times 0.15 \times 0.10$ mm

Data collection

Siemens P4 diffractometer
Absorption correction: ψ scan (*XSCANS*; Siemens, 1996)
 $T_{\min} = 0.340$, $T_{\max} = 0.535$

2385 measured reflections
1791 independent reflections
1343 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.097$
 $S = 1.05$
1791 reflections

98 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.61$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.73$ e Å⁻³

Table 1
Selected geometric parameters (Å, °).

Sn1—Cl1	2.4216 (13)	Sn1—Cl3	2.4212 (13)
Sn1—Cl2	2.4513 (14)		
Cl1—Sn1—Cl2	89.67 (5)	Cl3—Sn1—Cl2	90.06 (6)
Cl3—Sn1—Cl1 ⁱ	89.70 (5)	Cl3—Sn1—Cl2 ⁱ	89.94 (6)
Cl3—Sn1—Cl1	90.30 (5)		

Symmetry code: (i) $-x, -y, -z + 1$.

Table 2
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···Cl2 ⁱⁱ	0.86	2.45	3.234 (5)	151
C3—H3···Cl1 ⁱⁱⁱ	0.93	2.77	3.646 (6)	158
C5—H5···Cl1 ^{iv}	0.93	2.86	3.774 (7)	170

Symmetry codes: (ii) $-x + 1, -y, -z + 2$; (iii) $-x + 1, -y, -z + 1$; (iv) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS*; data reduction: *SHELXTL* (Sheldrick, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Al al-Bayt University and Al-Balqa'a Applied University are thanked for supporting this work

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2165).

References

- Al-Far, R. & Ali, B. F. (2007). *Acta Cryst. C* **63**, m137–m139.
- Ali, B. F. & Al-Far, R. (2007). *Acta Cryst. E* **63**, m892–m894.
- Ali, B. F., Al-Far, R. & Al-Sou'od, K. (2007). *J. Chem. Crystallogr.* **37**, 265–273.
- Ali, B. F., Al-Far, R. & Ng, S. W. (2007). *Acta Cryst. E* **63**, m2102–m2103.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Aruta, C., Licci, F., Zappettini, A., Bolzoni, F., Rastelli, F., Ferro, P. & Besagni, T. (2005). *Appl. Phys. A*, **81**, 963–968.
- Awwadi, F. F., Willett, R. D., Peterson, K. A. & Twamley, B. (2007). *J. Phys. Chem. A*, **111**, 2319–2328.

metal-organic compounds

- Bouacida, S., Merazig, H., Benard-Rocherulle, P. & Rizzoli, C. (2007). *Acta Cryst. E* **63**, m379–m381.
- Ellis, B. D. & Macdonald, C. L. B. (2006). *Acta Cryst. E* **62**, m1235–m1236.
- Fernandes, R. M. Jr, de Oliveira, G. M., Lang, E. S. & Vázquez-López, E. M. (2004). *Z. Anorg. Allg. Chem.* **630**, 2687–2691.
- Hill, C. L. (1998). *Chem. Rev.* **98**, 1–2.
- Kagan, C. R., Mitzi, D. B. & Dimitrakopoulos, C. D. (1999). *Science*, **286**, 945–947.
- Knutson, J. L., Martin, J. D. & Mitzi, D. B. (2005). *Inorg. Chem.* **44**, 4699–4705.
- Li, H.-T., Sun, R., Shi, H.-P. & Huang, S.-P. (2005). *Acta Cryst. E* **61**, m2088–m2089.
- Mitzi, D. B., Dimitrakopoulos, C. D. & Kosbar, L. L. (2001). *Chem. Mater.* **13**, 3728–3740.
- Raptopoulou, C. P., Terzis, A., Mousdis, G. A. & Papavassiliou, G. C. (2002). *Z. Naturforsch. Teil B*, **57**, 645–650.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *XSCANS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Willett, R. D. & Haddad, S. F. (2000). *Acta Cryst. C* **56**, e438.

supporting information

Acta Cryst. (2008). E64, m637–m638 [doi:10.1107/S160053680800901X]

Bis(2-bromopyridinium) hexachloridostannate(IV)

Basem Fares Ali, Rawhi Al-Far and Salim F. Haddad

S1. Comment

Noncovalent interactions play an important role in organizing structural units in both natural and artificial systems. Hybrid organic-inorganic compounds are of great interest owing to their ionic, electrical, magnetic and optical properties (Hill, 1998; Kagan *et al.*, 1999; Raptopoulou *et al.*, 2002). Tin metal-halo based hybrids are of particular interest as being materials with interesting optical and magnetic properties (Aruta *et al.*, 2005; Knutson *et al.*, 2005; Mitzi *et al.*, 2001; Kagan *et al.*, 1999). We are currently carrying out studies about lattice including different types of intermolecular interactions. Our strategy is to use aromatic compounds to encourage aryl···aryl packing arrangements of various types, using substituted pyridinium in order to facilitates associations, and halo salts that can involve in $X\cdots X$ interactions as well as $X\cdots$ aryl and $X\cdots$ H interactions. Within our research of hybrid compounds containing tin metal (Al-Far & Ali 2007; Ali, Al-Far & Al-Sou'od, 2007; Ali & Al-Far, 2007; Ali, Al-Far & Ng, 2007), the crystal structure of the title salt, (I), has been investigated.

The asymmetric unit of (I) contains one cation and one-half anion (Fig. 1). The $(\text{SnCl}_6)^{2-}$ anion lies on an inversion center, in a quasi-octahedral geometry (Table 1). The Sn—Cl bond lengths are almost invariant, but Sn—Cl2 is longer than the others (involved in the shortest hydrogen bonds). These lengths fall within the range of tin-chloride distances reported previously for compounds containing $(\text{SnCl}_6)^{2-}$ anions (Bouacida *et al.*, 2007; Ellis & Macdonald, 2006; Li *et al.*, 2005; Willett & Haddad, 2000). Bond lengths and angles within the cation are as expected (Allen *et al.*, 1987).

The packing of the structure (Fig. 2) can be described as layers of alternating anions (zigzag orientation) along the face parallel to b -axis and diagonal to ac plane. Each $(\text{SnCl}_6)^{2-}$ anion is surrounded by six cations *via* four equatorial (C,N)— $H\cdots Cl$ interactions (Table 2) and two axial $Cl\cdots Br$ interactions [$Cl3\cdots Br2^i = 3.4393$ (15) Å; symmetry code: (*i*) $-1/2 + x$, $-1/2 - y$, $-1/2 + z$; Fig. 3]. This arrangement of molecules appears as layers approximately parallel to [110]. It is noteworthy that structural and theoretical results (Awwadi *et al.*, 2007; and references therein), show the significance of linear $C-Y\cdots X$ (in this case $C-Cl\cdots Br$) synthons in influencing structures of crystalline materials and in use as potential building blocks in crystal engineering *via* supramolecular synthesis.


The intermolecular hydrogen bonds (Table 2) and $Cl\cdots Br$ interactions would therefore add some lattice stability. This is evident in the isostructurality with the reported Te analogue (Fernandes *et al.*, 2004).

S2. Experimental


Warm solution of SnCl_4 (1.0 mmol) dissolved in absolute ethanol (10 ml) and concentrated HCl (1 ml), was added dropwise to a stirred hot solution of 2-bromopyridine (1 mmol) dissolved in ethanol (10 ml). The mixture was treated with another 2 ml of concentrated HCl and refluxed for 2 h, then cooled, filtered off, and allowed to stand undisturbed at room temperature. The salt crystallized over 1 d as nice yellow block crystals (yield: 89.6%).

S3. Refinement


H atoms were positioned geometrically, with N—H = 0.86 Å (for NH) and C—H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

**Figure 1**

A view of the asymmetric unit of (I), with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A packing diagram of (I). Hydrogen bonds and Cl...Br interactions are shown as dashed lines.

**Figure 3**

Part of the cell contents of (I), showing Cl···Br and (C,N)—H···Cl intermolecular interactions (dashed lines) for one $(\text{SnCl}_6)^{2-}$ anion and six surrounding cations.

Bis(2-bromopyridinium) hexachloridostannate(IV)

Crystal data



$M_r = 649.41$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.0843$ (14) Å

$b = 10.6827$ (9) Å

$c = 10.6345$ (17) Å

$\beta = 109.843$ (11)°

$V = 970.8$ (2) Å³

$Z = 2$

$F(000) = 612$

$D_x = 2.222 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 90 reflections

$\theta = 1.6\text{--}27.4^\circ$

$\mu = 6.25 \text{ mm}^{-1}$

$T = 296$ K

Block, yellow

0.20 × 0.15 × 0.10 mm

Data collection

Siemens P4

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 3 pixels mm⁻¹

ω scans

Absorption correction: ψ scan
(*XSCANS*; Siemens, 1996)

$T_{\min} = 0.340$, $T_{\max} = 0.535$

2385 measured reflections

1791 independent reflections

1343 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$

$\theta_{\max} = 25.5^\circ$, $\theta_{\min} = 2.8^\circ$

$h = -1 \rightarrow 11$

$k = -12 \rightarrow 1$

$l = -12 \rightarrow 12$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.040$$

$$wR(F^2) = 0.097$$

$$S = 1.05$$

1791 reflections

98 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0417P)^2 + 0.8983P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\text{max}} < 0.001$$

$$\Delta\rho_{\text{max}} = 0.61 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\text{min}} = -0.73 \text{ e \AA}^{-3}$$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0145 (10)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.0000	0.0000	0.5000	0.0271 (2)
Br2	0.68293 (9)	0.00098 (6)	0.92745 (7)	0.0595 (3)
Cl1	0.15178 (16)	0.07315 (13)	0.36630 (13)	0.0376 (4)
Cl2	0.24265 (16)	-0.03276 (15)	0.68883 (14)	0.0474 (4)
Cl3	0.01440 (19)	-0.21235 (13)	0.42546 (16)	0.0500 (4)
N1	0.8532 (5)	0.1873 (5)	1.0891 (4)	0.0410 (11)
H1	0.7983	0.1643	1.1369	0.049*
C6	0.9552 (7)	0.2804 (5)	1.1327 (6)	0.0485 (15)
H6	0.9655	0.3203	1.2130	0.058*
C5	1.0445 (8)	0.3170 (7)	1.0591 (7)	0.0612 (19)
H5	1.1162	0.3820	1.0881	0.073*
C2	0.8330 (6)	0.1285 (5)	0.9736 (5)	0.0368 (12)
C4	1.0264 (8)	0.2556 (7)	0.9409 (7)	0.0621 (19)
H4	1.0880	0.2782	0.8904	0.075*
C3	0.9181 (7)	0.1612 (6)	0.8968 (6)	0.0518 (16)
H3	0.9039	0.1210	0.8160	0.062*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0278 (3)	0.0278 (3)	0.0285 (3)	0.0023 (2)	0.0134 (2)	0.0006 (2)
Br2	0.0574 (5)	0.0572 (4)	0.0634 (5)	-0.0230 (3)	0.0197 (3)	-0.0074 (3)
Cl1	0.0379 (7)	0.0438 (7)	0.0379 (7)	-0.0022 (6)	0.0216 (6)	0.0021 (6)
Cl2	0.0334 (7)	0.0680 (9)	0.0382 (8)	-0.0004 (7)	0.0088 (6)	0.0147 (7)
Cl3	0.0604 (10)	0.0319 (7)	0.0698 (10)	0.0009 (7)	0.0378 (8)	-0.0086 (7)
N1	0.040 (3)	0.050 (3)	0.035 (2)	-0.003 (2)	0.016 (2)	0.001 (2)
C6	0.042 (3)	0.045 (3)	0.051 (4)	0.000 (3)	0.005 (3)	-0.010 (3)
C5	0.046 (4)	0.050 (4)	0.076 (5)	-0.010 (3)	0.005 (3)	0.007 (4)
C2	0.031 (3)	0.036 (3)	0.041 (3)	-0.002 (2)	0.009 (2)	0.001 (2)
C4	0.050 (4)	0.076 (5)	0.069 (5)	-0.013 (4)	0.033 (3)	0.002 (4)
C3	0.054 (4)	0.062 (4)	0.050 (4)	-0.009 (3)	0.031 (3)	-0.011 (3)

Geometric parameters (\AA , $^\circ$)

Sn1—Cl1	2.4216 (13)	N1—H1	0.8600
Sn1—Cl2	2.4513 (14)	C6—C5	1.362 (9)
Sn1—Cl3	2.4212 (13)	C6—H6	0.9300
Sn1—Cl1 ⁱ	2.4216 (13)	C5—C4	1.378 (10)
Sn1—Cl2 ⁱ	2.4513 (14)	C5—H5	0.9300
Sn1—Cl3 ⁱ	2.4212 (13)	C2—C3	1.347 (8)
Br2—C2	1.871 (5)	C4—C3	1.375 (9)
N1—C6	1.331 (8)	C4—H4	0.9300
N1—C2	1.336 (7)	C3—H3	0.9300
Cl1—Sn1—Cl2	89.67 (5)	C2—N1—H1	118.9
Cl3—Sn1—Cl1 ⁱ	89.70 (5)	N1—C6—C5	119.6 (6)
Cl3—Sn1—Cl1	90.30 (5)	N1—C6—H6	120.2
Cl3—Sn1—Cl2	90.06 (6)	C5—C6—H6	120.2
Cl3—Sn1—Cl2 ⁱ	89.94 (6)	C6—C5—C4	118.6 (6)
Cl1—Sn1—Cl2 ⁱ	90.33 (5)	C6—C5—H5	120.7
Cl3 ⁱ —Sn1—Cl3	180.0	C4—C5—H5	120.7
Cl3 ⁱ —Sn1—Cl1 ⁱ	90.30 (5)	N1—C2—C3	120.5 (5)
Cl3 ⁱ —Sn1—Cl1	89.70 (5)	N1—C2—Br2	116.4 (4)
Cl1 ⁱ —Sn1—Cl1	180.0	C3—C2—Br2	123.1 (5)
Cl3 ⁱ —Sn1—Cl2 ⁱ	90.06 (6)	C3—C4—C5	120.7 (7)
Cl1 ⁱ —Sn1—Cl2 ⁱ	89.67 (5)	C3—C4—H4	119.6
Cl3 ⁱ —Sn1—Cl2	89.94 (6)	C5—C4—H4	119.6
Cl1 ⁱ —Sn1—Cl2	90.33 (5)	C2—C3—C4	118.4 (6)
Cl2 ⁱ —Sn1—Cl2	180.0	C2—C3—H3	120.8
C6—N1—C2	122.3 (5)	C4—C3—H3	120.8
C6—N1—H1	118.9	 	
C2—N1—C6—C5	-0.9 (9)	C6—C5—C4—C3	1.4 (11)
N1—C6—C5—C4	-0.1 (10)	N1—C2—C3—C4	0.6 (9)
C6—N1—C2—C3	0.7 (9)	Br2—C2—C3—C4	-179.4 (5)
C6—N1—C2—Br2	-179.4 (4)	C5—C4—C3—C2	-1.6 (11)

Symmetry code: (i) $-x, -y, -z+1$.Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1 ⁱⁱ —Cl2 ⁱⁱ	0.86	2.45	3.234 (5)	151
C3—H3 ⁱⁱⁱ —Cl1 ⁱⁱⁱ	0.93	2.77	3.646 (6)	158
C5—H5 ^{iv} —Cl1 ^{iv}	0.93	2.86	3.774 (7)	170

Symmetry codes: (ii) $-x+1, -y, -z+2$; (iii) $-x+1, -y, -z+1$; (iv) $-x+3/2, y+1/2, -z+3/2$.