

## N'-(5-Bromo-2-methoxybenzylidene)-3-hydroxybenzohydrazide methanol hemisolvate

Zhi Zhou

Department of Chemistry, Kaili College, Kaili Guizhou 556000, People's Republic of China  
Correspondence e-mail: zhou82zhi@126.com


Received 13 June 2008; accepted 16 June 2008

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.008 \text{ \AA}$ ;  $R$  factor = 0.062;  $wR$  factor = 0.170; data-to-parameter ratio = 16.4.

The asymmetric unit of the title compound,  $\text{C}_{15}\text{H}_{13}\text{BrN}_2\text{O}_3 \cdot 0.5\text{CH}_3\text{OH}$ , contains two Schiff base molecules and a methanol molecule of crystallization. The dihedral angles between the benzene rings in the two molecules are 23.8 (2) and 18.6 (2) $^\circ$ . In the crystal structure, molecules are linked through intermolecular N—H $\cdots$ O, O—H $\cdots$ O and O—H $\cdots$ N hydrogen bonds, forming a three-dimensional network.

### Related literature

For related literature, see: Zhou & Tang (2007); Zhou & Xiao (2007). For related structures, see: Ali *et al.* (2007); Butcher *et al.* (2007); He (2008); Jing & Yu (2007); Nie (2008).



### Experimental

#### Crystal data

$\text{C}_{15}\text{H}_{13}\text{BrN}_2\text{O}_3 \cdot 0.5\text{CH}_3\text{OH}$	$b = 11.177 (2) \text{ \AA}$
$M_r = 365.21$	$c = 22.607 (3) \text{ \AA}$
Monoclinic, $P2_1/n$	$\beta = 93.706 (3)^\circ$
$a = 12.906 (2) \text{ \AA}$	$V = 3254.3 (9) \text{ \AA}^3$

$Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 2.54 \text{ mm}^{-1}$   
 $T = 298 (2) \text{ K}$   
 $0.20 \times 0.18 \times 0.17 \text{ mm}$

#### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)  
 $T_{\min} = 0.630$ ,  $T_{\max} = 0.672$   
21623 measured reflections  
6725 independent reflections  
2610 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.104$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.170$   
 $S = 0.98$   
6725 reflections  
409 parameters  
2 restraints  
H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.47 \text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots\text{A}$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
N4—H4A $\cdots$ O2 <sup>i</sup>	0.902 (10)	2.045 (18)	2.923 (5)	164 (5)
N2—H2 $\cdots$ O7 <sup>i</sup>	0.894 (10)	1.977 (13)	2.866 (5)	173 (5)
O7—H7 $\cdots$ O3	0.82	1.96	2.737 (5)	157
O6—H6 $\cdots$ N1 <sup>ii</sup>	0.82	2.48	3.140 (5)	138
O6—H6 $\cdots$ O2 <sup>ii</sup>	0.82	2.06	2.777 (5)	146
O3—H3 $\cdots$ N3 <sup>iii</sup>	0.82	2.64	3.110 (6)	118
O3—H3 $\cdots$ O5 <sup>iii</sup>	0.82	1.92	2.692 (5)	157

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $x - 1, y + 1, z$ ; (iii)  $x + 1, y, z$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The author thanks Kaili College for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2575).

### References

- Ali, H. M., Zuraini, K., Wan Jeffrey, B. & Ng, S. W. (2007). *Acta Cryst. E63*, o1729–o1730.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Butcher, R. J., Jasinski, J. P., Narayana, B., Sunil, K. & Yathirajan, H. S. (2007). *Acta Cryst. E63*, o3652.
- He, L. (2008). *Acta Cryst. E64*, o82.
- Jing, Z.-L. & Yu, M. (2007). *Acta Cryst. E63*, o509–o510.
- Nie, Y. (2008). *Acta Cryst. E64*, o471.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Zhou, Z. & Tang, R.-R. (2007). *Acta Cryst. E63*, m2960.
- Zhou, Z. & Xiao, Z.-H. (2007). *Acta Cryst. E63*, m2012.

# supporting information

*Acta Cryst.* (2008). E64, o1298 [doi:10.1107/S1600536808018205]

## **N'-(5-Bromo-2-methoxybenzylidene)-3-hydroxybenzohydrazide methanol hemisolvate**

**Zhi Zhou**

### **S1. Comment**

Recently, we have reported two metal complexes with Schiff base ligands (Zhou & Tang, 2007; Zhou & Xiao, 2007). We report herein the crystal structure of the title Schiff base compound (I), Fig. 1.

The asymmetric unit of (I) consists of two Schiff base molecules and a methanol molecule of crystallization. The dihedral angles are 23.8 (2) ° and 18.6 (2) °, respectively, between the benzene rings (C1-C6) and (C10-C15) for molecule A, and (C16-C21), (C25-C30) for molecule B. All the bond values are comparable to the similar compounds (Ali *et al.*, 2007; Nie, 2008; He, 2008; Butcher *et al.*, 2007; Jing & Yu, 2007).


In the crystal structure, molecules are linked through intermolecular N—H···O, O—H···O and O—H···N hydrogen bonds (Table 1) to form a three-dimensional network (Fig. 2).

### **S2. Experimental**


2-Methoxy-5-bromobenzaldehyde (1.0 mmol, 215.0 mg) and 3-hydroxybenzohydrazide (1.0 mmol, 152.1 mg) were dissolved in methanol (30 ml). The mixture was stirred at reflux for 30 min to give a colourless solution. After keeping the solution in air for a few days, colourless block-like crystals were formed.

### **S3. Refinement**

H2 and H4A were located in a difference Fourier map and refined isotropically, with  $U_{\text{iso}}$  fixed at 0.08 Å<sup>2</sup>. Other H atoms were positioned geometrically and refined using a riding model with d(O—H) = 0.82 Å,  $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{O})$ , and d(C—H) = 0.93 - 0.96 Å,  $U_{\text{iso}} = 1.2$  or  $1.5U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of (I), with displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

The molecular packing of (I), viewed along the  $c$  axis. Hydrogen bonds are shown as dashed lines.

### *N'*-(5-Bromo-2-methoxybenzylidene)-3-hydroxybenzohydrazide methanol hemisolvate

#### Crystal data

$C_{15}H_{13}BrN_2O_3 \cdot 0.5CH_4O$

$M_r = 365.21$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 12.906 (2)$  Å

$b = 11.177 (2)$  Å

$c = 22.607 (3) \text{ \AA}$   
 $\beta = 93.706 (3)^\circ$   
 $V = 3254.3 (9) \text{ \AA}^3$   
 $Z = 8$   
 $F(000) = 1480$   
 $D_x = 1.491 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1690 reflections  
 $\theta = 2.4\text{--}24.1^\circ$   
 $\mu = 2.54 \text{ mm}^{-1}$   
 $T = 298 \text{ K}$   
Block, colourless  
 $0.20 \times 0.18 \times 0.17 \text{ mm}$

#### Data collection

Bruker SMART CCD area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Bruker, 2001)  
 $T_{\min} = 0.630$ ,  $T_{\max} = 0.672$

21623 measured reflections  
6725 independent reflections  
2610 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.104$   
 $\theta_{\max} = 26.5^\circ$ ,  $\theta_{\min} = 1.8^\circ$   
 $h = -16 \rightarrow 16$   
 $k = -13 \rightarrow 14$   
 $l = -28 \rightarrow 27$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.170$   
 $S = 0.98$   
6725 reflections  
409 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0522P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.47 \text{ e \AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.89733 (6)	-0.03746 (6)	0.19990 (3)	0.0896 (3)
Br2	-0.27606 (8)	0.47236 (11)	0.74505 (4)	0.1632 (5)
O1	0.4978 (3)	-0.1249 (3)	0.31384 (17)	0.0705 (11)
O2	0.8013 (2)	0.2809 (3)	0.44744 (14)	0.0499 (9)
O3	0.6978 (3)	0.6557 (3)	0.55948 (18)	0.0680 (11)
H3	0.7389	0.6728	0.5346	0.102*
O4	0.1231 (4)	0.3770 (4)	0.6320 (2)	0.0868 (14)
O5	-0.1758 (3)	0.7734 (3)	0.48927 (15)	0.0563 (10)
O6	-0.0715 (3)	1.1472 (3)	0.37821 (15)	0.0551 (10)

H6	-0.1226	1.1603	0.3970	0.083*
O7	0.5718 (3)	0.8519 (4)	0.5649 (3)	0.0984 (16)
H7	0.6018	0.7923	0.5537	0.148*
N1	0.6877 (3)	0.1219 (3)	0.38704 (17)	0.0435 (10)
N2	0.6465 (3)	0.1906 (4)	0.43049 (18)	0.0454 (11)
N3	-0.0639 (3)	0.6176 (4)	0.55665 (18)	0.0489 (11)
N4	-0.0209 (3)	0.6887 (4)	0.51548 (18)	0.0448 (11)
C1	0.6526 (4)	-0.0203 (5)	0.3091 (2)	0.0476 (14)
C2	0.5840 (4)	-0.1083 (5)	0.2848 (2)	0.0494 (14)
C3	0.6097 (5)	-0.1693 (5)	0.2347 (2)	0.0599 (16)
H3A	0.5641	-0.2261	0.2179	0.072*
C4	0.7018 (5)	-0.1475 (5)	0.2094 (2)	0.0623 (16)
H4	0.7181	-0.1892	0.1756	0.075*
C5	0.7690 (4)	-0.0647 (5)	0.2338 (2)	0.0532 (15)
C6	0.7453 (4)	-0.0001 (4)	0.2831 (2)	0.0466 (14)
H6A	0.7916	0.0571	0.2990	0.056*
C7	0.4258 (4)	-0.2131 (5)	0.2916 (3)	0.0795 (19)
H7A	0.4041	-0.1949	0.2512	0.119*
H7B	0.3664	-0.2133	0.3151	0.119*
H7C	0.4582	-0.2904	0.2936	0.119*
C8	0.6233 (4)	0.0517 (5)	0.3595 (2)	0.0485 (14)
H8	0.5560	0.0460	0.3717	0.058*
C9	0.7095 (4)	0.2717 (4)	0.4583 (2)	0.0403 (12)
C10	0.6619 (4)	0.3512 (4)	0.5019 (2)	0.0424 (13)
C11	0.7043 (4)	0.4646 (4)	0.5105 (2)	0.0441 (13)
H11	0.7613	0.4877	0.4901	0.053*
C12	0.6614 (4)	0.5429 (5)	0.5494 (2)	0.0475 (13)
C13	0.5772 (4)	0.5085 (5)	0.5794 (2)	0.0562 (15)
H13	0.5487	0.5610	0.6058	0.067*
C14	0.5352 (4)	0.3978 (5)	0.5707 (2)	0.0610 (16)
H14	0.4775	0.3758	0.5908	0.073*
C15	0.5772 (4)	0.3173 (5)	0.5321 (2)	0.0533 (15)
H15	0.5486	0.2414	0.5267	0.064*
C16	0.0358 (7)	0.3912 (5)	0.6599 (3)	0.075 (2)
C17	-0.0323 (5)	0.4802 (5)	0.6358 (2)	0.0563 (16)
C18	-0.1248 (5)	0.5010 (5)	0.6608 (3)	0.0716 (19)
H18	-0.1693	0.5598	0.6448	0.086*
C19	-0.1527 (7)	0.4368 (8)	0.7089 (3)	0.105 (3)
C20	-0.0841 (10)	0.3495 (9)	0.7327 (4)	0.134 (5)
H20	-0.1013	0.3062	0.7658	0.161*
C21	0.0056 (9)	0.3278 (7)	0.7085 (3)	0.116 (4)
H21	0.0491	0.2684	0.7247	0.140*
C22	0.1976 (6)	0.2885 (6)	0.6549 (3)	0.125 (3)
H22A	0.2229	0.3105	0.6942	0.187*
H22B	0.2546	0.2846	0.6297	0.187*
H22C	0.1645	0.2116	0.6559	0.187*
C23	0.0025 (4)	0.5525 (4)	0.5873 (2)	0.0501 (14)
H23	0.0719	0.5518	0.5784	0.060*

C24	-0.0810 (4)	0.7700 (4)	0.4857 (2)	0.0422 (13)
C25	-0.0264 (4)	0.8531 (4)	0.4470 (2)	0.0411 (13)
C26	-0.0762 (4)	0.9601 (4)	0.43118 (19)	0.0412 (12)
H26	-0.1412	0.9771	0.4445	0.049*
C27	-0.0283 (4)	1.0406 (5)	0.3956 (2)	0.0429 (13)
C28	0.0680 (4)	1.0164 (5)	0.3763 (2)	0.0561 (15)
H28	0.1007	1.0720	0.3532	0.067*
C29	0.1162 (4)	0.9099 (5)	0.3910 (2)	0.0577 (15)
H29	0.1806	0.8927	0.3768	0.069*
C30	0.0696 (4)	0.8282 (4)	0.4267 (2)	0.0455 (13)
H30	0.1030	0.7568	0.4370	0.055*
C31	0.6289 (6)	0.9488 (7)	0.5556 (5)	0.173 (5)
H31A	0.5889	1.0194	0.5622	0.260*
H31B	0.6491	0.9484	0.5154	0.260*
H31C	0.6898	0.9483	0.5823	0.260*
H2	0.5782 (11)	0.183 (5)	0.434 (2)	0.080*
H4A	0.0489 (9)	0.695 (5)	0.520 (2)	0.080*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.1071 (6)	0.0787 (5)	0.0892 (5)	0.0026 (4)	0.0552 (4)	0.0009 (4)
Br2	0.1725 (10)	0.2271 (12)	0.0980 (7)	-0.1320 (9)	0.0695 (6)	-0.0473 (7)
O1	0.057 (3)	0.071 (3)	0.084 (3)	-0.018 (2)	0.008 (2)	-0.022 (2)
O2	0.035 (2)	0.056 (2)	0.059 (2)	0.0003 (18)	0.0038 (17)	-0.0118 (18)
O3	0.067 (3)	0.046 (2)	0.096 (3)	-0.010 (2)	0.042 (2)	-0.018 (2)
O4	0.119 (4)	0.055 (3)	0.082 (3)	0.019 (3)	-0.028 (3)	0.003 (2)
O5	0.040 (2)	0.060 (2)	0.070 (3)	0.0084 (19)	0.0117 (19)	0.0173 (19)
O6	0.061 (3)	0.046 (2)	0.060 (2)	0.0045 (19)	0.0187 (19)	0.0093 (19)
O7	0.047 (3)	0.057 (3)	0.193 (5)	-0.005 (2)	0.028 (3)	-0.036 (3)
N1	0.047 (3)	0.036 (3)	0.048 (3)	0.000 (2)	0.004 (2)	-0.008 (2)
N2	0.039 (3)	0.044 (3)	0.054 (3)	-0.006 (2)	0.012 (2)	-0.017 (2)
N3	0.049 (3)	0.046 (3)	0.052 (3)	-0.010 (2)	0.005 (2)	-0.001 (2)
N4	0.033 (3)	0.048 (3)	0.054 (3)	-0.004 (2)	0.004 (2)	0.011 (2)
C1	0.060 (4)	0.041 (3)	0.042 (3)	0.002 (3)	0.000 (3)	0.001 (3)
C2	0.055 (4)	0.041 (3)	0.051 (4)	0.007 (3)	-0.004 (3)	-0.006 (3)
C3	0.081 (5)	0.045 (4)	0.051 (4)	0.001 (3)	-0.011 (3)	-0.006 (3)
C4	0.094 (5)	0.046 (4)	0.047 (4)	0.002 (4)	0.012 (3)	-0.009 (3)
C5	0.073 (4)	0.043 (4)	0.046 (3)	0.000 (3)	0.019 (3)	0.005 (3)
C6	0.054 (4)	0.036 (3)	0.050 (3)	-0.002 (3)	0.002 (3)	0.002 (3)
C7	0.056 (4)	0.071 (4)	0.110 (5)	-0.014 (4)	-0.009 (4)	0.000 (4)
C8	0.045 (3)	0.049 (4)	0.053 (3)	-0.001 (3)	0.014 (3)	-0.004 (3)
C9	0.041 (3)	0.036 (3)	0.044 (3)	0.002 (3)	0.004 (3)	0.001 (3)
C10	0.038 (3)	0.042 (3)	0.047 (3)	0.002 (3)	0.004 (2)	-0.005 (3)
C11	0.034 (3)	0.045 (3)	0.054 (3)	0.000 (3)	0.011 (2)	-0.004 (3)
C12	0.042 (3)	0.043 (3)	0.059 (3)	-0.002 (3)	0.015 (3)	-0.007 (3)
C13	0.064 (4)	0.050 (4)	0.058 (4)	-0.003 (3)	0.025 (3)	-0.018 (3)
C14	0.064 (4)	0.065 (4)	0.057 (4)	-0.011 (3)	0.028 (3)	-0.013 (3)

C15	0.059 (4)	0.049 (4)	0.053 (4)	-0.020 (3)	0.013 (3)	-0.011 (3)
C16	0.134 (7)	0.033 (4)	0.055 (5)	-0.025 (4)	-0.021 (5)	0.008 (3)
C17	0.075 (4)	0.049 (4)	0.044 (4)	-0.025 (3)	-0.002 (3)	0.003 (3)
C18	0.096 (5)	0.069 (5)	0.048 (4)	-0.043 (4)	-0.002 (4)	-0.001 (3)
C19	0.151 (8)	0.114 (7)	0.050 (5)	-0.074 (6)	0.016 (5)	-0.008 (5)
C20	0.235 (15)	0.115 (9)	0.052 (6)	-0.108 (10)	0.002 (7)	0.018 (5)
C21	0.219 (12)	0.061 (5)	0.063 (6)	-0.039 (7)	-0.034 (6)	0.020 (5)
C22	0.175 (8)	0.067 (5)	0.120 (6)	0.046 (5)	-0.082 (6)	-0.020 (4)
C23	0.054 (4)	0.042 (3)	0.054 (4)	-0.001 (3)	0.002 (3)	0.001 (3)
C24	0.038 (3)	0.041 (3)	0.047 (3)	-0.001 (3)	0.001 (3)	-0.004 (3)
C25	0.040 (3)	0.042 (3)	0.041 (3)	-0.006 (3)	0.002 (2)	-0.001 (3)
C26	0.040 (3)	0.048 (3)	0.035 (3)	0.000 (3)	0.000 (2)	-0.003 (3)
C27	0.050 (3)	0.038 (3)	0.042 (3)	0.002 (3)	0.005 (3)	-0.002 (3)
C28	0.051 (4)	0.057 (4)	0.062 (4)	-0.004 (3)	0.018 (3)	0.010 (3)
C29	0.043 (3)	0.061 (4)	0.071 (4)	0.003 (3)	0.022 (3)	0.005 (3)
C30	0.038 (3)	0.046 (3)	0.052 (3)	0.009 (3)	0.005 (3)	0.000 (3)
C31	0.093 (6)	0.077 (6)	0.357 (15)	-0.022 (5)	0.076 (8)	-0.013 (8)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Br1—C5	1.893 (5)	C10—C11	1.389 (6)
Br2—C19	1.879 (9)	C11—C12	1.382 (6)
O1—C2	1.342 (6)	C11—H11	0.9300
O1—C7	1.423 (6)	C12—C13	1.372 (7)
O2—C9	1.230 (5)	C13—C14	1.360 (7)
O3—C12	1.360 (5)	C13—H13	0.9300
O3—H3	0.8200	C14—C15	1.387 (7)
O4—C16	1.336 (8)	C14—H14	0.9300
O4—C22	1.452 (6)	C15—H15	0.9300
O5—C24	1.233 (5)	C16—C21	1.384 (9)
O6—C27	1.363 (5)	C16—C17	1.413 (8)
O6—H6	0.8200	C17—C18	1.374 (8)
O7—C31	1.335 (7)	C17—C23	1.456 (7)
O7—H7	0.8200	C18—C19	1.370 (9)
N1—C8	1.276 (5)	C18—H18	0.9300
N1—N2	1.379 (5)	C19—C20	1.401 (12)
N2—C9	1.346 (6)	C20—C21	1.335 (12)
N2—H2	0.894 (10)	C20—H20	0.9300
N3—C23	1.290 (6)	C21—H21	0.9300
N3—N4	1.369 (5)	C22—H22A	0.9600
N4—C24	1.347 (6)	C22—H22B	0.9600
N4—H4A	0.902 (10)	C22—H22C	0.9600
C1—C6	1.384 (7)	C23—H23	0.9300
C1—C2	1.411 (7)	C24—C25	1.484 (6)
C1—C8	1.465 (7)	C25—C30	1.377 (6)
C2—C3	1.380 (7)	C25—C26	1.393 (6)
C3—C4	1.374 (7)	C26—C27	1.380 (6)
C3—H3A	0.9300	C26—H26	0.9300

C4—C5	1.361 (7)	C27—C28	1.371 (7)
C4—H4	0.9300	C28—C29	1.374 (7)
C5—C6	1.380 (7)	C28—H28	0.9300
C6—H6A	0.9300	C29—C30	1.382 (6)
C7—H7A	0.9600	C29—H29	0.9300
C7—H7B	0.9600	C30—H30	0.9300
C7—H7C	0.9600	C31—H31A	0.9600
C8—H8	0.9300	C31—H31B	0.9600
C9—C10	1.489 (6)	C31—H31C	0.9600
C10—C15	1.379 (6)		
C2—O1—C7	117.8 (4)	C10—C15—C14	119.2 (5)
C12—O3—H3	109.5	C10—C15—H15	120.4
C16—O4—C22	118.1 (6)	C14—C15—H15	120.4
C27—O6—H6	109.5	O4—C16—C21	127.1 (8)
C31—O7—H7	109.5	O4—C16—C17	115.0 (6)
C8—N1—N2	114.9 (4)	C21—C16—C17	117.8 (8)
C9—N2—N1	117.3 (4)	C18—C17—C16	119.7 (6)
C9—N2—H2	126 (3)	C18—C17—C23	122.3 (6)
N1—N2—H2	116 (3)	C16—C17—C23	117.9 (6)
C23—N3—N4	114.1 (4)	C19—C18—C17	121.2 (7)
C24—N4—N3	119.1 (4)	C19—C18—H18	119.4
C24—N4—H4A	123 (3)	C17—C18—H18	119.4
N3—N4—H4A	115 (3)	C18—C19—C20	118.5 (9)
C6—C1—C2	119.1 (5)	C18—C19—Br2	120.6 (8)
C6—C1—C8	120.9 (5)	C20—C19—Br2	120.8 (7)
C2—C1—C8	119.9 (5)	C21—C20—C19	120.8 (9)
O1—C2—C3	125.8 (5)	C21—C20—H20	119.6
O1—C2—C1	115.2 (5)	C19—C20—H20	119.6
C3—C2—C1	119.1 (5)	C20—C21—C16	121.9 (10)
C4—C3—C2	120.9 (5)	C20—C21—H21	119.0
C4—C3—H3A	119.5	C16—C21—H21	119.0
C2—C3—H3A	119.5	O4—C22—H22A	109.5
C5—C4—C3	119.9 (5)	O4—C22—H22B	109.5
C5—C4—H4	120.1	H22A—C22—H22B	109.5
C3—C4—H4	120.1	O4—C22—H22C	109.5
C4—C5—C6	121.0 (5)	H22A—C22—H22C	109.5
C4—C5—Br1	119.8 (4)	H22B—C22—H22C	109.5
C6—C5—Br1	119.2 (4)	N3—C23—C17	119.4 (5)
C5—C6—C1	120.0 (5)	N3—C23—H23	120.3
C5—C6—H6A	120.0	C17—C23—H23	120.3
C1—C6—H6A	120.0	O5—C24—N4	122.0 (5)
O1—C7—H7A	109.5	O5—C24—C25	122.1 (5)
O1—C7—H7B	109.5	N4—C24—C25	116.0 (5)
H7A—C7—H7B	109.5	C30—C25—C26	119.9 (5)
O1—C7—H7C	109.5	C30—C25—C24	122.8 (5)
H7A—C7—H7C	109.5	C26—C25—C24	117.4 (4)
H7B—C7—H7C	109.5	C27—C26—C25	119.6 (5)

N1—C8—C1	121.7 (5)	C27—C26—H26	120.2
N1—C8—H8	119.2	C25—C26—H26	120.2
C1—C8—H8	119.2	O6—C27—C28	116.3 (5)
O2—C9—N2	121.6 (5)	O6—C27—C26	123.3 (5)
O2—C9—C10	121.7 (5)	C28—C27—C26	120.4 (5)
N2—C9—C10	116.7 (4)	C27—C28—C29	120.0 (5)
C15—C10—C11	119.9 (4)	C27—C28—H28	120.0
C15—C10—C9	122.5 (5)	C29—C28—H28	120.0
C11—C10—C9	117.5 (4)	C28—C29—C30	120.4 (5)
C12—C11—C10	119.8 (5)	C28—C29—H29	119.8
C12—C11—H11	120.1	C30—C29—H29	119.8
C10—C11—H11	120.1	C25—C30—C29	119.7 (5)
O3—C12—C13	117.0 (5)	C25—C30—H30	120.1
O3—C12—C11	123.1 (5)	C29—C30—H30	120.1
C13—C12—C11	119.9 (5)	O7—C31—H31A	109.5
C14—C13—C12	120.3 (5)	O7—C31—H31B	109.5
C14—C13—H13	119.8	H31A—C31—H31B	109.5
C12—C13—H13	119.8	O7—C31—H31C	109.5
C13—C14—C15	120.8 (5)	H31A—C31—H31C	109.5
C13—C14—H14	119.6	H31B—C31—H31C	109.5
C15—C14—H14	119.6		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N4—H4A···O2 <sup>i</sup>	0.90 (1)	2.05 (2)	2.923 (5)	164 (5)
N2—H2···O7 <sup>i</sup>	0.89 (1)	1.98 (1)	2.866 (5)	173 (5)
O7—H7···O3	0.82	1.96	2.737 (5)	157
O6—H6···N1 <sup>ii</sup>	0.82	2.48	3.140 (5)	138
O6—H6···O2 <sup>ii</sup>	0.82	2.06	2.777 (5)	146
O3—H3···N3 <sup>iii</sup>	0.82	2.64	3.110 (6)	118
O3—H3···O5 <sup>iii</sup>	0.82	1.92	2.692 (5)	157

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x-1, y+1, z$ ; (iii)  $x+1, y, z$ .