

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,4-Bis(chloromethyl)naphthalene

Muhammad Ilyas Tariq,^a M. Nawaz Tahir,^{b*} Ishtiaq Hussain^a and Ayesha Roohi^c^aDepartment of Chemistry, University of Sargodha, Sargodha, Pakistan, ^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan, and ^cInstitute of Chemistry, University of the Punjab, Lahore 54590, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 16 July 2008; accepted 19 July 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.054; wR factor = 0.167; data-to-parameter ratio = 16.3.

In the title molecule, $\text{C}_{12}\text{H}_{10}\text{Cl}_2$, the torsion angles $\text{C}_r-\text{C}_r-\text{C}_m-\text{Cl}$ around the C_m-C_r bonds have values of -104.1 (4) and -101.9 (4)°, where C_m is a methylene and C_r is a ring C atom. The molecules related by translation along the b axis are arranged into stacks by $\pi-\pi$ interactions between unsubstituted and substituted aromatic rings of the naphthalene ring system (centroid-centroid distance = 3.940 Å).

Related literature

For related literature, see: Basaran *et al.* (1992); Gabe & Glusker (1971); Garriz *et al.* (2004); Ikeda *et al.* (1987); Kazakov (2003); Li *et al.* (2004); Mitchell & Iyer (1989); Tariq *et al.* (2008); Zhang *et al.* (1989, 2007).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_{10}\text{Cl}_2$
 $M_r = 225.10$

 Monoclinic, $P2_1/c$
 $a = 13.6887$ (11) Å

 $b = 4.5835$ (3) Å
 $c = 17.8278$ (13) Å
 $\beta = 109.666$ (4)°
 $V = 1053.31$ (13) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.57$ mm⁻¹
 $T = 296$ (2) K
 $0.25 \times 0.08 \times 0.04$ mm

Data collection

 Bruker Kappa APEX2
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.943$, $T_{\max} = 0.974$

 10310 measured reflections
 2073 independent reflections
 1220 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.167$
 $S = 1.05$
 2073 reflections

 127 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.79$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³

Data collection: APEX2 (Bruker, 2007); cell refinement: APEX2; data reduction: SAINT (Bruker, 2007); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2003); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

The authors acknowledge the Higher Education Commission, Islamabad, Pakistan, for funding the purchase of the diffractometer at GCU, Lahore.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2159).

References

- Basaran, R., Dou, S.-Q. & Weiss, A. (1992). *Ber. Bunsenges. Phys. Chem.* **96**, 1688–1692.
 Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc. Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Gabe, E. J. & Glusker, J. P. (1971). *Acta Cryst.* **B27**, 1925–1930.
 Garriz, P. I., Alvarez, H. L. & Colavita, G. M. (2004). *Acta Hort.* **636**, 325–330.
 Ikeda, T., Uchida, T., Shimizui, T. & Aoki, K. (1987). Jpn. Patent 62 158 243.
 Kazakov, P. V. (2003). *Khim. Farm. Zh.* **37**, 23–25.
 Li, J., Huang, X. & Huang, H. (2004). *Huanan Nongye Daxue Xuebao Ziran Kexueban*, **25**, 10–12.
 Mitchell, R. H. & Iyer, V. S. (1989). *Synlett*, **1**, 55–57.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Tariq, M. I., Siddiqui, H. L., Hussain, I. & Afzal, S. (2008). *J. Chem. Soc. Pak.* **30**, 472–476.
 Zhang, M., Su, P., Meng, X.-G. & Xu, X.-M. (2007). *Acta Cryst.* **E63**, o951–o952.
 Zhang, S., Xiao, S., Lai, H. & Fan, J. (1989). *Shiyou Huagong*, **18**, 263–265.

supporting information

Acta Cryst. (2008). E64, o1596 [doi:10.1107/S1600536808022757]

1,4-Bis(chloromethyl)naphthalene

Muhammad Ilyas Tariq, M. Nawaz Tahir, Ishtiaq Hussain and Ayesha Roohi

S1. Comment

Naphthalene acetic acid (NAA) is well known growth regulator/stimulator for different varieties of fruits and vegetables (Garriz *et al.*, 2004; Li *et al.*, 2004). Its synthesis has developed a great interest among the chemists and several methods have been reported. One of them is chloromethylation of naphthalene using methylene chloride in the presence of catalysts (Kazakov, 2003; Mitchell & Iyer, 1989; Zhang *et al.*, 1989). During the synthesis of NAA, using formaline and a mixture of acids as a source of insertion of methylene group (Ikeda *et al.*, 1987; Tariq *et al.*, 2008), the title compound has been isolated.

The crystal structures of 1,4-bis(bromomethyl)benzene (Zhang *et al.*, 2007), 1,4-bis(chloromethyl)benzene (Basaran *et al.*, 1992) and 9,10-bis(chloromethyl)anthracene (Gabe & Glusker, 1971) were published but no analogous derivatives of naphthalene have been reported.

The bond lengths in the naphthalene system are in the range of 1.344 (6)–1.425 (5) Å. The Cl atoms deviate in opposite directions from the plane of the naphthalene ring by 1.660 (6) Å and 1.559 (6) Å. The closest contacts of Cl atoms with neighbouring molecules are: 3.491 (5) Å for C11 \cdots C9ⁱ and 3.5581 (16) Å for C12 \cdots Cl2ⁱⁱ [symmetry codes: (i) $-x, 1 - y, -z$; (ii) $1 - x, 1 - y, 1 - z$].

S2. Experimental

A mixture of naphthalene (40.0 g), paraformaldehyde (35.0 g), glacial acetic acid (82.0 ml), H₃PO₄ (52.0 ml) and concentrated HCl (114.0 ml) was heated in a water bath at 358 K with vigorous stirring for 2 h. Thereafter, the mixture was cooled to room temperature. A solid product was obtained and isolated. It was thoroughly washed with water, ether and n-hexane, respectively in order to remove unreacted material. The product was further purified in hot methanol. Needle-shaped colorless crystals (m.p. 394–396 K) were obtained by recrystallization from ethyl acetate.

S3. Refinement

H atoms were positioned geometrically, with C—H = 0.93 and 0.97 Å for aromatic and methylene C-atoms, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$.




Figure 1

The *ORTEP* diagram of the title compound with displacement ellipsoids at the 50% probability level. H-atoms are shown by small circles of arbitrary radii.




Figure 2

The crystal packing diagram.

1,4-Bis(chloromethyl)naphthalene

Crystal data

$C_{12}H_{10}Cl_2$

$M_r = 225.10$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 13.6887$ (11) Å

$b = 4.5835$ (3) Å

$c = 17.8278$ (13) Å

$\beta = 109.666$ (4)°

$V = 1053.31$ (13) Å³

$Z = 4$

$F(000) = 464$

$D_x = 1.419$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2073 reflections

$\theta = 1.6$ – 26.0 °

$\mu = 0.57$ mm⁻¹

$T = 296$ K

Needle, colourless

$0.25 \times 0.08 \times 0.04$ mm

Data collection

Bruker Kappa APEX2

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 7.40 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.943$, $T_{\max} = 0.974$

10310 measured reflections

2073 independent reflections

1220 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.049$

$\theta_{\max} = 26.0$ °, $\theta_{\min} = 1.6$ °

$h = -16 \rightarrow 16$

$k = -5 \rightarrow 5$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.167$
 $S = 1.05$
 2073 reflections
 127 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0763P)^2 + 0.5487P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.79 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.15383 (10)	0.5096 (3)	-0.06255 (6)	0.0825 (5)
C12	0.41801 (9)	0.6216 (3)	0.40488 (6)	0.0694 (4)
C1	0.2211 (3)	0.3836 (7)	0.1364 (2)	0.0438 (11)
C2	0.2846 (3)	0.4596 (8)	0.0908 (2)	0.0479 (12)
C3	0.3638 (3)	0.6492 (9)	0.1214 (2)	0.0566 (14)
C4	0.3843 (3)	0.7720 (8)	0.1975 (3)	0.0572 (14)
C5	0.3249 (3)	0.7058 (7)	0.2432 (2)	0.0472 (11)
C6	0.2416 (3)	0.5067 (7)	0.2133 (2)	0.0428 (11)
C7	0.1764 (3)	0.4271 (8)	0.2572 (2)	0.0537 (12)
C8	0.0976 (3)	0.2349 (10)	0.2267 (3)	0.0635 (16)
C9	0.0776 (3)	0.1120 (9)	0.1520 (3)	0.0642 (16)
C10	0.1366 (3)	0.1849 (8)	0.1080 (2)	0.0546 (12)
C11	0.2650 (3)	0.3370 (10)	0.0095 (2)	0.0681 (17)
C12	0.3470 (3)	0.8534 (9)	0.3220 (2)	0.0631 (16)
H3	0.40549	0.69924	0.09161	0.0680*
H4	0.43949	0.90101	0.21711	0.0681*
H7	0.18769	0.50745	0.30737	0.0642*
H8	0.05609	0.18432	0.25656	0.0759*
H9	0.02356	-0.02056	0.13242	0.0771*
H10	0.12179	0.10348	0.05762	0.0655*
H11A	0.25290	0.12868	0.01017	0.0818*
H11B	0.32575	0.36693	-0.00605	0.0818*
H12A	0.28193	0.91087	0.32822	0.0755*
H12B	0.38692	1.02904	0.32298	0.0755*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0917 (9)	0.0956 (9)	0.0466 (7)	0.0066 (7)	0.0055 (6)	0.0041 (6)
C12	0.0783 (8)	0.0692 (7)	0.0492 (6)	0.0150 (5)	0.0065 (5)	0.0007 (5)
C1	0.043 (2)	0.0421 (18)	0.042 (2)	0.0109 (16)	0.0085 (16)	0.0062 (16)
C2	0.050 (2)	0.050 (2)	0.043 (2)	0.0115 (18)	0.0149 (17)	0.0073 (17)
C3	0.053 (2)	0.064 (2)	0.055 (3)	0.005 (2)	0.021 (2)	0.015 (2)
C4	0.048 (2)	0.049 (2)	0.064 (3)	-0.0019 (18)	0.005 (2)	0.010 (2)
C5	0.053 (2)	0.0411 (19)	0.041 (2)	0.0119 (17)	0.0071 (18)	0.0081 (16)
C6	0.041 (2)	0.0400 (18)	0.044 (2)	0.0095 (16)	0.0097 (16)	0.0065 (16)
C7	0.052 (2)	0.058 (2)	0.051 (2)	0.0138 (19)	0.0173 (19)	0.0106 (18)
C8	0.051 (2)	0.069 (3)	0.075 (3)	0.009 (2)	0.027 (2)	0.020 (2)
C9	0.054 (3)	0.061 (2)	0.070 (3)	-0.005 (2)	0.011 (2)	0.011 (2)
C10	0.056 (2)	0.051 (2)	0.049 (2)	0.0012 (18)	0.0074 (19)	0.0033 (18)
C11	0.079 (3)	0.072 (3)	0.055 (3)	0.014 (2)	0.025 (2)	0.004 (2)
C12	0.072 (3)	0.049 (2)	0.057 (3)	0.010 (2)	0.007 (2)	-0.0009 (19)

Geometric parameters (\AA , $^\circ$)

C11—C11	1.810 (4)	C8—C9	1.386 (7)
C12—C12	1.814 (4)	C9—C10	1.344 (6)
C1—C2	1.419 (6)	C3—H3	0.9300
C1—C6	1.421 (5)	C4—H4	0.9300
C1—C10	1.425 (5)	C7—H7	0.9300
C2—C3	1.353 (6)	C8—H8	0.9300
C2—C11	1.492 (5)	C9—H9	0.9300
C3—C4	1.407 (6)	C10—H10	0.9300
C4—C5	1.365 (6)	C11—H11A	0.9700
C5—C6	1.417 (5)	C11—H11B	0.9700
C5—C12	1.496 (5)	C12—H12A	0.9700
C6—C7	1.419 (6)	C12—H12B	0.9700
C7—C8	1.357 (6)		
C11...C10	3.464 (4)	C10...H11A	2.7400
C11...C9 ⁱ	3.491 (5)	C11...H10	2.6200
C12...C12 ⁱⁱ	3.5581 (16)	C12...H7	2.6400
C12...C7	3.578 (4)	H3...H11B	2.2900
C11...H12A ⁱⁱⁱ	3.0500	H3...C12 ^{viii}	3.0800
C11...H10	2.9800	H4...H12B	2.3100
C12...H12B ^{iv}	3.0400	H4...C4 ^{viii}	2.9200
C12...H3 ^v	3.0800	H7...C12	3.0900
C12...H7	3.0900	H7...C12	2.6400
C1...C4 ^{iv}	3.521 (5)	H7...H12A	2.2100
C4...C1 ^{vi}	3.521 (5)	H8...C8 ^{ix}	3.0300
C6...C9 ^{vi}	3.504 (6)	H10...C11	2.9800
C6...C12 ^{iv}	3.595 (5)	H10...C11	2.6200
C7...C12	3.578 (4)	H10...H11A	2.2300

C7...C12 ^{iv}	3.448 (6)	H11A...C3 ^{iv}	3.0100
C9...C6 ^{iv}	3.504 (6)	H11A...C10	2.7400
C9...C11 ⁱ	3.491 (5)	H11A...H10	2.2300
C10...C11	3.464 (4)	H11B...H3	2.2900
C12...C7 ^{vi}	3.448 (6)	H12A...C7	2.7200
C12...C6 ^{vi}	3.595 (5)	H12A...C7 ^{vi}	2.8400
C3...H11A ^{vi}	3.0100	H12A...C8 ^{vi}	2.9600
C4...H4 ^v	2.9200	H12A...H7	2.2100
C7...H12A	2.7200	H12A...C11 ^x	3.0500
C7...H12A ^{iv}	2.8400	H12B...C12 ^{vi}	3.0400
C8...H8 ^{vii}	3.0300	H12B...H4	2.3100
C8...H12A ^{iv}	2.9600		
C2—C1—C6	119.7 (3)	C4—C3—H3	119.00
C2—C1—C10	122.3 (3)	C3—C4—H4	119.00
C6—C1—C10	118.0 (3)	C5—C4—H4	119.00
C1—C2—C3	119.3 (3)	C6—C7—H7	120.00
C1—C2—C11	121.3 (3)	C8—C7—H7	120.00
C3—C2—C11	119.4 (4)	C7—C8—H8	119.00
C2—C3—C4	121.2 (4)	C9—C8—H8	119.00
C3—C4—C5	121.4 (4)	C8—C9—H9	120.00
C4—C5—C6	119.0 (3)	C10—C9—H9	120.00
C4—C5—C12	119.2 (4)	C1—C10—H10	119.00
C6—C5—C12	121.8 (4)	C9—C10—H10	119.00
C1—C6—C5	119.4 (4)	C11—C11—H11A	109.00
C1—C6—C7	118.3 (3)	C11—C11—H11B	109.00
C5—C6—C7	122.3 (3)	C2—C11—H11A	110.00
C6—C7—C8	120.6 (3)	C2—C11—H11B	109.00
C7—C8—C9	121.5 (4)	H11A—C11—H11B	108.00
C8—C9—C10	119.8 (4)	C12—C12—H12A	109.00
C1—C10—C9	121.8 (3)	C12—C12—H12B	109.00
C11—C11—C2	111.0 (3)	C5—C12—H12A	109.00
C12—C12—C5	112.6 (3)	C5—C12—H12B	109.00
C2—C3—H3	119.00	H12A—C12—H12B	108.00
C6—C1—C2—C3	-0.1 (5)	C2—C3—C4—C5	-0.3 (6)
C6—C1—C2—C11	-179.5 (3)	C3—C4—C5—C6	0.7 (6)
C10—C1—C2—C3	-179.8 (4)	C3—C4—C5—C12	-177.1 (4)
C10—C1—C2—C11	0.8 (6)	C4—C5—C6—C1	-0.7 (5)
C2—C1—C6—C5	0.5 (5)	C4—C5—C6—C7	-180.0 (4)
C2—C1—C6—C7	179.8 (3)	C12—C5—C6—C1	177.0 (3)
C10—C1—C6—C5	-179.8 (3)	C12—C5—C6—C7	-2.3 (5)
C10—C1—C6—C7	-0.5 (5)	C4—C5—C12—C12	-101.9 (4)
C2—C1—C10—C9	179.3 (4)	C6—C5—C12—C12	80.4 (4)
C6—C1—C10—C9	-0.4 (6)	C1—C6—C7—C8	1.0 (6)
C1—C2—C3—C4	0.0 (6)	C5—C6—C7—C8	-179.8 (4)
C11—C2—C3—C4	179.4 (4)	C6—C7—C8—C9	-0.5 (6)

C1—C2—C11—C11	75.3 (4)	C7—C8—C9—C10	-0.5 (7)
C3—C2—C11—C11	-104.1 (4)	C8—C9—C10—C1	1.0 (6)

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+3/2, z-1/2$; (iv) $x, y-1, z$; (v) $-x+1, y-1/2, -z+1/2$; (vi) $x, y+1, z$; (vii) $-x, y+1/2, -z+1/2$; (viii) $-x+1, y+1/2, -z+1/2$; (ix) $-x, y-1/2, -z+1/2$; (x) $x, -y+3/2, z+1/2$.