

Poly[(3-nitrobenzoato)(μ_3 -1,2,4-triazolato)cobalt(II)]

Xu-Liang Qi

Liaocheng Vocational and Technical College, Liaocheng 252000, ShanDong, People's Republic of China
Correspondence e-mail: q200801@sina.com


Received 30 November 2008; accepted 22 December 2008

Key indicators: single-crystal X-ray study; *T* = 296 K; mean σ (C–C) = 0.003 Å;
R factor = 0.023; *wR* factor = 0.059; data-to-parameter ratio = 15.2.

In the title compound, $[\text{Co}(\text{C}_2\text{H}_2\text{N}_3)(\text{C}_7\text{H}_4\text{NO}_4)]_n$, the Co^{II} atom is five-coordinated by three triazolate ligands and one bidentate 3-nitrobenzoate anion in a distorted trigonal-bipyramidal geometry. The triazolate ligand bridges the Co^{II} atoms, generating a two-dimensional net parallel to the *ab* plane, in which both the Co^{II} atom and the triazolate ligand act as three-connected nodes. Two weak intermolecular C–H···O hydrogen bonds connect the nets.

Related literature

For metal-triazole complexes, see: Park *et al.* (2006); Yang *et al.* (2008); Zhai *et al.* (2007). For Co–O and Co–N bond lengths, see: Zhang *et al.* (2008).



Experimental

Crystal data

$[\text{Co}(\text{C}_2\text{H}_2\text{N}_3)(\text{C}_7\text{H}_4\text{NO}_4)]$

M_r = 293.11

Data collection

Bruker SMART 1K CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2004)
 R_{int} = 0.029
 T_{\min} = 0.802, T_{\max} = 0.826

19233 measured reflections
2477 independent reflections
2245 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)]$ = 0.023
 $wR(F^2)$ = 0.059
 S = 1.04
2477 reflections

163 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.28 \text{ e } \text{\AA}^{-3}$

Table 1
Selected bond lengths (Å).

Co1–O1	2.3314 (12)	Co1–N2 ⁱ	2.0118 (12)
Co1–O2	2.0008 (12)	Co1–N3 ⁱⁱ	2.0385 (12)
Co1–N1	2.0232 (12)		

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, -z + 1$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, z$.

Table 2
Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
C3–H3···O2 ⁱⁱⁱ	0.93	2.54	3.250 (3)	134
C8–H8···O4 ^{iv}	0.93	2.46	3.372 (2)	169

Symmetry codes: (iii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + \frac{1}{2}, -y + 1, z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2370).

References

- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Park, H., Moureau, D. M. & Parise, J. B. (2006). *Chem. Mater.* **18**, 525–531.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yang, E.-C., Liu, Z.-Y., Wang, X.-G., Batten, S. R. & Zhao, X.-J. (2008). *CrystEngComm*, **10**, 1140–1143.
- Zhai, Q.-G., Lu, C.-Z., Wu, X.-Y. & Batten, S. R. (2007). *Cryst. Growth Des.* **7**, 2332–2342.
- Zhang, J., Chew, E., Chen, S.-M., Pham, J. T. H. & Bu, X.-H. (2008). *Inorg. Chem.* **47**, 3495–3497.

supporting information

Acta Cryst. (2009). E65, m134 [doi:10.1107/S1600536808043596]

Poly[(3-nitrobenzoato)(μ_3 -1,2,4-triazolato)cobalt(II)]

Xu-Liang Qi

S1. Comment

Recently, more and more attention is paid on the coordination chemistry about trz ligand or analogy ligand (Park *et al.*, 2006; Yang *et al.*, 2008; Zhai *et al.*, 2007), driven by their intriguing topological matrix and potential applications.

The asymmetric unit of I is shown in Fig. 1. The Co^{II} atom is five-coordinated by two *L* (3-nitrobenzoate anion) O atoms, three trz N atoms to give rise to a distorted trigonal-bipyramidal geometry. The Co—O/N bond lengths of 2.0008 (12)–2.3314 (12) Å (Table 1) are in the normal range (Zhang *et al.*, 2008). The trz and *L* ligand adopt bridging and bidentate coordinated modes, respectively. As shown in Fig. 2a, the Co^{II} atoms are combined together by trz ligands to generate a two-dimensional net parallel to the *ab* plane with the *L* ligands ligated on the two-dimensional net up and down. From a topological point of view, if considering the trz ligands and cobalt ions as three-connected nodes.


Moreover, besides the presence of two weak intermolecular C—H···O hydrogen bonds, see Table 2, there is not other obvious supramolecular interactions between two-dimensional nets,

S2. Experimental


CoCl₂ (1.0 mmol), 3-nitrobenzoic acid (1 mmol) and triazole (1 mmol) were dissolved in water (10 ml). The solution was heated in a 25 ml Teflonlined reaction vessel at 433 K for *ca* 3 days and then cooled to room temperature. Purple crystals of the title compound were obtained in a yield of 78%.

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

An *ORTEP* view of the asymmetric unit with 50% thermal ellipsoids for non-H atoms [symmetry codes: (A) $-x + 1/2, y - 1/2, z$; (B) $x + 1/2, -y + 3/2, -z + 1$].

**Figure 2**

a) View of the two-dimensional net onto the ab plane, formed by cobalt ions and trz ligands; b) View of the two-dimensional net built on three-connected trz and cobalt nodes.

Poly[(3-nitrobenzoato)(μ_3 -1,2,4-triazolato)cobalt(II)]

Crystal data

$[\text{Co}(\text{C}_2\text{H}_2\text{N}_3)(\text{C}_7\text{H}_4\text{NO}_4)]$

$M_r = 293.11$

Orthorhombic, $Pbca$

Hall symbol: $-P\bar{2}ac\bar{2}ab$

$a = 9.2419 (18)$ Å

$b = 10.377 (2)$ Å

$c = 22.597 (5)$ Å

$V = 2167.1 (8)$ Å³

$Z = 8$

$F(000) = 1176$

$D_x = 1.797$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 15896 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 1.60 \text{ mm}^{-1}$
 $T = 296 \text{ K}$

Block, purple
 $0.14 \times 0.12 \times 0.12 \text{ mm}$

Data collection

Bruker SMART 1K CCD area-detector
dифрактометр
Radiation source: sealed tube
Graphite monochromator
Detector resolution: 8.192 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 2004)
 $T_{\min} = 0.802$, $T_{\max} = 0.826$

19233 measured reflections
2477 independent reflections
2245 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.1^\circ$
 $h = -12 \rightarrow 11$
 $k = -13 \rightarrow 12$
 $l = -29 \rightarrow 29$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.059$
 $S = 1.04$
2477 reflections
163 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0278P)^2 + 1.1998P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0918 (2)	0.62096 (17)	0.17212 (8)	0.0358 (4)
C2	-0.0058 (2)	0.7078 (2)	0.14920 (8)	0.0470 (5)
H2	-0.0198	0.7141	0.1085	0.056*
C3	-0.0823 (3)	0.7853 (2)	0.18746 (9)	0.0522 (6)
H3	-0.1473	0.8457	0.1728	0.063*
C4	-0.0620 (2)	0.77306 (18)	0.24798 (8)	0.0406 (4)
H4	-0.1157	0.8240	0.2738	0.049*
C5	0.1165 (2)	0.60861 (16)	0.23241 (7)	0.0322 (4)
H5	0.1843	0.5503	0.2468	0.039*
C6	0.03721 (19)	0.68583 (16)	0.27043 (7)	0.0295 (3)
C7	0.05354 (18)	0.67429 (16)	0.33614 (7)	0.0290 (3)
C8	0.32376 (16)	0.75394 (14)	0.51070 (7)	0.0253 (3)
H8	0.3425	0.6826	0.5344	0.030*
C9	0.22994 (17)	0.87843 (13)	0.44808 (6)	0.0231 (3)
H9	0.1693	0.9118	0.4190	0.028*
Co1	0.08538 (2)	0.615646 (17)	0.442261 (8)	0.01829 (7)
N1	0.21652 (13)	0.75890 (11)	0.47077 (5)	0.0227 (2)
N2	0.39953 (13)	0.86141 (12)	0.51260 (5)	0.0225 (3)
N3	0.33828 (13)	0.94272 (11)	0.47154 (5)	0.0209 (2)
N4	0.1706 (2)	0.53740 (17)	0.13114 (7)	0.0459 (4)
O1	-0.00457 (13)	0.75383 (12)	0.36970 (5)	0.0367 (3)
O2	0.12664 (15)	0.58069 (12)	0.35684 (5)	0.0351 (3)

O3	0.2746 (2)	0.47839 (18)	0.14938 (7)	0.0683 (5)
O4	0.1257 (2)	0.53020 (16)	0.08006 (6)	0.0627 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0517 (11)	0.0329 (9)	0.0228 (8)	-0.0083 (7)	0.0032 (7)	-0.0016 (6)
C2	0.0745 (14)	0.0453 (11)	0.0211 (8)	-0.0027 (10)	-0.0091 (8)	0.0050 (8)
C3	0.0778 (16)	0.0444 (12)	0.0345 (10)	0.0148 (10)	-0.0150 (9)	0.0058 (9)
C4	0.0592 (11)	0.0331 (9)	0.0295 (9)	0.0079 (8)	-0.0040 (8)	0.0000 (7)
C5	0.0422 (9)	0.0299 (8)	0.0246 (8)	-0.0020 (7)	-0.0011 (7)	0.0013 (6)
C6	0.0418 (9)	0.0261 (8)	0.0208 (7)	-0.0037 (7)	-0.0026 (6)	0.0022 (6)
C7	0.0371 (8)	0.0284 (8)	0.0214 (7)	-0.0058 (6)	-0.0020 (6)	0.0014 (6)
C8	0.0290 (7)	0.0196 (7)	0.0274 (7)	-0.0033 (6)	-0.0042 (6)	0.0061 (6)
C9	0.0269 (7)	0.0190 (7)	0.0234 (7)	-0.0015 (5)	-0.0031 (5)	0.0034 (5)
Co1	0.02203 (12)	0.01445 (11)	0.01838 (11)	-0.00039 (6)	0.00086 (7)	0.00145 (7)
N1	0.0263 (6)	0.0181 (6)	0.0238 (6)	-0.0032 (5)	-0.0026 (5)	0.0025 (5)
N2	0.0247 (6)	0.0185 (6)	0.0243 (6)	-0.0015 (5)	-0.0038 (5)	0.0050 (5)
N3	0.0246 (6)	0.0163 (6)	0.0217 (6)	-0.0005 (4)	-0.0008 (4)	0.0039 (5)
N4	0.0611 (11)	0.0437 (9)	0.0330 (8)	-0.0104 (8)	0.0139 (7)	-0.0058 (7)
O1	0.0453 (7)	0.0413 (7)	0.0234 (6)	0.0066 (6)	-0.0001 (5)	-0.0033 (5)
O2	0.0541 (7)	0.0304 (6)	0.0207 (5)	0.0052 (6)	-0.0027 (5)	0.0020 (5)
O3	0.0667 (11)	0.0791 (12)	0.0590 (10)	0.0140 (10)	0.0135 (8)	-0.0168 (9)
O4	0.1046 (13)	0.0585 (10)	0.0249 (7)	-0.0100 (9)	0.0117 (8)	-0.0095 (7)

Geometric parameters (\AA , ^\circ)

C1—C2	1.376 (3)	C8—N2	1.3175 (19)
C1—C5	1.388 (2)	C8—N1	1.3414 (19)
C1—N4	1.463 (2)	C8—H8	0.9300
C2—C3	1.376 (3)	C9—N3	1.3149 (19)
C2—H2	0.9300	C9—N1	1.3478 (18)
C3—C4	1.386 (3)	C9—H9	0.9300
C3—H3	0.9300	Co1—O1	2.3314 (12)
C4—C6	1.385 (2)	Co1—O2	2.0008 (12)
C4—H4	0.9300	Co1—N1	2.0232 (12)
C5—C6	1.385 (2)	Co1—N2 ⁱ	2.0118 (12)
C5—H5	0.9300	Co1—N3 ⁱⁱ	2.0385 (12)
C6—C7	1.497 (2)	N2—N3	1.3759 (16)
C7—O1	1.243 (2)	N4—O3	1.212 (2)
C7—O2	1.272 (2)	N4—O4	1.229 (2)
C2—C1—C5	122.56 (17)	N1—C9—H9	123.7
C2—C1—N4	118.45 (17)	O2—Co1—N2 ⁱ	132.33 (5)
C5—C1—N4	118.98 (17)	O2—Co1—N1	109.04 (5)
C1—C2—C3	118.87 (17)	N2 ⁱ —Co1—N1	105.25 (5)
C1—C2—H2	120.6	O2—Co1—N3 ⁱⁱ	95.03 (5)
C3—C2—H2	120.6	N2 ⁱ —Co1—N3 ⁱⁱ	103.60 (5)

C2—C3—C4	119.79 (19)	N1—Co1—N3 ⁱⁱ	109.64 (5)
C2—C3—H3	120.1	O2—Co1—O1	60.06 (5)
C4—C3—H3	120.1	N2 ⁱ —Co1—O1	88.82 (5)
C6—C4—C3	120.72 (18)	N1—Co1—O1	89.18 (5)
C6—C4—H4	119.6	N3 ⁱⁱ —Co1—O1	153.26 (5)
C3—C4—H4	119.6	C8—N1—C9	102.90 (12)
C6—C5—C1	117.93 (16)	C8—N1—Co1	128.95 (10)
C6—C5—H5	121.0	C9—N1—Co1	127.60 (10)
C1—C5—H5	121.0	C8—N2—N3	106.17 (12)
C4—C6—C5	120.10 (15)	C8—N2—Co1 ⁱⁱⁱ	124.78 (10)
C4—C6—C7	118.83 (15)	N3—N2—Co1 ⁱⁱⁱ	128.36 (9)
C5—C6—C7	121.04 (15)	C9—N3—N2	105.91 (11)
O1—C7—O2	120.80 (14)	C9—N3—Co1 ^{iv}	125.40 (10)
O1—C7—C6	120.56 (15)	N2—N3—Co1 ^{iv}	128.05 (9)
O2—C7—C6	118.63 (15)	O3—N4—O4	123.81 (19)
N2—C8—N1	112.47 (13)	O3—N4—C1	118.64 (17)
N2—C8—H8	123.8	O4—N4—C1	117.55 (19)
N1—C8—H8	123.8	C7—O1—Co1	82.34 (10)
N3—C9—N1	112.56 (13)	C7—O2—Co1	96.61 (10)
N3—C9—H9	123.7		
C5—C1—C2—C3	-0.2 (3)	C7—Co1—N1—C9	25.56 (14)
N4—C1—C2—C3	178.90 (19)	N1—C8—N2—N3	0.24 (17)
C1—C2—C3—C4	-1.2 (3)	N1—C8—N2—Co1 ⁱⁱⁱ	-170.88 (10)
C2—C3—C4—C6	1.7 (3)	N1—C9—N3—N2	-0.56 (17)
C2—C1—C5—C6	1.2 (3)	N1—C9—N3—Co1 ^{iv}	170.84 (10)
N4—C1—C5—C6	-177.94 (16)	C8—N2—N3—C9	0.19 (16)
C3—C4—C6—C5	-0.7 (3)	Co1 ⁱⁱⁱ —N2—N3—C9	170.89 (10)
C3—C4—C6—C7	-178.94 (19)	C8—N2—N3—Co1 ^{iv}	-170.91 (10)
C1—C5—C6—C4	-0.7 (3)	Co1 ⁱⁱⁱ —N2—N3—Co1 ^{iv}	-0.21 (18)
C1—C5—C6—C7	177.51 (16)	C2—C1—N4—O3	166.47 (19)
C4—C6—C7—O1	-10.4 (2)	C5—C1—N4—O3	-14.4 (3)
C5—C6—C7—O1	171.35 (16)	C2—C1—N4—O4	-14.4 (3)
C4—C6—C7—O2	168.80 (16)	C5—C1—N4—O4	164.78 (17)
C5—C6—C7—O2	-9.4 (2)	O2—C7—O1—Co1	-4.03 (15)
N2—C8—N1—C9	-0.55 (17)	C6—C7—O1—Co1	175.16 (15)
N2—C8—N1—Co1	171.29 (10)	O2—Co1—O1—C7	2.54 (10)
N3—C9—N1—C8	0.68 (17)	N2 ⁱ —Co1—O1—C7	-139.56 (10)
N3—C9—N1—Co1	-171.31 (10)	N1—Co1—O1—C7	115.17 (10)
O2—Co1—N1—C8	-113.76 (13)	N3 ⁱⁱ —Co1—O1—C7	-20.77 (16)
N2 ⁱ —Co1—N1—C8	99.90 (13)	O1—C7—O2—Co1	4.68 (18)
N3 ⁱⁱ —Co1—N1—C8	-10.96 (14)	C6—C7—O2—Co1	-174.52 (13)
O1—Co1—N1—C8	-171.55 (13)	N2 ⁱ —Co1—O2—C7	53.70 (13)
O2—Co1—N1—C9	56.19 (14)	N1—Co1—O2—C7	-79.98 (11)
N2 ⁱ —Co1—N1—C9	-90.15 (13)	N3 ⁱⁱ —Co1—O2—C7	167.23 (10)

N3 ⁱⁱ —Co1—N1—C9	158.99 (12)	O1—Co1—O2—C7	−2.47 (9)
O1—Co1—N1—C9	−1.60 (13)		

Symmetry codes: (i) $x-1/2, -y+3/2, -z+1$; (ii) $-x+1/2, y-1/2, z$; (iii) $x+1/2, -y+3/2, -z+1$; (iv) $-x+1/2, y+1/2, z$.

Hydrogen-bond geometry (\AA , °)

D—H···A	D—H	H···A	D···A	D—H···A
C3—H3···O2 ^v	0.93	2.54	3.250 (3)	134
C8—H8···O4 ^{vi}	0.93	2.46	3.372 (2)	169

Symmetry codes: (v) $-x, y+1/2, -z+1/2$; (vi) $-x+1/2, -y+1, z+1/2$.